

# Effect of Industrial Pollutants on Physical Efficiency of Post-Menopausal Women in Urban Area

Dipak Kumar Bhuniya<sup>1</sup>, Sudipta Bhuniya<sup>2</sup>

Department of Physiology, Midnapore College, Mindnapore-721 101, West Bengal, India

Corresponding Author

Dr. Dipak Kumar Bhuniya Associate Professor, Department of Physiology, Midnapore College  
Mindnapore-721 101, West Bengal, India, Mob: +91-8016441997, E-mail: drdkbhuniya1960@gmail.com

**Abstract:** Present study evaluated the relationship of physical fitness, aerobic capacity and heart responses of Postmenopausal women (n=100) living in rural and urban area. Body fat percent was estimated by Siri equation putting the value of iliac skin fold thickness and arm skinfold thickness using Harpenden Skin fold Calipers.  $VO_2$  max was determined by treadmill exercise with direct measurement of oxygen consumption. Blood pressure was measured by Sphygmomanometer and the radial pulse rate (Beats/Min) was measured. Anthropometric data (Height, body mass) was taken by using Anthropometer rod, weighing machine and data were compared among the subjects of both groups. Blood Haemoglobin of both groups was measured by spectrophotometrical technique and compared. Rural post-menopausal women had significantly ( $p<0.05$ ) posses higher lean mass, physical efficiency and aerobic capacity than urban menopausal women. Urban menopausal women showed a significantly ( $p<0.05$ ) lower cardio respiratory responses to exercise than women living in rural area. Insignificant ( $p>0.05$ ) differences of hemoglobin concentration and body weight of rural women was noticed in respect to post-menopausal urban women. Good nutrients have stimulatory where as Industrial pollutants have some detrimental roles on muscle strength and physical efficiency in either sex of human beings in all age group.


**Keywords:** anthropometric, ergometry, nutrient, pollutant, post menopausal, haemoglobin

## 1. Introduction

Health is nothing but a state of complete physical, mental developed social wellbeings and not merely the absence of diseases or infirmity (WHO,1996). Good health causes better efficiency to combat the pollution tide in world. Health mainly depends on food nutrients. Deficiency of nutrient creates ill health. Besides that mental pleasure is also an important factor for health condition. On the other hand anxiety, stress, manmade pollutants affect the health condition of people living particularly in urban area affected mainly by vehicle exhaust, unburned hydrocarbons, industrial wastes, municipal sewage, nuclear waste and water pollutants which coastally increases the concentration of Lead, Chromium, Mercury, Arsenic, Fluoride, Silicon and other toxic metals in surrounding areas (Thawley et al., 1977; Pocock et al., 1989; Leggett, 1993; WHO, 1996). Particularly Lead affects hemoglobin synthesis by inhibiting the  $\delta$ -Amino Leuvilinic Acid Synthase enzyme and creates anaemic condition. It causes lung disorder and arrest foetal development (Granick et al., 1973; Klein et al., 1980; Lakatta, 1993). Alongside it causes fatigue condition in urban people. The metallic contaminant destroys soil bacteria and other organisms acting on water to impure it, resulting death of animals and plants. Arsenic and Lead particularly make the water undrinkable and also their toxicity damage the chromosome and process of heredity. It is mentioned that 60  $\mu$ g of Arsenic and 40  $\mu$ g Lead/100 mL of blood cause of damaging brain cells (National Academy of Sciences/Institute of Medicine, 2003). Slight increase in mercury level can prevent the movement of Gill filament and there by arrest respiration in aquatic animals. Other metals like Cobalt, Zn, Ni, Cu also cause different health

hazards including Gastrointestinal disorders (Suttle and Mills,1966; Thawley et al., 1977). High concentration of CFCs and carcinogenic chemicals also affect health condition. So, to protect our health condition preventive measure should be adopted immediately. Other factors like Age, Sex, regular physical exercise may affect the body efficiency (Londeree and Moeschberger,1982). Relationship between health and efficiency with some controlling factors is shown by following schematic diagram:

**Scheme 1:**



Health is related with body fitness in terms of physical efficiency and it may be expressed either Gross efficiency or Net efficiency.

Gross Efficiency (%) = (External work output/ total energy used for work) X 100

Net Efficiency (%) = (External work output/ total energy used – Basal requirement) X 100

Mechanical Efficiency (%) = (Actual Mechanical work /input of energy) X 100

So, to combat the environmental hazards sufficient and good nutrients should be included in their daily diet.

For taking adequate calorific valued food good nutrient everyone has some idea about the computation of calorie requirement as well as nutrient requirement of an individual and it can be computed by the following ways:

**Table 1:** Computation of Calorie requirement

If the individual's age -50 years, body weight- 50 kg, BMR- 60 Kcal/Hr.			
Activities	Duration	@KCal	Energy (KCal)
Sleep	8 Hrs.	90% of BMR	= 60 X 0.9 X 8
<b>Non Occupational activities:</b>			
Washing, bathing, Dressing, Eating	1.5 Hrs	2/Kg bw/hour	50 X 1.5 X 2
Sitting, Standing	3 Hrs	1.4/Kg	50 X 3 X 1.4
Writing, Card Playing, Gossiping,	2 Hrs	1.5/Kg bw	50 X 2 X 1.5
Walking, Running	1.5 Hrs	3/Kg bw/hour	50 X 1.5 X 3
<b>Occupational activities:</b>			
Sedentary/Moderate/Heavy	8 Hrs.	4/Kg bw/hour	50 X 4 X 8
<b>Total = 2767</b>			
NUTRIENT REQUIREMENT			
Nutrient	Rate	Amount	KCal
Protein	1 gm/Kg/Day	50 gm	205
Fat	15% of total calorie	49 gm	456
Carbohydrate	(Total calories– calories for Fat & Protein) / 4.1	583gm	2390

## 2. Materials and Methods

For present study hundred (100) rural women from different villages of Midnapore districts and hundred (100) urban women from Howrah district (Industrial belt) were selected. Some physiological parameters were taken on those subjects participating in exercise including Harvard Step test and Trade mill test. The subjects were randomly chosen and belong to menopausal and post-menopausal group (Age between 45 -54 years).

Physiological parameters of such women were noted from anthropometric measurement i.e. height and weight using Standard laboratory techniques (Maiti et al., 2011). Arm circumference, Chest circumference during inspiration and expiration were also measured by standard laboratory methods (Singh and Bishnoi, 2005). Skin fold thickness of subjects of both group were measured by Harpenden Skin fold Calipers (Jason and David, 2010). And the experimental data collected from rural and urban women group has been shown in **Table 2**.

Blood pressure (systolic and diastolic) of rural women and urban women were measured by sphygmomanometer and radial pulse rate of both groups were also recorded. Measured data has been shown in **Table 3**.

Resting and exercised heart rate of both groups were measured from carotid pulsation, maximal oxygen uptake of both groups was measured during continuous Treadmill exercise, Physical Fitness Index (%) of the subjects of both group was measured by Harvard step test, blood Haemoglobin concentration of both rural women and urban women was determined by Spectrophotometrical technique

and measured data of the above experiments has been shown in **Table 4**.

## 3. Results and Discussion

The present study shows for the first time to our knowledge that physical efficiency of post-menopausal Women decreased with advancing age. As lower concentration of oestrogen may affects muscular calcium metabolism that also may cause of decrement of physical efficiency in women (Aloia et al., 1991).

Our present investigation on post-menopausal women living in Urban and rural area provides an impressive evidence that muscular efficiency can be restored by physical exercise ignoring lower concentration of sex hormones and food nutrients (Pocock et al., 1989).

Good nutrients have enormous role on production of sufficient energy for muscular contraction in this Physiological State of women (WHO, 1996).

But higher concentration of heavy metals develop toxicity that decreases oxygenation of haemoglobin in blood and aerobic respiration of tissues of human body that also one of the reason of lower physical efficiency of women living in urban area (National academy of Sciences/Institute of Medicine, 2003).

Interesting findings in our investigation are that the measured data of body mass, Arm circumference, iliac, chest circumference during expiration higher in menopausal women living in rural area in comparison to that data

collected from women living in urban area. But body fat percent and skin fold thickness of post-menopausal women of urban zone is higher than that of post-menopausal women in rural area (**Table-2**).

Another finding was observed that myocardial strength, maximal oxygen uptake or vital capacity and physical fitness index of rural menopausal women is higher than that of urban menopausal women (Landers and Petruzzewllo, 1994). Blood haemoglobin concentration of both rural women and urban women were measured but shows insignificant ( $P>0.05$ ) variation (**Table-3, 4**). Regular physical exercise increases oxygen uptake capacity (Dalsky et al., 1988), oxygenation of myoglobin and also increases oxygen consumption of muscle (Leggett, 1993; Landers and Petruzzewllo, 1994). In spite of regular consumption of sufficient quality nutrients postmenopausal women living in Industrial belt shows lower physical efficiency due to greater concentration of environmental pollutants (Leggett, 1993). So, it may be concluded that the aged women are able to increase their heart rate and oxygen uptake to its maximum level like other women belonging to lower age groups, if they involved in regular physical exercise like yoga, Meditation in a suitable environmental condition. Rise of concentration of environmental pollutants may one of the major causes of muscular fatigue of people living in urban area.

**Table 2:** Anthropometric measurements of rural women and urban women. In each vertical column data are expressed as Mean  $\pm$  SEM

Variables	Rural women (n=100)	Urban Women(n=100)
Age (years)	51.21 $\pm$ 1.98	50.27 $\pm$ 0.63
Height (cm)	151.65 $\pm$ 1.17	145.60 $\pm$ 0.89
Weight (Kg)	52.25 $\pm$ 1.94	50.27 $\pm$ 0.63
Arm Circumference(cm)	24.62 $\pm$ 0.60	21.89 $\pm$ 0.39
Head Circumference (cm)	52.09 $\pm$ 0.94	52.24 $\pm$ 0.30
Chest Circumference (cm)		
Inspiration	80 $\pm$ 1.02	74.22 $\pm$ 2.21
Expiration	74.25 $\pm$ 0.88	72.86 $\pm$ 0.78
Total Skin fold Thickness (mm)		
Iliac	9.0 $\pm$ 0.02	8.6 $\pm$ 0.28
Arm	5.0 $\pm$ 0.05	8.0 $\pm$ 0.02
% of Body Fat	16.9 $\pm$ 0.01	18.4 $\pm$ 0.03
Body density = 1.0764-(0.3081) X iliac Skin fold -0.3088 X Arm Skin Fold		
Percent of body fat = {(4.570/BD)-4.142} X 100		
Total Fat (Kg)= (Body weight (Kg) X Percent of fat)/ 100		

**Table 3:** Heart response of Rural & Urban women. In vertical columns data are expressed as Mean  $\pm$  SEM

Variables		Rural Women (n=100)	Urban Women (n=100)	Level of Significance
Blood Pressure (mm of Hg)	Systolic	138.0 $\pm$ 2.31	140.3 $\pm$ 3.36	p>0.05
	Diastolic	84.92 $\pm$ 2.30	80.12 $\pm$ 1.99	p>0.05
Pulse Rate(beats/min.)		67.0 $\pm$ 3.10	72.26 $\pm$ 2.39	p>0.05

**Table 4:** Cardio-respiratory response after exercise and blood Haemoglobin concentration. In vertical columns data are expressed as Mean  $\pm$  SEM.

Variables	Rural Women(n=100)	Urban Women (n=100)	Level of Significance
Resting Heart Rate (beats/min)	67.28 $\pm$ 2.64	72.26 $\pm$ 1.41	p>0.05
Maximum Heart Rate (beats/min)	146.0 $\pm$ 1.04	166.0 $\pm$ 2.41	P<0.05
Maximal Oxygen uptake (lit/min)	2.84 $\pm$ 0.08	1.52 $\pm$ 0.02	p<0.05
Physical Fitness Index (%)	68.0 $\pm$ 0.02	54.0 $\pm$ 0.32	P<0.05
Hemoglobin Concentration (gm%)	9.2 $\pm$ 0.84	10.8 $\pm$ 1.02	p>0.05

## References

- [1] Aloia, J.F., DMcGown, M., Vaswani, A.N., Ross, P and Cohen, S.H. (1991). Relationship of menopause to skeletal and muscle mass. American journal of Clinical Nutrition, 53:1378—1383.
- [2] Dalsky, G., Stocke, K.S., Eshani, A.A., Slatopolsky, E., and Lee, W.C. (1988). Weight bearing exercise training and lumbar bone mineral content in post-menopausal women. Annals of Internal Medicine, 108:824-828.
- [3] Doblen, W.V. (1959). Anthropometric determination of fat free body weight. Acta Medical Scientific Journal, 37:165.
- [4] Granick, J.L., Sassa, S., Granick R.D. and Kappas, A. (1973). Studies in Lead poisoning ii: Correlation between the ratio of activated to inactivated d-aminolevulinic acid dehydrates of whole blood and the blood lead level. Biochemia Medica, 8:149-159.
- [5] Jason, R. B. and DAVID J.S (2010). Validity of 2 skinfold calipers in estimating percent body fat of college-aged men and women. Journal of Strength and Conditioning Research. 24:3448–3456.
- [6] Klein, J.P., Forester, H.V., Stewart, R.D. and Wu, A. (1980). Haemoglobin affinity for oxygen during short term exhaustive exercise. Journal of Applied Physiology, 48:236.
- [7] Lakatta, E.G. (1993). Cardiovascular regulatory mechanism in advanced age. Physiological Review. 73:413.
- [8] Landers, D.M.; Petruzzello, S.J. In: Physical activity, fitness, and anxiety. Bouchard, C., Shephard, Roy J. and Stephens, T (Eds). (1994). Physical activity, fitness, and health: International proceedings and consensus statement. , pp. 868-882.
- [9] Leggett, R.W. (1993). An age specific Kinetic model for Lead metabolism in human health perspective. 101:593-616.
- [10] Londeree, B.R. and Moeschberger, M.L. (1982). Effect of age and other factors on maximal heart rate. Research Quarterly for Exercise & Sport. 53:297.
- [11] Maier, W.F. and Costa, L.G. (1990). Toxicology. 51:175-188.
- [12] Maiti, S., De, D., Chatterjee, K., Jana, K., Ghosh, D. and Paul, S.P. Prevalence of stunting and thinness among early adolescent school girls of Paschim Medinipur district, West Bengal. International Journal of Biological and Medical Research. 2011; 2(3): 781-783.

- [13] National academy of Sciences/Institute of Medicine. (2003). Dietary reference intakes for vitamin -A, vit-K,As,Cr,Cu,I,Fe,Mn,Mo.,Ni.,Zn. Food and Nutrition Board, Institute of Medicine, Washington, DC.
- [14] Pocock, N., Eisman, J., Gwinn,T., Sambrook, P., Kelly, P., Freund , J. and Yeates M. (1989). Muscle strength, physical fitness, and weight but not age predict femoral neck bone mass. *Journal of Bone Mineral Research*. 4:441-448.
- [15] Singh, S. and Bishnoi, I.(2005).Trend of growth in Mid-arm circumference and head circumference of pre-schol female children of Faizabaad District, UP. *Indian Journal of Preventive and Social Medicine*. 36:143-146.
- [16] Suttle, N.F and C.F.Mills. (1966). Studies of the toxicity of copper to pigs: Effects of Zinc and iron salts on development of copper toxicosis. *Biological Journal of Nutrition*.20:135-149.
- [17] Thawley, D.G., Prattand S.E and Selby, L.A. (1977). Antagonistic effect of Zinc on increased urinary  $\delta$ -aminolevulinic acid excretion in lead intoxicated rats. *Environmental Research*. 14:463-475.
- [18] Torra, M. and Rodailans, M.(1989). *Toxicology*.42:467.
- [19] WHO (World Health Organisation). (1996). Trace elements in human health and nutrition, chapter10: Manganese.Geneva, Pp. 163-167.
- [20] Wittenberg, B.A., Wittenberg, J.B. and Caldwell, P.R. (1975). Role of Myoglobin in the Oxygen Supply to Red Skeletal Muscle. *The Journal of Biological Chemistry*.250:9038-9043.