

Evaluations of Two Isolated *Paecilomyces* Against *Phthorimaea operculella* (Lepidoptera:Gelechiidae) Under Laboratory and Field Conditions

Sabbour M. M.¹, Singer, S. M.²

¹Department, Pests & Plant Protection; Agricultural, Biological Research Division, National Research Centre, El-Tahrir St. - Dokki, Cairo,

²Department of vegetable crops research, Biological Research Division, National Research Centre, El-Tahrir St. - Dokki, Cairo,

Abstract: Under laboratory conditions, the LC₅₀ recorded 122X10⁴ and 106X10⁴ conidia/ml for the fungi, *Paecilomyces fumosoroseus* and *Paecilomyces lilaceus*, respectively. Under semi field conditions the corresponding LC₅₀ 136 X10⁴ and 119 X10⁴ conidia/ml. *fumosoroseus* caused a significant decrease in infestation reached to 21± 9.45 and 20± 8.12 after 90 days of the post applications as compared to 89±13.03 and 79 ± 11.12 individuals in the control during season 2012 and 2013, respectively. The weight of the potato recorded during 2012 were, 5879± 8.28 and 5667± 7.18 kg/ feddan in the plots treated with *P. lilaceus* and *P. fumosoroseus* respectively as compared to 3398± 4.29 kg/ feddan in the control during season 2012. During season 2013 the highest weight scored 5992± 8.21kg/feddan after treatment with *P. lilaceus*.

Keywords: *Paecilomyces lilacinus*, *Paecilomyces fumosoroseus*, *Phthorimaea operculella*

1. Introduction

Potato tuber moth *Phthorimaea operculella* PTM attacks solanaceous crops with potato being favored. Foliar injury is due to the larvae (tubeworm) mining into leaflets, causing them to form transparent blisters, then move into stem tissue causing death. Tubers are marred when larvae reach tubers by two major means. Upon hatching from eggs laid on leaves, larvae can drop to the ground and burrow through cracks in the soil to a tuber, entering it through the eye. This is common after vine desiccation. Another common way is that the female PTM lays its eggs directly on exposed tubers at or near the eye. When the larvae hatch, they just enter the tuber through the eye making a slender tunnel along the surface or deep into the tuber (pictured). A tunnel can be detected by mounds of worm excrement (frass) appearing black at the entrance (pictured). Tunnels do not heal and are entryways for diseases most notably soft rot and dry rot. In Egypt it is one of the most economic important pests. Larvae cause severe damage to vegetable crops of family Solanaceae (Sarhan 2004; Soliman et al., 2008; Abul-Nasr et al., 1971 & 1972; Sabbour, 2002). The entomopathogenic *P. carneus* is found on a wide range of material, and especially in soil. It is sometimes isolated from insects, though it appears to be a weak insect pathogen. Some isolates produce several metabolites of the antibiotic group cephalosporins. *P. farinosus* is also commonly isolated from soil. It is a well-known insect pathogen, and there has been interest in its use as an agent of biological control. Sabbour and Abdel-Rahman, 2013, Sabbour et al, 2011, Sabbour, 2002, Sahab et al, 2014 control the corn porwers by different entomopathogenic fungi under laboratory and field conditions. Entomopathogenic fungi are found worldwide associated to insects and phytophagous mite populations, contributing to biological control of these arthropods on several economically important crops (Sabbour and Sahab, 2007). Commercial products have been developed with entomopathogenic fungi (Alves and Pereira, 1998). Quintela and McCoy (1998) reported that fungal

concentrations of 10⁶ and 10⁷ conidia/ml of *B. bassiana* and *N. rileyi* affected the larval development, movement and mobility of corn borers larvae during the seedlings and vegetative stages of corn plant under laboratory; greenhouse and field conditions. Success of a pest control program using *B. bassiana* however depends on conidia survival in the field environment (Benz, 1987). Conidia survival may be affected either by environmental factors (Furlong and Pell, 1997) or chemical products used to protect plants (Anderson and Roberts, 1983). Abdel-Rahman, et al. (2006) controlled the cereal aphids with the fungus *B. bassiana* and found that the infestation was reduced after fungal applications under laboratory and field conditions.

The present study aims to evaluate the pathogenicity of the two soil isolates of entomopathogenic fungus, *Paecilomyces lilacinus* and *Paecilomyces fumosoroseus* against potato tuber moth.

2. Materials and Methods

2.1 Tested Insects

Standard laboratory colony of the potato tuber moth *P. operculella* was reared on potato tubers *Solanum tuberosum* as a natural host plant under controlled conditions (26±2°C and 70±5% R.H). Eggs were obtained from the stock culture and kept in Petri-dishes till larval hatch. The rearing technique by El-Sherif (1966) was adopted. Pupae were individually kept in specimen tubes (1×3cm) till adult emergence. Adult moth were kept in oviposition cages that consist of chimney glass (8cm in diameter and 16cm height), the lower rim of which rested on the bottom of a Petri-dish lined with a disk of filter paper (Watman) and the upper rim covered with muslin. Each cage was provided with a small piece of cotton soaked in 5% honey solution as food supply. The deposited eggs were collected and kept in Petri-dishes till larval hatching. Groups of newly hatched

larvae were transferred into Petri-dishes containing fresh pieces of potato. Larval development was allowed to continue until the adult emergence

2.2 Entomopathogenic Fungi

The fungus, *Paecilomyces lilacinus*, and *Paecilomyces fumosoroseus* isolated from the Egyptian soil from Ismailia governorate. They were reproduced on potato dextrose agar (PDA) plus 0.4% yeast extracts (PDAY) and poured onto sterilized Petri-dishes (Alves *et al.*, 1998). Plating was performed according to the full dish method. The conidia were transferred from the Eppendorf vial to dish containing medium by platinum loop and then streaked. Plates were incubated at 25°C with 12 hours photo phase for fungus growth and sporulation. After ten days, conidia were scraped and transferred to conical flasks (200 ml) containing 200 ml sterilized distilled water with 0.02% the speeder sticker (tween, 80). Conidial concentrations in the suspensions were quantified directly under the optical microscope with a haemocytometer. Then the suspensions were standardized until the direct concentration 1×10^7 conidia/ml was obtained.

2.3 Efficacy of Entomopathogenic Fungi against Pests Larvae

Spores of the entomopathogenic fungi; *Paecilomyces lilacinus* and *Paecilomyces fumosoroseus*, collected from the surface of mycelium growth and spore suspensions with 2 drops of tween 80 were prepared and adjusted at 1×10^7 conidia/ml. Conidial viability was determined by counting germ tubes produced on PDAY medium after 18 hrs, using light microscope at 400 x. Conidial viability was 95-100%. The surface of cultures was gently brushed in the presence of 20ml of sterilized water in order to free the spores and the suspension was filtered through muslin. Six concentrations of spore suspensions were prepared *i.e.*, 10^7 , 10^6 , 10^5 , 10^4 , 10^3 , and 10^2 conidia/ml. Piece of corn leaves were dipped in the prepared suspensions and left for drying under laboratory conditions then placed in Petri-dishes (one/dish). For each concentration (4 replicates/ each), ten L3 larvae of each of the tested insects were transferred into each Petri-dish. Control larvae were fed on untreated castor leaves. Percentages of mortality were calculated according to Abbott, while LC_{50} was calculated throughout probit analysis. The experiment was carried out under laboratory conditions at $26^\circ\text{C} \pm 2$ and 60-70 % RH. Physiological and metabolic characteristics of *P. fumosoroseus* and *P. lilacinus*

2.4 Semi-Field (green house) Trials

Potato plant as planted in the green house in 40 plots in each artificial infestation was made by spraying the plant with the bioinsecticides *Paecilomyces lilacinus* and *P. fumosoroseus* at the concentrations of 8.25×10^8 conidia/ml for each of the fungi. Control samples were sprayed by water only. The plants were examined every two days, the percentage of infestation was calculated until the end of the experiment. Each treatment was replicated 4 times. The percent mortality was counted and corrected according to Abbott, 1925 while LC_{50} s were calculated through probit analysis after Finney 1964.

2.5 Field Trials

Field trials were carried out at Nobaria region (Behera Governorate), Egypt during the two successive corn seasons 2012 and 2013 to study the effectiveness of the tested fungi on corn borers. Corn (variety Giza 2) was cultivated by end of May during the two seasons in an area of about half feddan. Fungi were applied as single treatments in randomized plots. Regular agricultural practices were performed and no chemical control was used during the study period. Weeds were removed by hand. Five plots were sprayed with water as control. Samples from each treatment were collected weekly and transferred to the laboratory for investigation. Percentages of infection were estimated.

2.6 Yield Assessment:

Yield data in treated and untreated plots in the corn harvest seasons (2012 and 2013), represented by weight in kgs were determined. Yield loss was estimated according to the following equation:

$$\text{Yield loss} = \frac{\text{Potential yield} - \text{Actual yield}}{\text{Potential yield}} \times 100$$

Potential yield is *Paecilomyces lilacinus* treatment (the best result among the tested pathogens) was considered the standard for comparison with the other ones (Actual yield).

3. Results and Discussion

Data in table 1 show that under laboratory conditions, the LC_{50} recorded 122×10^4 and 106×10^4 conidia/ml for the fungi, *Paecilomyces fumosoroseus* and *Paecilomyces lilacinus*, respectively. Under semifield conditions the corresponding LC_{50} 136×10^4 and 119×10^4 conidia/ml (Table 2). These results agree with Sabbour & Shadia Abd El-Aziz, (2002 and 2010), proved that applications with bioinsecticides increased the yield and decreased the infestation with insect pests. Similar results were obtained by Sabbour, (2003 and 2007), Sabbour and Sahab (2005 and 2007); Sabbour and Ismail (2001) and Ismail and Sabbour (2002). loss of the yield by Sabbour & Shadia Abd El-Aziz, (2002 and 2010), Shadia Abdel Aziz & Nofel (1998), proved that applications with bioinsecticides increased the yield and decreased the infestations. Sabbour and Sahab (2005 and 2007), reported that the different entomopathogenic fungi could reduce many insect pests under laboratory and field conditions and causing yield increase under field conditions. Sabbour, (1992 and 1995, 2003 and 2006), Sabbour and Sahab (2005 and 2007) reported that many insect pests could be controlled by the bioinsecticides under field conditions. The same obtain by Sabbour and Abdel-rahman 2007, who recorded LC_{50} s of the tested fungi against the *Phyllotreta cruciferae*, *Pegomya hyoscami* and *Cassida vittata* ranged between 5.4×10^6 and 1.43×10^7 spores/ml.

Table 3 show that in the field, the plots treated with *P. fumosoroseus* caused a significant decrease in infestation reached to 21 ± 9.45 and 20 ± 8.12 after 90 days of the post applications as compared to 89 ± 13.03 and 79 ± 11.12 individuals in the control during season 2012 and 2013, .

respectively. The infestations with potato tuber moth were significantly decreased to 26 ± 10.11 , 21 ± 9.12 and 20 ± 8.28 individuals after treatments with the fungus *P. lilacinus* after 20, 50 and 90 days, respectively during season 2012. During season 2013 the corresponding number of infestation reached to 23 ± 12.11 , 23 ± 9.11 and 25 ± 9.71 individuals, respectively. The same results obtained by Sabbour 2003, (20014a&b), 2013. Magda Mahmoud Sabbour and Shadia El-Sayed Abd-El-Aziz. 2014, Magda Sabbour, 2001, Sabbour (2002 a &b), Magda Sabbour and Ismail 2002, Sabbour and Sahab 2005 & 2007, 20011. These results agree with Sabbour & Shadia Abd El-Aziz, (2002 and 2010) Sabbour and Soliman (2014 a&b).

Table 4, show that the weight of the potato recorded during 2012 were, 5879 ± 8.28 and 5667 ± 7.18 kg/ feddan in the plots treated with *P. lilaceous* and *P. fumosoroseus* respectively as compared to 3398 ± 4.29 kg/ feddan in the control during season 2012 during season 2013 the highest weight recorded 5992 ± 8.21 kg/feddan after teament with *P. lilaceous*. (Sabbour & Sahab 2005, 2007, Tanda and Kaya, 1993 and Sahab and Sabbour 2011) found that the fungi *B. bassiana*, *M. anisopliae*, *Pacilomyces fumosoroseus* *Verticillium*

lecanii; reduced insect infestations of cabbage and tomato pests under laboratory and field conditions. Sabbour and Abdel-Rahman 2013 found that, in all treatments the number of corn pests were significantly decreased. loss of the yield by Sabbour & Shadia Abd El-Aziz, (2002 and 2010), proved that applications with bioinsecticides increased the yield and decreased the infestations. They found that the infestation was reduced after fungi applications under laboratory and field conditions. Sabbour & Sahab (2005, 2007 and 2011) found that the fungi reduced insect infestations of cabbage and tomato pests under laboratory and field conditions. These results agree with Sabbour & Shadia Abd El-Aziz, (2002 and 2010), proved that applications with bioinsecticides increased the yield and decreased the the infestation with insect pests.

4. Acknowledgment

This research was supported by Agric. Department, National Research Center, Cairo, Egypt. Project No (10120601).

Table 1: Effect of the entomopathogenic fungi, against (PTM) the target insect pests larvae under laboratory conditions.

Insects	LC ₅₀	slope	variance 95%	confidence limits
<i>Paecilomyces fumosoroseus</i>	122X10 ⁴	0.1	1.01	99-166
<i>Paecilomyces lilacinus</i>	106X10 ⁴	0.2	1.00	110-189

Table 2: Effect of the entomopathogenic fungi against the target insect pests larvae under semifield conditions.

Insects	LC ₅₀	slope	variance 95%	confidence limits
<i>Paecilomyces fumosoroseus</i>	136 X10 ⁴	1.01	0.02	111-176
<i>Paecilomyces lilacinus</i>	119 X10 ⁴	0.1	1.01	134-187

Table 3: Effect of different treatments on the target insect pests under field conditions

Post 1 st Application Date	Treatments	Number of infestation (means) ± s.e during both two seasons 2012/2013	
20	Control	52 ± 11.5	55 ± 7.89
50		53 ± 9.33	54 ± 5.56
90		89 ± 13.03	79 ± 11.12
20	<i>P. fumosoroseus</i>	31 ± 9.12	24 ± 8.23
50		28 ± 11.20	27 ± 7.66
90		21 ± 9.45	20 ± 8.12
20	<i>P. lilacinus</i>	26 ± 10.11	23 ± 12.11
50		21 ± 9.12	23 ± 9.11
90		20 ± 8.28	25 ± 9.71
F value		25.23	
Lsd%		.4141	

References

- [1] Abbott, W.W. A method of computing the effectiveness of an insecticide. J. Econ. Entomol 1925: 18(2) (2): 265-267.
- [2] Abdel-Rahman, M. A., A. Ahmed, Y. Abdel-Mallek and G.A. Hamam 2006. Comparative abundance of entomopathogenic fungi of cereal aphids in Assiut Egypt. J. Boil. Pest Cont.; 16(2): 39-43.
- [3] Abdel-Rahman, M.A.A. Seasonal prevalence of entomopathogenic fungi attacking cereal aphids infesting wheat in southern Egypt. Inter. Symposium. Agric. Agadir-Morocco. 2001: 7 (10): 381-389.
- [4] Abdel-Rahman, M.A.A and Abdel-Mallek, A.Y... Paramilitary records on entomopathogenic fungi attacking cereal aphids infesting wheat plants in southern Egypt. First Conference for safe Alternatives to pesticides for pest managements, Assiut. 2001(12): 183-190.
- [5] Abdel-Rahman, M.A.A.; Abdel-Mallek, A.Y.; Omar S.A. and Hamam, A.H... Natural occurrence of entomopathogenic fungi on cereal aphids at Assiut. A comparison study between field and laboratory observations. Egypt. J. Boil. Sci., 2004:14(2): 107-112.
- [6] Castillo, M.A.; Moya, P.; Hernandez, E. and Primo-Yufer, E. Susceptibility of *Ceratitis capitata* Wiedemann (Diptera: Tephritidae) to entomopathogenic fungi and their extract. Biol. Cont. 2000: 19(2): 274-282.
- [7] El-Husseini, M.M.; Shahira, S. Marie; A.M. Amal; A. El-Zoghby; Sahar S. Ali, Naglaa, A.M. Omar; E.A. Agamy; H.E. Abou Bakr; M.S. Nada; Sherin Tamer; Hannah, M. Kamal and A.M. Ibrahim. Isolation, Production and use of

Table 4: Assessments of damage caused in potato field after the fungi treatment

Treatments	Season 2012 Wt of potato crop (kg feddan) yield loss%	Season 2013 Wt of potato crop (kg/ feddan) yield loss%
<i>P. lilaceous</i>	5879 ± 8.28-	5992 ± 8.21-
<i>P. P. fumosoroseus</i>	5667 ± 7.183	5887 ± 6.201
control	3398 ± 4.2942	3255 ± 3.2645
F value	28.6	26.8
Lsd5% =	15.1	14.1

- entomopathogenic fungi for controlling the sugar beet insect pests in Egypt. Egypt. J. Biol. Pest Control. 2004;14(1): 265-275.
- [8] Espin, G.A. T. laghi De .S.M., Messias, C.L. and Pie-Drabuena, A.E. Pathogenicidad de *Metarhizium anisopliae* diferentes fases de desenvolvimiento de *Ceratitiscapitata* (Wied.) (Diptera: Tephritidae). Revista Brasileira de Entomologia. 1989 :33(2): 17-23.
- [9] Finney, D.J... Probit Analysis, Cambridge: Cambridge University Press 1971.
- [10] Qiao, Meihua; Daniel E. Snyder, Jeffery Meyer, Alan G. Zimmerman, Meihua Qiao, Sonya J. Gissendanner, Larry R. Cruthers, Robyn L. Slone, Davide R. Young (12 September 2007). "Preliminary Studies on the effectiveness of the novel pulicide, spinosad, for the treatment and control of fleas on dogs". *Veterinary Parasitology*. 2007 3(2): 345–351. Retrieved 3 May 2012.
- [11] Rice, R.E. 2000. Bionomics of the olive fruit fly *Bactrocera (Dacus) oleae*. Univ. of California Plant Prot. Quart. 2000: 4(2): 10:-1-5.
- [12] Roberts, D.W. and R.A. Humber. 1981. Entomogenous fungi In Biology of Conidial Fungi (G.T. Cole and W.B. Kendrick, eds). 1981: 1(2): 201-236. Academic Press, New York.
- [13] Rombach, M.C.; Aguda, R.M. and Robert, D.W. Production of *Beauveria bassiana* in different liquid media and subsequent conditions mycelium. Entomo. 1988: 3(33): 315-234.
- [14] Sabbour M.M. and Abdel-Rahman. 2013. Efficacy of isolated *Nomuraea rileyi* and Spinosad against corn pests under laboratory and field conditions in Egypt Annual review and research in biology, 3(4): 903-912. Sabbour, M.M., M. Ragei and A. Abd-El Rahman, 2011. Effect of Some Ecological Factors on The Growth of *Beauveria bassiana* and *Paecilomyces fumosoroseus* Against Corn Borers. Australian Journal of Basic and Applied Sciences, 5(11): 228-235, 2011
- [15] Sabbour, Magda M, 2002. Evaluation studies of some bio-control agents against corn borers in Egypt. Annal Agric. Sci. Ain Shams Univ. Cairo, 47(3): 1033-1043
- [16] Sabbour, M.M. and Shadia E. Abed El-Aziz. 2002, Efficacy of some botanical oils formulated with microbial agents against the cotton leafworm and greasy cutworm attaching cotton plants. Bull. ENT.Soc. Egypt. 2002: 5(28): 135-151.
- [17] Sabbour, M.M. and Sahab, A.F. Efficacy of some microbial control agents against cabbage pests in Egypt. Pak. J. Biol. Sci. 2005:5(8): 1351-1356.
- [18] Sabbour, M.M. and Sahab, A.F. 2007. Efficacy of some microbial control agents against *Agrotisipsilon* and *Heliothis armigera* in Egypt. Bull. N.R.C. Egypt. 2007: 13(33): 165-174
- [19] Sabbour, M.M. and Shadia, E. Abd-El-Aziz. 2010. Efficacy of some bioinsecticides against *Bruchidius incarnatus* (BOH.) (Coleoptera: Bruchidae) Infestation during storage. 2010 J. Plant Prot. Res. 2010: 50 (1): 28-34.
- [20] Sabbour, M. Mand Abdel-Rahman, A . 2007. Evaluations of some terpenes and entomopathogenic fungi on three sugar beet insect pests. J. Boil. Pest. Cont. 17:22-29
- [21] Sabbour, M. M, (2008). Evaluations of some microbial control agents against olive moth *Prays oleae* under field conditions under publication.
- [22] Sabbour, M.M and Nayera, Y. Soliman, 2014a,. Evaluations of three *Bacillus thuringiensis* against *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Egypt. **Volume 3 Issue 8, August 2014**. 2319-7064.
- [23] Sabbour, M.M. and Nayera, Y. Soliman. 2014b. Preliminary Investigations Into The Biological Control Of Red Palm Weevil *Rhynchophorus ferrugineus* By Using three isolates of the fungus *Lecanicillium (Verticillium) lecanii* In Egypt **Volume 3 Issue 8, August 2014**. 2319-7064
- [24] Sabbour, M.M. 2009. Evaluation of two entomopathogenic fungi against some insect pests infesting tomato crops in Egypt , IOBC/wprs Bulletin, Vol. 49: 273-278.
- [25] Sabbour, M.M., M. Ragei and A. Abd-El Rahman, 2011. Effect of Some Ecological Factors on The Growth of *Beauveria bassiana* and *Paecilomyces fumosoroseus* Against Corn Borers. Australian Journal of Basic and Applied Sciences, 5(11): 228-235, 2011
- [26] Sabbour, M.M . (2012). Evaluations of some bioagents against the rice weevil *Sitophilus oryzae* under laboratory and store conditions. *Integrated Protection of Stored Products. IOBC-WPRS Bulletin Vol. 81, pp. 135-142*
- [27] Sabbour M.M. and M.A. Abd-El-Raheem. 2013. Repellent Effects of *Jatropha curcas*, canola and Jojoba Seed oil, against *Callosobruchus maculatus* (F.) and *Callosobruchus chinensis* (L.). Journal of Applied Sciences Research, 9(8): 4678-4682, 2013
- [28] Sahab, A .F and Sabbour, M.M, (2011). Virulence of four entomopathogenic fungi on some cotton pests with especial reference to impact of some pesticides, nutritional and environmental factors on fungal growth. Egypt. J. biol pest cont. 21 (1): 61-67.
- [29] Sahab ,A.F; Sabbour , M.M., Attallah, A.G. and Abou-Serreh, Nivin. 2014. Genetic analysis of the entomopathogenic fungus *Beauveria bassiana* to the corn borers tested by UV as physical mutagen. *International Journal of ChemTech Research* Vol.6, No.5, pp 3228-3236, .
- [30] Sameh A. Moustafa; Ahmed E. Abd El-Mageed; Mostafa M. El-Metwally and Nabil M. Ghanim. Efficacy of Spinosad, Lufenuron and Malathion against olive fruit fly, *Bactrocera oleae* (Gmelin) (Diptera: Tephritidae) Egypt. Acad. J. biolog. Sci. 2009: 2 (2): 171- 178.
- [31] Shadia E. Abdel Aziz and Nofel, M.A. The efficacy of bacteria, fungi and natural products in baits against the greasy cutworm *Agrotisipsilon* (Hufn.) (Lepidoptera: Noctuidae) in Egypt. J. Egypt. Ger. Soc. Zool., 1998: 27(2). 129-139.
- [32] Tanda, Y. and Kaya, H.K... Insect Pathology. Academic Press, San Diego, CA, USA. 1993: 3(4): 123-129.
- [33] Sabbour M.M. 2014 a. Biocontrol of the Tomato Pinworm *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae) in Egypt. Middle East Journal of Agriculture Research. 2077-4605.
- [34] Sabbour, M.M. 2014 b. Efficacy of some microbial control agents and inorganic insecticides against red flour beetle *Tribolium castaneum* and confused flour beetle, *Tribolium confusum* (Coleoptera: Tenebrionidae) Integrated Protection of Stored Products. IOBC-WPRS Bulletin Vol. 98, 2014. pp. 193-201.