

Occurrence of Multiple Antibiotic Resistances in *Escherichia coli* Isolated from Children of Diarrheal Symptoms

M. S. Al- Hussaini¹, A. M. Saadabi², R. Mohammed Abdallah³

^{1,2}Department of Clinical Laboratory Sciences, College of Applied Medical Sciences, Shaqra University, Kingdom of Saudi Arabia

³Faculty of Science & Technology, Al-Neelain University, Khartoum, Sudan

Abstract: This study was carried out to evaluate the resistance patterns of pathogenic *Escherichia coli* which cause diarrhea in children less than 5 years. Stool samples were collected from young children suffering from diarrhea in different hospitals and health centers, and the bacteria was then isolated and identified. The susceptibility of pathogenic *Escherichia coli* for seven known used antibiotics was then tested. The study showed that among every 3 young children suffering from diarrhea there is about one child his case was severe and infected with pathogenic *Escherichia coli* and need antibiotic medication, so the ratio between non-pathogenic and pathogenic *Escherichia coli* during this study was (2:1). The bacteria under study shows 100% resistance for Chloramphenicol, 62.5% resistance and 37.5% intermediate sensitivity for Erythromycin and intermediate sensitivity of 72.5% for Amoxicillin. Also it showed 75% sensitivity and 25% intermediate sensitivity for Amoxicillin / Clavulanic acid. The susceptibility testing for Nalidixic Acid, Ampicillin / Sulbactam and Cotrimoxazole showed strong sensitivity of 100% against pathogenic *Escherichia coli*.

Keywords: Antibiotic resistance, *Escherichia coli*, diarrheagenic bacteria, Diarrheal Symptoms

1. Introduction

Acute diarrhea is one of the most commonly reported illnesses in the world, second only to respiratory tract infections (Levine, 1987). Worldwide, it is the leading cause of mortality in children under the age of four years. Diarrheal diseases are major causes of morbidity and mortality, with attack rates ranging from two to twelve or more illnesses per person per year in developed and developing countries (Atif *et al.*, 2000, Kosek *et al.*, 2003). The causes of diarrhea include a wide array of viruses, bacteria, and parasites, many of which have been recognized only in the last decade. The enterotoxigenic *Escherichia coli* and tavirus predominate in developing areas (Alrifai *et al.*, 2009; Bhan *et al.*, 1989).

The use of antimicrobial drugs is now threatened by the global emergence of resistant strains. The process is fostered by the remarkable genetic diversity and transfer of resistance among microbes, the selective forces of antibiotic use, and powerful socioeconomic and marketing forces (Alrifai *et al.*, 2009). In developing countries, acquired bacterial resistance to antimicrobial agents is common in isolates from healthy persons and from persons with community-acquired infections. Complex socioeconomic and behavioral factors associated with antibiotic resistance, particularly regarding diarrheal and respiratory pathogens, in developing tropical countries, include misuse of antibiotics by health professionals, unskilled practitioners, and laypersons; poor drug quality; unhygienic conditions accounting for spread of resistant bacteria; and inadequate surveillance (Ochoa *et al.*, 2008). Knowledge of drug resistance in bacteria is indispensable for the proper selection of antimicrobial drugs. Saudi Arabia allots a substantial amount of its health-expenditure allocation to the purchase of drugs, especially antibiotics (Cristian Pérez *et al.*, 2010).

The sensitivity of *Escherichia coli* isolated from children suffering from diarrhea has been reported from different parts of the world with varying results. According to an epidemiological study in Vietnam, *Escherichia coli* is most commonly isolated microorganism in clinical samples from a patient with diarrhea and it shows a high prevalence of resistance to antibiotics (Trung *et al.*, 2005). The prevalence of resistance to commonly used antibiotics has also been investigated in groups of children from four developing countries; Peru, Belize, Zaire and Sudan (Vila, 1999). Enterobacteriaceae species isolated from faeces of children with diarrhea were sensitive to ampicillin, tetracycline, cotrimoxazole, streptomycin and chloramphenicol. Overall, the highest prevalence of resistance was to cotrimoxazole (56% of children) and the lowest was to chloramphenicol (19% of children).

The objectives of the present work are to determine the incidence of antimicrobial-resistant, enteropathogenic *Escherichia coli* among children of less than five years at a number of hospitals and clinics of the area.

2. Materials and Methods

Study population and sample collection

Samples in this study were taken from young children (under 5 years), at a number of hospitals and clinics of the area. The children were considered as suffering from diarrhea if they had passed the clinical tests in the last 12 h. All of the procedures of laboratory work were carried out in faculty of Science, Khartoum and College of Medical Applied Sciences, Shaqra University. Serotyping of all samples was performed in ministry of health Laboratories, Khartoum.

2.1 Isolation of Bacteria

Samples were collected aseptically in clean, sterile bottles and immediately cultured on xylose lysine deoxycholate (XLD) agar overnight at 37 °C. Samples of yellow color were then cultured on plates containing Eosin Methylene Blue (EMB) agar to differentiate and select *E.coli* by incubating at 37°C (overnight). From EMB, subcultures of each sample were done on nutrient agar for biochemical tests.

2.2 Biochemical tests

Testing for indole production is important in the identification of enterobacteria. Most strains of *E. coli* breakdown the amino acid tryptophan with the release of indole. The test organism was cultured in a medium containing tryptophan. Indole production was detected by using Kovac's or Ehrlich's reagent, which contains 4 (p)-dimethylaminobenzaldehyde. This reagent reacts with the indole to produce a red-colored compound. The test organism was inoculated in a medium test tube containing 3 ml of sterile peptone water, and incubated at 35–37°C for up to 48 h. followed by testing for indole by addition of 0.5 ml of Kovac's reagent, shaken gently and examined for red color in the surface layer within 10 minutes (Vila, 1999).

2.3 Sero-typing

A drop of normal saline was placed on a clean microscope slide, inoculation from culture was added and a suspension was made, followed by addition of a drop of antisera. The slides were then rotated for 10 min and examined under the microscope to detect any clumping. Different reagents for different strains of *E.coli* were used that included: Enter pathogenic *E.coli* (EPEC), Enteroinvasive *E.coli* (EIEC), Enterotoxigenic *E.coli* (ETEC) and Enterohemorrhagic *E.coli* (VTEC). From each bacterial culture, four suspensions in four different slides were made for the four reagents, each reagent testing specific strains of *E.coli* (Sehgal, 1999).

2.4 Susceptibility of isolated bacteria to different antibiotics

In this study, sensitivity of the different isolates to a number of antibiotics that are commonly used in hospitals and community was studied using the standard disc diffusion technique. Mueller Hinton sensitivity testing agar medium was prepared and sterilized as instructed by the manufacturer and poured into sterile Petri dishes. The inoculums were prepared by emulsifying 3-5 colonies of the test organism in about 3-5 ml of sterile normal saline. A sterile cotton swab was dipped into this diluted culture; the swab was pressed against the side of the tube to remove excess fluid, and streaked across the medium in three directions by rotating the plate approximately 60° to ensure even distribution. (Williams and Ryan, 1998; Robins-Browne, 1987). A paper discs saturated with a known amount of the antibiotic were used. The antibiotic discs were carefully placed on the inoculated plates; each disc was lightly pressed down to ensure its contact with the agar. After overnight incubation at 37 °C aerobically, the culture was examined for zones of inhibition of bacterial growth around the respective discs.

2.5 Interpretation of zone sizes

Using the interpretative chart, the zone sizes of each antibiotic was determined to classify the organisms as resistant, intermediate/moderately sensitive, sensitive (susceptible) (Robins-Browne, 1987).

3. Results

The total samples collected in the present study were 45, out of this, 30 were *E.coli* (showed metallic green on EMB) and 10 of them did not give positive result (metallic green) on EMB and the indole test (Table 1). Other remaining 5 samples didn't ferment the sugar on XLD (XLD media did not change to the yellow color) so they were rejected (Table 1).

All of the *E. coli* samples were subjected to sero-typing tests in order to classify those strains to: EHEC, EPEC, VTEC and EIEC. The results of sero-typing tests are summarized in Table 2 and Figures 4 and 5 shows the clumping of bacterial cells as a result of reaction with antisera.

Antibiotic sensitivity tests were done using the standard disk diffusion method. A measurement for zone of inhibition of bacterial growth was measured in millimeters (Table 3).

Table 1: Identification of stool samples (Culture media and biochemical testing)*

Sample number	XLD	EMB	Indole test
1	yellow	Metallic green	+ve
2	yellow	Metallic green	+ve
3	yellow	Metallic green	+ve
4	yellow	Metallic green	+ve
5	yellow	Metallic green	+ve
6	yellow	Metallic green	+ve
7	yellow	Metallic green	-ve
8	yellow	Metallic green	+ve
9	yellow	Metallic green	+ve
10	yellow	Metallic green	+ve
11	yellow	Metallic green	+ve
12	yellow	Metallic green	-ve (Suspected to be new strain)
13	yellow	Metallic green	+ve
14	yellow	Pink (unusual case)	+ve
15	yellow	Metallic green	+ve
16	yellow	Metallic green	+ve
17	yellow	Pink (unusual case)	+ve
18	yellow	Metallic green	+ve
19	yellow	Metallic green	+ve
20	yellow	Metallic green	+ve
21	yellow	Metallic green	+ve
22	yellow	Metallic green	+ve
23	yellow	Metallic green	+ve
24	yellow	Metallic green	+ve
25	yellow	Metallic green	+ve
26	yellow	Metallic green	+ve
27	yellow	Metallic green	+ve
28	yellow	Metallic green	-ve (Suspected to be new strain)
29	yellow	Metallic green	+ve
30	yellow	Metallic green	+ve

*XLD = Xylose Lysine Deoxycholate agar EMB= Eosin Methylene Blue

Methylene Blue.

+ve= appearance of Red surface layer, -ve=No appearance of Red surface layer

Table 2: Identification of bacterial pathotypes using four different types of antisera

Sample Number	EIEC	VTEC	EPEC	ETEC
1	-ve	-ve	-ve	-ve
2	-ve	-ve	-ve	-ve
3	-ve	-ve	-ve	-ve
4	-ve	-ve	+ve	-ve
5	-ve	-ve	-ve	-ve
6	-ve	-ve	-ve	-ve
7	-ve	-ve	-ve	-ve
8	-ve	-ve	-ve	-ve
9	-ve	-ve	-ve	-ve
10	-ve	-ve	+ve	-ve
11	-ve	-ve	-ve	-ve
12	-ve	-ve	-ve	-ve
13	-ve	-ve	+ve	-ve
14	-ve	-ve	-ve	-ve
15	-ve	-ve	-ve	-ve
16	-ve	-ve	+ve	-ve
17	-ve	-ve	-ve	-ve
18	-ve	-ve	-ve	-ve
19	-ve	-ve	-ve	-ve
20	-ve	-ve	-ve	-ve
21	-ve	-ve	-ve	-ve
22	-ve	-ve	-ve	-ve
23	-ve	-ve	+ve	-ve
24	-ve	-ve	+ve	-ve
25	-ve	-ve	-ve	-ve
26	-ve	-ve	-ve	-ve
27	-ve	-ve	-ve	-ve
28	-ve	-ve	-ve	-ve
29	-ve	-ve	+ve	-ve
30	-ve	-ve	+ve	-ve

*EPEC=Enteropathogenic E.coli

EIEC= Entero invasive E.coli

ETEC= Entero Toxogenic E.coli

VTEC=Entero homoraggic E.coli

-ve = no clumping appears

+ve -= clumping appearance under microscope

Table 3: Inhibition zones for seven common used antibiotics against pathogenic E.coli

Samples containing pathogenic E.coli	Na	Ac	As	Co	E	C	AX
Sample No. 4	30	19	24	30	14	0	25
Sample No. 10	29	18	24	32	13	0	24
Sample No. 13	25	18	21	27	9	0	20
Sample No. 16	28	20	24	30	12	0	23
Sample No. 23	29	19	24	30	12	0	20
Sample No. 24	28	18	23	31	12	0	23
Sample No. 29	30	14	24	32	13	0	23
Sample No. 30	29	14	23	31	12	0	25

Na= Nalidixic Acid (30 mcg)

Ac= Amoxycillin / Clavulanic acid (30 mcg) As =Ampicillin / Sulbactam (10/10 mcg)

Co= Co-Trimoxazole (Trime thoprim/Sulphamethoxazole) (1.25/ 23.75 mcg) E= Erythromycin (15 mcg)

C= Chloramphenicol (30 mcg)

AX= Amoxicillin (25 mcg)

Inhibition zones in (mm)

Table 4: Resistance patterns for seven common used antibiotics against pathogenic E.coli*

Sample number containing pathogenic E.coli	Na	Ac	As	Co	E	C	AX
4	S	S	S	S	I	R	S
10	S	S	S	S	I	R	S
13	S	S	S	S	R	R	I
16	S	S	S	S	R	R	I
23	S	S	S	S	R	R	I
24	S	S	S	S	R	R	I
29	S	I	S	S	I	R	I
30	S	I	S	S	R	R	S

Na= Nalidixic Acid (30 mcg)

Ac=Amoxycillin / Clavulanic acid (30 mcg) As =Ampicillin / Sulbactam (10/10 cg)

Co =Co-Trimoxazole (Trime thoprim/Sulphame thoxazole) (1.25/ 23.75 mcg) E= Erythromycin (15 mcg)

C= Chloramphe nicol (30 mcg) AX= Amoxicillin (25 mcg)

S= Sensitive

I= Intermediate R= Resistant



Figure 1: Samples cultured on EMB media shown metallic green E.coli colonies




Figure 2: Testing samples for indole production



Figure 3: Positive indole sample (red surface layer)



Figure 4: Clumping of bacterial cells on slide as a result of reacting with antisera




Figure 5: C lumping of bacterial cells under microscope as a result of reaction with antisera.



Figure 6: Susceptibility of pathogenic E.coli for seven common used Antibiotics

4. Discussion

The present study determined the resistance of E. coli isolated from stools of young children. The isolation and identification of bacteria were done on samples collected under strict sterile conditions to avoid contamination.

Antimicrobial resistance has become a major medical and public health problem. The main factor responsible for development and spread of bacterial resistance is injudicious use of antimicrobial agents, which has resulted in most gram positive and gram negative bacteria continuously developing resistance to the antimicrobials in regular use at different time periods (Urassa, 1997). The antimicrobial sensitivity is routinely performed to identify the best possible regimen for the treatment of bacterial infections. There are several reports from different parts of the world about antimicrobial sensitivity with varying results indicating the bacterial resistance to antimicrobials differs in different areas. In

Sudan, antimicrobial sensitivity tests were performed on four-hundred and ninety-seven bacterial isolates from patients with diarrhea, and enteropathogenic *Escherichia coli*. The results showed high resistance rates against the commonly used antimicrobial agents; ampicillin, amoxicillin, chloramphenicol, tetracycline, cotrimoxazole and nalidixic acid (Atif et al., 2000). Similarly, a study in Vietnam on 162 diarrheagenic *E. coli* isolates reported 77.2% resistance to chloramphenicol and 19.1% resistant to nalidixic acid (Trung et al., 2005).

The prevalence and antimicrobial susceptibility of diarrheagenic *Escherichia coli* in 346 children from Tanzania showed that diarrhea was due to enteropathogenic *E. coli* and there was high-level resistance to ampicillin, tetracycline, cotrimoxazole, and chloramphenicol but susceptible to quinolones (Vila et al., 1999). As mentioned earlier, the prevalence of resistance to commonly used antibiotics in groups of children from four developing countries; Peru, Belize, Zaire and Sudan (Vila et al., 1999) showed that isolated microorganism were sensitive to ampicillin, tetracycline, cotrimoxazole, streptomycin and chloramphenicol. In the present study, all samples containing enteropathogenic *E. coli* were resistant to chloramphenicol, 62.5% of them were resistant to erythromycin, and intermediate sensitivity of 72.5% was observed for amoxicillin. The results of the present study were similar to those reported by Vila et al. (1999) and Atif et al. (2000). However, the sensitive of pathogenic *Escherichia coli* to nalidixic was different from that reported by Trung et al. (2005), where they found 19.1% of isolated pathogenic *Escherichia coli* resistant to nalidixic acid and in the present study no resistance was observed.

During informal discussion with children's mothers, we learnt that 90% of young children had used different classes of antibiotics during the last 6 months. We believe that use of these antibiotics might have led to evolution of new resistant strains of bacterial pathogens. There was also strong correlation between hygiene, child activity and the infection with diarrhea. It is well known that indiscriminate use of some antibiotics, especially in young children leads to development of resistance to other classes of antibiotics apart from serious adverse effects associated with their use.

The susceptibility of enteropathogenic *Escherichia coli* for cotrimoxazole this study was high (100% effectiveness against bacteria under study). This result was contradictory to that reported in isolates from four different countries by Vila et al., (1999), where high level of resistance was observed for cotrimoxazole. The Ampicillin/Sulbactam combination was also very effective with 0% resistance. This wide spectrum antibiotic is most commonly used to treat diarrheal diseases.

Chloramphenicol failed to prevent the growth of enteropathogenic *E. coli* indicating that this drug is not suitable for use against these bacteria any more.

From the results of the present study, it is recommended that a routine program must be developed for regular sensitivity screening of commonly used antibiotics in the treatment of diarrhea. The regular determination of antibiotic sensitivity

will help in reducing incidence of bacterial resistance and in prevention of adverse effects due to indiscriminate use of ineffective antibiotics. Furthermore, the use of highly effective antibiotics has a possible health risk, particularly for children under five. The routine screening of sensitivity will also help in selection of antibiotics with less adverse effects, especially in young children.

5. Conclusion

The enteropathogenic *E. coli* strains isolated from children under five years of age were found to be highly resistant to chloramphenicol, while amoxicillin showed a weak effect. Intermediate effect for amoxicillin/clavulanic acid and a very weak effect for erythromycin were observed. The *E. coli* strains were high sensitive to nalidixic acid, ampicillin/sulbactam and cotrimoxazole.

References

- [1] Alrifai S.B., A .Alsaadi , Y.A. Mahmood , A.A. Ali , L.A. Al-Kaisi, (2009). Prevalence and etiology of nosocomial diarrhoea in children < 5 years in Tikrit teaching hospital, East Mediterr. Health J. 15(5):1111-1118.
- [2] Atif A. Ahmed, H. Osman, A. M. Mansour, A. M. Hassan, B. A. Abdalla, Z. Karrar and H. S. Hassan (2000). Antimicrobial agent resistance in bacterial isolates from patients with Diarrhea and urinary tract infection in the Sudan, Am. J. Trop. Med. Hyg., 63(5): 259-263.
- [3] Behiry I.K., E.A. Abada, E.A. Ahmed, R.S. Labeeb (2011). Enteropathogenic *Escherichia coli* associated with diarrhea in children in Cairo, Egypt, Scientific World Journal; 11:2613-2619.
- [4] Bhan M.K., P .Raj, M.M. Levine (1989). Enteropathogenic *Escherichia coli* association with persistent diarrhea in a cohort of rural children in India. J Infect Dis, 159:1061-1064.
- [5] Cristian Pérez, Oscar G. Gómez-Duarte and María L. Arias, (2010).
- [6] Diarrheagenic *Escherichia coli* in children from Costa Rica, Am. J. Trop. Med. Hyg., 83(2): 292-297.
- [7] Jordi Vila, Martha Vargas, Climent Casals, (1999). Antimicrobial Resistance of Diarrheagenic *Escherichia coli* Isolated from children under the Age of 5 Years from Ifakara, Tanzania. ANTIMICROBIAL AGENTS AND CHEMOTHERAPY, p. 3022-3024.
- [8] Kosek M, C. Bern, R.L. Guerrant (2003). The global burden of diarrhoeal disease, as estimated from studies published between 1992 and 2000, Bull World Health Organ. 81(3):197-204.
- [9] Levine M. M. (1987). *Escherichia coli* that cause diarrhea: enterotoxigenic, enteropathogenic, enteroinvasive, enterohemorrhagic and enteroadherent. J Infect Dis; 15:377-89.
- [10] Ochoa T.J., F. Barletta, C. Contre ras, E. Mercado (2008). New insights into the epidemiology of enteropathogenic *Escherichia coli* infection, Trans R Soc Trop Med Hyg. 102(9):852-6.
- [11] Robins-Browne R. M. (1987). Traditional enteropathogenic *Escherichia coli* of infantile diarrhea, Rev Infect Dis. 9(1):28-53.

- [12] Sehgal R. (1999). Combating antimicrobial resistance in India. *JAMA*; 281:1081–2.
- [13] Shears P., C.A. Hart, G.S. Suliman (1988). A preliminary investigation of antibiotic resistance in Enterobacteriaceae isolated from children with diarrhoea from four developing countries, *Ann Trop Med Parasitol*. 82(2):185-8.
- [14] Shears P, G. Suliman, C.A. Hart (1988). Occurrence of multiple antibiotic resistance and R plasmids in Enterobacteriaceae isolated from children in the Sudan, *Epidemiol Infect*. 100(1):73-81.
- [15] Trung Vu Nguyen, Phung Van Le, Chinh Huy Le, and Andre j Weintraub, (2005). Antibiotic Resistance in Diarrheagenic Escherichia coli and Shigella Strains Isolated from children in Hanoi, Vietnam, *Antimicrobial Agents and Chemotherapy*, 18: 816–819.
- [16] Urassa W, E. Lyamuya, F. Mhalu (1997). Recent trends on bacterial resistance to antibiotics, *East Afr Med J*. 74(3):129-33.
- [17] Vila J, M. Vargas, C. Casals, H. Urassa, H. Mshinda, D. Schelleberg, J. Gascon, (1999). Antimicrobial resistance of diarrheagenic Escherichia coli isolated from children under the age of 5 years from Ifakara, Tanzania, *Antimicrob Agents Chemother*. 43(12):3022-4.