









88-223 °C and 60-225 while at high RF power, the temperature varied from 143-278 °C and from 140-179 °C at the electrode and tumor surface, consequently. Lesions continue to grow while temperature envelopes collapse after ablation since sufficiently high temperature is present to accrue tissue damage. In accordance to our results when the temperature of the electrode increases the tumor tissue was ablated and this was under ultrasound guidance and followed up with CT and alpha fetoprotein and a sudden increase in electrical impedance occurred. The efficacy of RFA can be assessed with contrast-enhanced ultrasound; contrast enhanced dynamic CT (CECT) or contrast-enhanced magnetic resonance imaging (MRI). A successful RFA treated area of HCC is nonenhancing on CECT and tends to be larger than the original tumor.<sup>(19)</sup> In this study we clarify the important role of contrast enhanced dynamic CT, all patients had pass through a routine CT abdomen with contrast and ultrasound examinations before and after treatment. The study was done under guidance of ultrasound. The texture of the tumor varied from before and after RF treatment this texture is monitored using abdominal CT with contrast before and after the treatment this variation in texture is caused because of that HCC is a malignant tumor which is supplied with blood vessels so when injecting the patient with contrast media before RF treatment the tumor appears a bright (hyper dense), but after Applying the RF treatment the texture appears to be black (hypodense), this means that the radiofrequency waves had applied the appropriate temperature to this tumor and destroyed all of its blood supplies. Also this is proved by laboratory investigation by measuring AFP before and after the treatment. The effectiveness of RFA. For lesions larger than 3 cm could be increased by several applications or adjuvant therapies should be considered to increase the diameter of coagulation hyperthermic cellular sensitization by chemotherapy<sup>(20)</sup>, vascular clamping<sup>(21)</sup>, or cellular ischemia (chemoembolization)<sup>(22)</sup> and alteration of electric and thermal conductivity (instillation of saline injection)<sup>(23)</sup>. Recent progress in bipolar electrodes could provide better control of the shape of large coagulations<sup>(24)</sup>. HCC is the third leading cause of cancer-related death and the fifth - sixth most common cancer<sup>(25-26)</sup>. The tissue itself is directly heated, rather than the probe itself.<sup>(27)</sup> Above ~50°C, protein denaturation results in irreversible tissue damage.<sup>(28)</sup> Above ~70 °C, coagulation occurs, where collagens are converted to glucose. Above 100 °C, water vapor develops, and tissue charring can occur, this is undesirable, since it raises the impedance and makes the treatment ineffective. Therefore, impedance between the probe and the dispersive electrode is measured during the treatment, and the generator is shut down automatically if the impedance exceeds a certain value (~ 200 to 1000 W, depending on generator type).<sup>(29)</sup>

## 7. Summary and Conclusion

Hepatocellular Carcinoma is one of the most common and dangerous diseases that attacks the human liver, it is considered the fifth cause of death worldwide, in Egypt a high incidence of HCC had been reported, there is a strong association between HCV infection & cirrhosis and HCC. RF ablation (RFA) is a simple, effective, and less expensive technique with a low morbidity compared with surgical treatment RFA can produce significant long-term survival rates and excellent local control. For cirrhotic patients with

early-stage, unrespectable HCC Yet, variable factors determine its efficacy e.g. the size of the lesion, its morphology, the operator expertise and low incidence of major complications is usually associated.

## References

- [1] Llovet JM, Burroughs A, Bruix J. Hepatocellular carcinoma. *Lancet* 2003;362:/1907-17.
- [2] Lounsberry W, Goldschmint V, Linke C, et al: The early histologic changes following electrocoagulation. *J. Urol.* 1961; 86:321-9.
- [3] Lee CS, Sheu JC, Wang M, Hsu HC. Long-term outcome after surgery for asymptomatic small hepatocellular carcinoma. *Br J Surg* 1996;/83:/330-3.
- [4] Bruix J, Sherman M. Management of hepatocellular carcinoma. *Hepatology* 2005;/42:/1208-36.
- [5] Llovet JM, Fuster J, Bruix J. Intention-to-treat analysis of surgical treatment for early hepatocellular carcinoma: resection versus transplantation. *Hepatology* 1999;/30:/1434-40.
- [6] Mazzaferro V, Regalia E, Doci R, Andreola S, Pulvirenti A, Bozzetti F, et al. Liver transplantation for the treatment of small hepatocellular carcinomas in patients with cirrhosis. *N Engl J Med* 1996;/334:/693-9.
- [7] Tungjitkusolmun S, Woo EJ, Cao H, Tsai JZ, Vorperian VR, Webster JG. Thermal – electrical finite element modelling for radio frequency cardiac ablation: effects of changes in myocardial properties. *Med Biol Eng Comput.* 2000; 38:562–568.
- [8] Livraghi T, Goldberg SN, Lazzaroni S, Meloni F, Solbiati L, Gazelle GS. Small hepatocellular carcinoma: treatment with radio-frequency thermal ablation versus ethanol injection. *Radiology* 1999;/210:/655-61.
- [9] Lu DS, Yu NC, Raman SS, Limanond P, Lassman C, Murray K, et al. Radiofrequency ablation of hepatocellular carcinoma: treatment success as defined by histologic examination of the explanted liver. *Radiology* 2005;/234:/954-60.
- [10] Wakai T, Shirai Y, Suda T, Yokoyama N, Sakata J, Cruz P, et al. Long-term outcomes of hepatectomy vs percutaneous ablation for treatment of hepatocellular carcinoma 54 cm. *World J Gastroenterol* 2006; 12:546-52.
- [11] LeVeen RF. Laser hyperthermia and radiofrequency ablation of hepatic lesions. *Sem Intervent Radiol* 1997; 14:313-324.
- [12] McGahan J, Browning P, Brock J, et al: Hepatic ablation using radiofrequency electrocautery. *Invest. Radiol.* 1990; 25:267-70.
- [13] d'Arsonval A. Action physiologique des courants alternatifs. *Comp. Rend. Soc. Biol.* 1891; 43- 283.
- [14] Clark W, Morgan J, Asnia E: Electrothermic methods in treatment of neoplasms and other lesions with clinical and histological observations. *Radiology.* 2:233-246; 1924.
- [15] Beer E: Removal of neoplasms of urinary bladder; a new method employing high frequency currents through a cauterizing cystoscope. *JAMA.* 1910; 54:1768-9.

- [16] Poon RT, Ng KK, Lam CM: Learning curve for radiofrequency ablation of liver tumors: prospective analysis of initial 100 patients in a tertiary institution. *Ann Surg.* 2004; 239:441-49.
- [17] Tungjtkusolmun S: Finite element analyses for a study of hepatic cancer tissue destruction using monopolar and bipolar radiofrequency ablation. *IJA Biomed Eng* 2009; 2:33-38.
- [18] Mulier S, Mulier P, Ni Y: Complications of radiofrequency coagulation of liver tumours. *Br J Surg.* 2002; 89:1206-22.
- [19] Curley SA, Izzo F, Ellis LM, et al: Radiofrequency ablation of hepatocellular cancer in 110 patients with cirrhosis. *Ann Surg.* 2000; 232:381-91.
- [20] Haemmerich D, Tungjtkusolmun S, Staelin ST, Lee FT, Jr, Mahvi DM, Webster JG. Finite-element analysis of hepatic multiple probe radio-frequency ablation. *IEEE Trans Biomed Eng.* 2006; 49:836-42.
- [21] Berjano EJ, Alio JL, Saiz J. Modeling for radio-frequency conductive keratoplasty: implications for the maximum temperature reached in the cornea. *Physiol Meas.* 2005; 26:157-72.
- [22] Wiley JD, Webster JG. Analysis and control of the current distribution under circular dispersive electrodes. *IEEE Trans Biomed Eng.* 1982;29:381-385.
- [23] Overmyer KM, Pearce JA, DeWitt DP. Measurements of temperature distributions at electro- surgical dispersive electrode sites. *Trans ASME, J Biomechanical Engineering.* 1979; 101:66-72.
- [24] Kim Y, Webster JG, Tompkins WJ. Simulated and experimental studies of temperature elevation around electrosurgical dispersive electrodes. *IEEE Trans Biomed Eng.* 1984; 31:681-92.
- [25] Shiina S, Tateishi R, Arano T, et al. Radiofrequency ablation for hepatocellular carcinoma: 10-year outcome and prognostic factors. *Am J Gastroenterol* 2012;107:569-577.
- [26] Lencioni R, Crocetti L. Local-regional treatment of hepatocellular carcinoma. *Radiology* 2013;262:43-58.
- [27] Elias D, Santoro R, Ouellet JF, et al: Simultaneous percutaneous right portal vein embolization and left liver tumor radiofrequency ablation prior to a major right hepatic resection for bilateral colorectal metastases. *Hepatogastroenterology.* 2004; 51:1788-91.
- [28] Merkle EM, Goldberg SN, Boll DT, et al: Effects of superparamagnetic iron oxide on radio-frequency-induced temperature distribution: in vitro measurements in polyacrylamide phantoms and in vivo results in a rabbit liver model. *Radiology.* 1999; 212:459-66.
- [29] De Baere T, Bessoud B, Dromain C, et al: Percutaneous radiofrequency ablation of hepatic tumors during temporary venous occlusion. *AJR Am J Roentgenol.* 2002; 178:53-9.