

# Yield Performance of Different Genotypes of Mustard under Agro-Climatic Conditions of Bahawalpur, Pakistan

Malik Muhammad Yousaf<sup>1</sup>, Mumtaz Hussain<sup>2</sup>, Dr. Nazim Hussain<sup>3</sup>, M. Jahangir Shah<sup>4</sup>, Bashir Ahmed<sup>5</sup>, Hafiz Muhammad Zia Ullah Ghazali<sup>3</sup>

<sup>1, 2, 4, 5</sup> Arid Zone Research Institute (AZRI), Bahawalpur, Pakistan

<sup>3</sup>Department of Agronomy, Faculty of Agriculture Sciences and Technology, BZU Multan, Pakistan

<sup>6</sup>Regional Agricultural Research Institute (RARI) Bahawalpur, Pakistan

**Abstract:** *The study regarding yield performance of ten genotypes of mustard under agro-climatic conditions of Bahawalpur region was conducted at Experimentation Station of Arid Zone Research Institute (AZRI), Bahawalpur. Ten genotypes of mustard were tested for their yield potential under agro-climatic conditions of Bahawalpur. The objective of this study was to find out the high yielding genotypes of mustard best suited for southern Punjab. Significant differences were observed for yield Kg ha<sup>-1</sup>, number of siliques per plant, number of branches per plant, silique's length and thousand grains weight. The genotype AZ-roya produced the highest seed yield of 2001.2 Kg ha<sup>-1</sup> followed by Khanpur raya (1838.4 Kg ha<sup>-1</sup>) and BARD-1 (1741.1 Kg ha<sup>-1</sup>). The genotype RL-18 produced the lowest seed yield of 1380.5 Kg ha<sup>-1</sup>.*

**Keywords:** Agro-climatic, Genotypes, Mustard, Yield.

## 1. Introduction

Pakistan is chronically deficient in the production of edible oil and spending huge amount of foreign exchange on its import. About 35 % of the total demand of edible oil is met through local production of oilseed crops and the remaining 65 % requirement is met through import. One disturbing aspect of this crisis issue is an annual increase in the gap between consumption and domestic production of edible oil. Edible oil is the single largest food import item, total cost involved for the import of edible oil during 2004-05 was 53.481 billion rupees, which is great burden to the national economy of the country.

In the country, after cotton, rapeseed-mustard is the second most important source of oil, but mustard is dominating species and grown on about 80 % area. The total area under rapeseed-mustard is (243.00) thousands hectares with the annual production of (205.00) thousands tonnes. In the country, the average yield of rapeseed-mustard is (843) kg ha<sup>-1</sup>, which is very low as compared to other countries of the world (Economic Survey of Pakistan 2004-05). By identifying and introducing the high yielding genotypes of rapeseed-mustard can increase the seed yield kg ha<sup>-1</sup>. So, the replacement of traditional low yielding genotypes with the improved and high yielding genotypes is very necessary for increasing the domestic production of edible oil.

The previous studies showed that yield of rapeseed-mustard can be increased by introducing and adapting the newly developed and high yielding genotypes of rapeseed-mustard (Anjum *et al*, 2005) (Ozer and Oral, 1997), (khan *et al*, 1999), and (Sharma and Manchanda 1997). In the present study ten improved genotypes of rapeseed-mustard were compared for their yield potential, resistant to insects, diseases, drought tolerance, resistant to shattering and

lodging under the existing climatic conditions of Bahawalpur region.

## 2. Materials and Methods

The proposed study was conducted on clay-loam soil at Experimentation Station of Arid Zone Research Institute (AZRI), Bahawalpur. The experiment was laid out according to Randomized Complete Block Design (RCBD), having three replications during Rabi season 2013-13 at AZRI Bahawalpur. A total number of ten genotypes of mustard, i.e. (G<sub>1</sub>) AZ-roya, (G<sub>2</sub>) Khanpur raya, (G<sub>3</sub>) RL-18, (G<sub>4</sub>) BARD-1, (G<sub>5</sub>) ZBJ-96006, (G<sub>6</sub>) RL-9, (G<sub>7</sub>) RD-7, (G<sub>8</sub>) KH-33, (G<sub>9</sub>) OR-15 and (G<sub>10</sub>) Varyana were included in this study. The crop was planted in rows spaced 45 cm and 10 cm plant-to-plant distance. There were 6 rows in each plot having row length of 10 meters. The net plot size was 2.7 x 10m<sup>2</sup>. The nitrogen and phosphorus were applied @ 75 and 50 kg ha<sup>-1</sup>. All P<sub>2</sub>O<sub>5</sub> and one third of nitrogen were applied at the time of sowing while the remaining doses of nitrogen fertilizer were applied at 1<sup>st</sup> and 2<sup>nd</sup> irrigation. A total number of three irrigations were applied to this crop. All other agronomic practices were kept normal and uniform for all treatments.

The data on following parameters i.e. days to 50% flowering completion, days to physiological maturity, Plant height (cm), number of branches per plant, number of siliques per plant, number of seeds per silique, thousand grains weight and seed yield kg ha<sup>-1</sup> were recorded at different growth stages of this crop. Days to 50% flowering completion were noted from the sowing date to the time when the 50% flowering was completed. Plant height was measured at physiological maturity of crop as an average from 10 randomly selected plants from each plot. The collected data were statistically analyzed by using Fisher's analysis of variance technique and the treatment means were compared

by using LSD test at 0.05 probability level (Steel and Torrie, 1984).

### 3. Results and Discussion

The results regarding the days to 50% flowering completion were found non-significant among different genotypes of mustard. However, the genotype BARD-1 has taken maximum days (110) to 50% flowering completion. While, minimum days (92) to 50% flowering completion were found in AZ-*raya*. Similar findings has been reported by Kukushkin-VA (1992) who reported that the duration of this phase is correlated with seed weight per fertile node, branching height, number of siliques on the central inflorescence, number of seed per silique and also lodging resistance (Table-1). The statistical analysis of the data revealed that days to physiological maturity were non-significantly affected by different genotypes of mustard. However, maximum days (155) to physiological maturity were recorded in genotype ZBJ-96006, on contrary, minimum days (127) to physiological maturity were noted in case of genotype AZ-*raya* (Table-1). In case of plant height, the analysis of results showed that non-significant differences were found among all tested genotypes of mustard under study. The highest plant height of 194.3 cm was noted in BARD-1, while the lowest plant of 163.5 cm was recorded in AZ-*raya* (Table-1). These results are contrary with Wright *et al.* (1995).

The highly significant differences were found with respect to number of branches per plant, number of siliques per plant, silique length (cm), thousands grain weight (g) and seed yield  $\text{kg ha}^{-1}$  (Table-1). These findings are against with those of Wang Guohuai *et al* (2002). The results regarding number of branches per plant revealed that, the said trait was highly significantly affected by different genotypes of mustard (Table-1). The maximum number of 12.5 branches per plant was recorded in genotype AZ-*raya*. On contrary the genotype varyana produced (9.8) minimum number of branches per plant. The genotypes Khanpur *raya* and RL-18, produced (11.4) and (10.8) number of branches per plant, remained at par with one another and statistically differed from all other genotypes. Where as the genotypes Kh-33, OR-15, ZBJ-96006, RL-9, RD-7 and BARD-1 produced (10.5), (10.5), (10.4), (10.0) and (9.9) number of branches per plant, remained at par with one another and statistically differed from all other genotypes included in this study. These results are in favor of Sharma and Manchanda (1997) and Munir and Mc Neilly (1986).

The statistical analysis of the data of number of siliques per plant as affected by different genotypes of mustard is represented in tabel-1. It is indicated that there were highly significant differences in number of siliques per plant produced by different genotypes under study. The highest number of siliques per plant (386.3) was obtained from the genotype AZ-*raya* followed by BARD-1 and Khanpur *raya* bearing (360.2) and (357.9) siliques per plant. On the other hand, the lowest number of siliques per plant (315.7) was produced by the genotype Varyana and it statistically at par with the genotype RL-18 having (325.9) numbers of siliques per plant. The genotypes OR-15, RD-71, Kh-33, RL-9 and


ZBJ-96006 produced (348.7), (340.9), (340.6), (338.6) and (330.8) number of siliques per plant, respectively and were at par with each other. This character might be attributed by genetic potential of different genotypes. These results are also supported by Wang Guohuai *at al* (2002) and Munir and Mc Neilly (1986).

Thousand grains weight contributes materially towards final yield, which varies from genotype-to-genotype. The highly significant differences among different genotypes were found regarding this character (Table-1). The maximum thousand grains weight of 5.5 g was found in genotype AZ-*raya*. In contrast, the lowest thousand grains weight of 3.2 g was noted in RL-9 and statistically at par with RL-18, RD-71 and Kh-33, all having the same vale (3.3 g) thousand grains weight. The genotypes Khanpur *raya*, BARD-1, ZBJ-96006, OR-15 and Varyana produced (4.5), (4.0), (4.0), (3.7) and (3.7) thousand grains weight (g) and remained at par with one another and statistically differ from all other treatments.

The final seed yield  $\text{kg ha}^{-1}$  is a function of the interplay of the various yield parameters. There were highly significant differences among different genotypes of mustard (Table-1). The mustard genotype AZ-*raya* produced highest yield of 2001.2  $\text{kg ha}^{-1}$ . On the contrary, the genotype RL-18 produced lowest yield of 1380.5  $\text{kg ha}^{-1}$ , this genotype did not significantly differ from the genotype Varyana that produced yield of 1442.6  $\text{kg ha}^{-1}$ . The genotypes Khanpur *raya* and BARD-1 produced yield of 1838.4 and 1741.1  $\text{kg ha}^{-1}$ , were at par with another, but statistically differ from other genotypes. The genotypes OR-15 and KH-33 produced yield of 1654.2 and 1596.4  $\text{kg ha}^{-1}$ , remained at par with another and statistically differed from all other genotypes. The genotypes RL-9, RD-71 and ZBJ-96006 produced yield of 1494.2, 1467.3 and 1445.6  $\text{kg ha}^{-1}$  respectively, were remained at par with one another but statistically differed from the other genotypes included this study. These results are in agreement with those of Abraham. T, and Lal R.B *et al.* (2002), Volliod (1982) and Santonoceto (1997). These results are in line with those of Allen and Morgan (1975) and Redmann *et al.* (1994) The differences in the yield of ten genotypes were due to best performance of genotypes under the existing agro-climatic conditions of Bahawalpur and this difference in yield might be due to well adaptation and genetic potential of genotypes of mustard.

### 4. Conclusion

In the light of results obtained, it may be concluded that the genotype AZ-*raya* performed well under existing climatic conditions of Bahawalpur. It produced maximum seed yield of 2001.2  $\text{kg ha}^{-1}$ . The results revealed that the genotype AZ-*raya* performed better incase of yield components such as number of branches per plant, number of siliques per plant and thousand grains weight. It is proved through experimentation, the genotype AZ-*raya* is a medium size, short duration and early maturing genotype of mustard. Due to that good character, the genotype is resistant to drought, lodging, shattering, insect pest and other yield reducing factors



**Figure 1:** Yield potential of different genotypes of mustard under agro-climatic conditions of Bahawalpur, Pakistan during Rabi 2012-13.

**Table 1:** Comparison of Means of Yield and Yield Components of Ten Genotypes of Mustard

Treatment/Genotypes	DFC 50%	DPM	Plant height (cm)	Branches / Plant	Siliqua's length (cm)	Siliqua/plant	1000 Grains wt. (g)	Yield kg/ha <sup>-1</sup>
AZ-roya	92.00	135	163.5	12.5 A	4.4 A	386.3A	5.5A	2001.2A
Khanpur raya	98.5	140	183.4	11.4 B	4.3AB	357.9AB	4.5B	1838.4B
RL-18	94.5	140	190.6	10.8 BC	3.7D	325.9CD	3.3C	1380.5G
BARD-1	110.00	150.4	194.3	9.9 DE	4.1ABC	360.2AB	4.0BC	1741.1BC
ZBJ-96006	105	155	182.6	10.4 CDE	4.1ABC	330.8BCD	4.0BC	1445.6FG
RL-9	105	145	182.3	10.0 DE	4.0ABCD	338.6BCD	3.2	1494.2EF
RD-71	105	155	178.9	10.5 CDE	3.9CD	340.9BCD	3.3	1467.3FG
KH-33	99.5	155	178.7	10.5 CDE	3.8CD	340.6BCD	3.3C	1596.4D
OR-15	108.4	155	186.9	9.8 E	4.0ABCD	348.7BC	3.7BC	1654.2CD
Varyana	108.5	155	185.1	0.72	4.0ABCD	315.7D	3.7BC	1442.6FG
<b>LSD</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>9.02</b>	<b>0.39</b>	<b>30.49</b>	<b>0.88</b>	<b>112.7</b>
<b>CV</b>					<b>5.73</b>	<b>12.29</b>	<b>13.44</b>	<b>4.67%</b>

NS: Non-Significant

DFC: Days to 50% flowering completion

DPM: Days to physiological maturity

CV: Coefficient of variance

Means with the same letters are not significantly different.

## References

- Abraham, T. and R.B.Lal, 2002. Sustainable and enhancement of yield potential of mustard through integrated nutrient management in legume based cropping system for the inceptisols. Cruciferae newsletter. No.24, 99-100. Dryland Research Station, Dhiansar, Bari Brahma Jammu, 181, 133, India.
- Allen, E.J. and D.G. Morgan, 1975. A quantitative comparison of the growth, development and yield of different varieties of oilseed rape. J. Agric. Sci., 85: 159-174.
- Anjum, R., M. Yousaf, M. Jahangir, M. Hussain, N. Nawaz and A. Ahmed, 2005. Adaptation and yield potential of different genotypes of rapeseed and mustard under agro-climatic conditions of Bahawalpur, Pakistan. Int. J. Agri. Biol., 7:609-611
- Anonymous, 2005. Agricultural statistics of Pakistan. Govt. of Pakistan, Ministry of Food, Agriculture and Livestock, Food, Agriculture and Livestock Division (Economic wing), Islamabad
- Khan, A., M. Rahim, and M. Khan, 1998. Yield performance of Brassica napus L. Varieties at Swat Valley bottom. Cruciferae newsletter No. 20, 91-92.
- Kiekushkin, V.A., 1992. Evaluation of some yield traits in introduced Swede rape varieties. Byullelen-Glavnogo-Botanicheskogo-Sada, 63:42-4
- Munir, M. and McNeilly, 1986. Variation in yield and yield components in six varieties of spring oilseed rape. Pak. J. Agric. Res., 7: 21-27.
- Ozer, H. and E. Oral, 1997. Phenological and yield characteristics of some rape (*Brassica napus* var *Oleifera* L.) cultivars grown at Erzurum. Turkish. J. Agric. And Forestry, 21: 319-325
- Redmann, R. E., Q. I., M. Q. and M. Belyk, 1994. Growth of transgenic and standard canola (*Brassica napus* L.) varieties in response to salinity. Can. J. Plant Sci., 74: 797-799
- Santonoceto, C., 1997. A four year research study on the development and yield of various rape (*Brassica napus* L. var. *Oleifera* DC) cultivars in Calabria. Sementi-Ellette, 43: 15-19.
- Sharma, S.K. and H.R. Manchanda, 1997. Relative performance of yellow sarson and toria grown at different salinity levels with different chloride and sulphate ratio. Indian. J. of Agri. Sci., 67:1, 1-5

- [12] Steel, R. G. D. and J. H. Torrie, 1980. Principles and procedures of Statistics, McGraw Hill Book Co., New York.
- [13] Vollioud, P., 1982. Results of variety trials with winter rape in Switzerland from 1973 to 1981. *Revue Suisse Agric.*, 14: 15-18.
- [14] Wright, P. R., J. M. Morgan, R. S. Jessop and A. Cass, 1995. Comparative adaptation of canola (*Brassica napus*) and Indian mustard (*Brassica juncea*) to soil water deficits yield and yield components. *Field Crop Res.*, 42: 1-13.

### Author Profile



**Malik Muhammad Yousaf** is Director (since 2012 continued) of Arid Zone Research Institute, Bahawalpur that is one of the prime multi-disciplinary institute of PARC. Established strong linkages with different international organizations, universities, research departments, NGOs and farming communities to boost-up the research activities of the institute. Reviewed the previous research activities and problem oriented research with special reference to introduction, selection and propagation of drought, heat and salinity tolerance in medicinal plants, pulses, cereals, oilseeds, arid horticultural plants, fodder crops, range grasses, shrubs and trees. In the capacity of the director AZRI as well as Principal Scientist, 10 projects have been developed and executed during last five years and 05 projects are currently undergoing at various implementation stages.