

- proteins. Collagen XVII as a prototype. *Matrix Biol.* 22, 299–309.
- [9] Franzke C. W., Tasanen K., Borradori L., Huotari V., Bruckner-Tuderman L. (2004) Shedding of collagen XVII/BP180. Structural motifs influence cleavage from cell surface. *J. Biol. Chem.* 279, 24521–24529.
- [10] Franzke C. W., Bruckner P., Bruckner-Tuderman L. (2005) Collagenous transmembrane proteins. Recent insights into biology and pathology. *J. Biol. Chem.* 280, 4005–4008.
- [11] Franzke C. W., Bruckner-Tuderman L., Blobel C. P. (2009) Shedding of collagen XVII/BP180 in skin depends on both ADAM10 and ADAM9. *J. Biol. Chem.* 284, 23386–23396.
- [12] Giudice G. J., Emery D. J., Zelickson B. D., Anhalt G. J., Liu Z., Diaz L. A. (1993) Bullous pemphigoid and herpes gestationis autoantibodies recognize a common noncollagenous site on the BP180 ectodomain. *J. Immunol.* 151, 5742–5750.
- [13] Grigoryan G., Keating A. E. (2008) Structural specificity in coiled-coil interactions. *Curr. Opin. Struct. Biol.* 18, 477–483.
- [14] Hirako Y., Yoshino K., Zillikens D., Owaribe K. (2003) Extracellular cleavage of bullous pemphigoid antigen 180/type XVII collagen and its involvement in hemidesmosomal disassembly. *J. Biochem.* 133, 197–206.
- [15] Hirako Y., Nishizawa Y., Sitaru C., Opitz A., Marcus K., Meyer H. E., Butt E., Owaribe K., Zillikens D. (2003) The 97-kDa (LABD97) and 120-kDa (LAD-1) fragments of bullous pemphigoid antigen 180/type XVII collagen have different N termini. *J. Invest. Dermatol.* 121, 1554–1556.
- [16] Hofmann S. C., Voith U., Schönau V., Sorokin L., Bruckner-Tuderman L., Franzke C. W. (2009) Plasmin plays a role in the *in vitro* generation of the linear IgA dermatosis antigen LAD97. *J. Invest. Dermatol.* 129, 1730–1739.
- [17] Huovila A. P., Turner A. J., Pelto-Huikko M., Kärkkäinen I., Ortiz R. M. (2005) Shedding light on ADAM metalloproteinases. *Trends Biochem. Sci.* 30, 413–422.
- [18] Kammerer R. A. (1997) α -Helical coiled-coil oligomerization domains in extracellular proteins. *Matrix Biol.* 15, 555–565.
- [19] Landschulz W. H., Johnson P. F., McKnight S. L. (1988) The leucine zipper. A hypothetical structure common to a new class of DNA-binding proteins. *Science* 240, 1759–1764.
- [20] Lin L., Betsuyaku T., Heimbach L., Li N., Rubenstein D., Shapiro S. D., An L., Giudice G. J., Diaz L. A., Senior R. M., Liu Z. (2012) Neutrophil elastase cleaves the murine hemidesmosomal protein BP180/type XVII collagen and generates degradation products that modulate experimental bullous pemphigoid. *Matrix Biol.* 31, 38–44.
- [21] Lupas A., Van Dyke M., Stock J. (1991) Predicting coiled from protein sequences. *Science* 252, 1162–1164.
- [22] Mason J. M., Arndt K. M. (2004) Coiled-coil domains. Stability, specificity, and biological implications. *Chem Bio Chem.* 5, 170–176
- [23] McAlinden A., Smith T. A., Sandell L. J., Ficheux D., Parry D. A., Hulmes D. J. (2003) α -Helical coiled-coil oligomerization domains are almost ubiquitous in the collagen superfamily. *J. Biol. Chem.* 278, 42200–42207.
- [24] McDonald L., Robertson D. H., Hurst J. L., Beynon R. J. (2005) Positional proteomics. Selective recovery and analysis of N-terminal proteolytic peptides. *Nat. Methods* 2, 955–957.
- [25] McDonald L., Beynon R. J. (2006). Preparation of amino-terminal peptides as a strategy for proteome simplification and characterization. *Nat. Protoc.* 1, 1790–1798.
- [26] McDonnell A. V., Jiang T., Keating A. E., Berger B. (2006) Paircoil2. Improved prediction of coiled coils from sequence. *Bioinformatics* 22, 356–358 .
- [27] Nebenführ A., Ritzenthaler C., Robinson D. G. (2002) Brefeldin A. Deciphering an enigmatic inhibitor of secretion. *Plant Physiol.* 130, 1102–1108.
- [28] Nishie W., Sawamura D., Goto M., Ito K., Shibaki A., McMillan J. R., Sakai K., Nakamura H., Olasz E., Yancey K. B., Akiyama M., Shimizu H. (2007) Humanization of autoantigen. *Nat. Med.* 13, 378–383.
- [29] Nishie W., Lamer S., Schlosser A., Licarete E., Franzke C. W., Hofmann S. C., Jackow J., Sitaru C., Bruckner-Tuderman L. (2010) Ectodomain shedding generates neoepitopes on collagen XVII, the major autoantigen for bullous pemphigoid. *J. Immunol.* 185, 4938–4947.
- [30] Nishie W., Kiritsi D., Nyström A., Hofmann S. C., Bruckner-Tuderman L. (2011) Dynamic interactions of epidermal collagen XVII with the extracellular matrix. Laminin 332 as a major binding partner. *Am. J. Pathol.* 179, 829.
- [31] Overall C. M., Blobel C. P. (2007) In search of partners. Linking extracellular proteases to substrates. *Nat. Rev. Mol. Cell Biol.* 8, 245–257.
- [32] Reiss K., Saftig P. (2009) The “a disintegrin and metalloprotease” (ADAM) family of sheddases. Physiological and cellular functions. *Semin. Cell Dev. Biol.* 20, 126–137.
- [33] Schäcke H., Schumann H., Hammami-Hauasli N., Raghunath M., Bruckner-Tuderman L. (1998) Two forms of collagen XVII in keratinocytes. A full-length transmembrane protein and a soluble ectodomain. *J. Biol. Chem.* 273, 25937–25943.
- [34] Snellman A., Tuomisto A., Koski A., Latvanlehto A., Pihlajaniemi T. (2007) The role of disulfide bonds and α -helical coiled coils in the biosynthesis of type XIII collagen and other collagenous transmembrane proteins. *J. Biol. Chem.* 282, 14898–14905.
- [35] Steiner H., Fluhrer R., Haass C. (2008) Intramembrane proteolysis by γ -secretase. *J. Biol. Chem.* 283, 29627–29631.
- [36] Tasanen K., Tunggal L., Chometon G., Bruckner-Tuderman L., Aumailley M. (2004) Keratinocytes from patients lacking collagen XVII display a migratory phenotype. *Am. J. Pathol.* 164, 2027–2038.
- [37] Thomas G. (2002) Furin at the cutting edge. From protein traffic to embryogenesis and disease. *Nat. Rev. Mol. Cell Biol.* 3, 753–766.
- [38] Van den Bergh F., Fu C. L., Olague-Marchan M., Giudice G. J. (2006) The NC16A domain of collagen XVII plays a role in triple helix assembly and stability. *Biochem. Biophys. Res. Commun.* 350, 1032–1037.

- [39] Veit G., Zimina E. P., Franzke C. W., Kutsch S., Siebolds U., Gordon M. K., Bruckner-Tuderman L., Koch M. (2007) Shedding of collagen XXIII is mediated by furin and depends on the plasma membrane microenvironment. *J. Biol. Chem.* 282, 27424–27435.
- [40] Wolf E., Kim P. S., Berger B. (1997) MultiCoil. A program for predicting two- and three-stranded coiled coils. *Protein Sci.* 6, 1179–1189.
- [41] Zimina E. P., Bruckner-Tuderman L., Franzke C. W. (2005) Shedding of collagen XVII ectodomain depends on plasma membrane microenvironment. *J. Biol. Chem.* 280, 34019–34024.