

Taxonomic Review of Indian Pipunculids in Comparison to Pipunculids of Other Countries

Shailendra Kumar Amogh¹, E.M. Ross², Prakash Michael³

Department of Biological Sciences, Sam Higginbottom Institute of Agriculture, Technology & Sciences, Allahabad, Uttar Pradesh, India

Department of Zoology, RLSY College, Bettiah, BRABU, Muzaffarpur, Bihar, India

Abstract: The taxonomy of Pipunculids is largely neglected in India. Brunetti, 1912, 1915 & 1923 and Subramaniam, 1922 worked on Indian Pipunculids. Since 1923 no work was done on the Indian Pipunculids till 1971. Then Hardy, 1971, 1972 & with Delfinado, 1975 worked on Indian Pipunculids. Kapoor et al., 1977, 1985 & 1987 worked on Indian Pipunculids. Again Michael, 1997 worked on Indian Pipunculids. After that no work was done on taxonomy of Indian Pipunculids. The Present review will enable researchers to work in this field and to explore new species of Pipunculids.

Keywords: Pipunculids, Taxonomy, Auchenorrhyncha, Vena spuria, Syrphids

1. Introduction

The Pipunculid is a small but very distinct fly belongs to the family Pipunculidae and order diptera with a world wide distribution. They are distinguished by their large spherical or hemispherical head which is extremely mobile and composed almost entirely of compound eyes. These flies are therefore called big headed flies. The family name is based on the genus *Pipunculus* Latreille (Latreille, 1802). According to Cammerson (1974), the name *Pipunculus* means “little pumpkin” due to the characteristic and peculiar head resembling pumpkin. The Pipunculids are relatives of Syrphids (hover flies or flower flies). They can be differentiated from Syrphids by large compound eyes that occupy most of their hemispherical head, the distinctive wing venation (no vena spuria, cell r4+5), the chitinized post spiracular plate found in the larvae. During their larval stage they are known as endoparasitoids of several families of Auchenorrhyncha (Homoptera) such as rice leaf hoppers. Over 1428 species of Pipunculids have been described worldwide (Papeet et al., 2011).

2. Early work on Pipunculidae

The first Pipunculid species were described by Bose, 1792. Meigen (1800) described the genus *Dorillas*. Latreille (1802) erected the genus *Pipunculus*. Meigen (1803) erected genus *Micronensis* Meigen. Fallen (1810) discovered genus *Cephalopus* Fallen. Walker (1834) erected genus *Chalarus* Walker. Macquart (1834) also erected a genus *Atlenevra* Macquart. Zetterstedt (1838) erected genus *Nephrocerus* Zetterstedt. Rondani (1856) discovered genus *Alloneura* Rondani and *Prothecus* Rondani. Later, Loew (1865, 1872) described 8 new species from America and Europe including *P. subvirescens*. Becker (1884) worked on Egyptian Pipunculids. Johannesson (1884) described 2 new species of *Nephrocerus* from New York. Kowarz (1887) described 14 species (of which 4 new) from Bohemia. Giard (1889) worked on *Atlenevraspuria* Meigen (Now *Chalarusspurius*). William (1893) described 6 new species of *Pipunculus* from north America. Hough (1899) described 3 new species of *Pipunculus* while studied on Pipunculids of North America. Becker (1900) published comprehensive

work on Pipunculids and described 6 new species from the Palaearctic Region. Verrall (1901) worked on British Pipunculidae of which 5 are new ones belonging to 4 genera *Chalarus*, *Verrallia*, *Nephrocerus* and *Pipunculus*. Kertesz (1903) described 6 new species of the genus *Pipunculus* from Sri Lanka.

Perkins (1905) described 26 new species from Australia and 5 new species from Hawaii. He in 1906 also described 1 new species, *P. eutrichodus* from Queensland. Becker (1907) reported 1 new species of *Pipunculus* from Tunis. De Meijere (1907) described one new species from Holland and 2 new species of *Pipunculus* from Java. Kertesz (1907) described 1 new species from Formosa and two from Africa. Becker (1908) described 2 new species of *Pipunculus* from Hungary. Strobl (1909) added *P. argyrostictus* as a new species from Monistrol. Becker (1910) added one new species from Coarsica. Perkin (1910) reported 4 new species from Hawaii. Wahlgren (1910) studied on the Swedish Pipunculidae and described 1 new species. Cresson (1911, 1912) described 9 new species of *Pipunculus* from North America.

Brunetti (1912) was first to work on Indian Pipunculids and described 12 new species from India. Kertesz (1912, 1915) described 15 new species from Formosa. Malloch (1912, 1913) described 4 new species from North America. Banks (1915) described 4 new species of *Pipunculus* and *Verrallia* from United States. Brunetti (1915) reported on new species *Pipunculusuniformis* from Shimla (H.P.), India. Johnson (1915) described one new species of *Nephrocerus* from United States. Knab (1915) also described 2 new species from United States. Johnson (1919) published a revised list of Diptera of Jamaica which included 2 new species of *Pipunculus*. Collin (1920) worked on British Pipunculidae and reported 1 new species. Becker (1921) described 6 new species from Palaearctic Region. Brunetti (1922) reported *P. annulifemur* parasitizing leaf hopper as a new species from Bangalore, India. Lamb (1922) described 4 new species from Seychelles. Brunetti (1923) published his comprehensive work on these flies “Fauna of British India” in which 19 were truly Indian (out of 24). Becker (1924) described 5 new species from Formosa. Tonnoir (1925)

Volume 4 Issue 1, January 2015

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

reported 3 new species from New Zealand. Brunetti (1927) reported 4 new species from Pahang. Shanon (1927) added 4 new species from Argentina.

Curran (1927) studied on North American Diptera and described 2 new species of Pipunculus from Canada. He in 1928 worked on African Pipunculidae. Frey (1932) studied on England Pipunculids. Curran (1934) added few new species from Africa. Sack (1935) revised the Palaearctic *Dorylaidae* and described new species. Enderlein (1936) erected 2 new genera i.e. *Cephalosphaera* for *Pipunculuscompestris* Laterrille and *Jassidophaga* for *Verrallia pilosa* Zetterstedt. Collin (1938) erected new genera and species from Nearctic and Palaearctic Region. Aczel (1939) erected 3 new tribes *Chalarine*, *Dorylaini* and *Nephrocerini* and a new genus *Tomosvaryella* or *Dorilassylvatica* Meigen from Hungary.

Hardy (1939) while working on Nearctic Pipunculidae described 4 new species and 2 new sub species of the genus *Pipunculus* and one new species of the genus *Verrallia* from U.S.A. Hardy *et al.* (1939) added 8 more new species from U.S.A. Duda (1940) described 4 new species of *Pipunculus* from Germany and also erected a new genus *Strandimyza*. Aczel (1940) erected 4 new genera i.e. *Anacephalops*, *Claraeola*, *Collinas* and *Eudorylas* and also described a new species of *Cephalosphaera* from Europe. Hardy (1940) described 7 new species of *Tomosvaryella* and 1 new species of *Dorylas* from U.S.A. Brooks and Jagges (1941) worked on the Pipunculidae of Iowa, USA. Collin (1941) worked on Pipunculidae of Ussuri region of USSR describing 4 new species of *Pipunculus* and 1 of *Verallia*. Hardy (1943) worked on Nearctic *Dorilaidae* and described 27 new species and a new genus *Allomethus*.

Rapp (1943) described 2 new species from Quebec and 1 from New Jersey. He again worked on Pipunculidae of New Jersey in 1944. In 1945, he studied the types of genera & sub genera of Pipunculids. Rapp & Cooper (1943) published a catalogue of African Pipunculids.

Aczel (1944) revised the Palaearctic species of *Tomosvaryella* giving new species. Aczel (1948) published a monograph of *Dorilaidae* of Europe. Aczel (1952) brought a catalogue of Pipunculids of Neotropical region.

Collin (1956) worked on Pipunculids describing many new species of *Alloneura*, *Cephalops* and *Pipunculus* from Palaearctic region. He again in 1958, described 2 new species from Yugoslavia (1 of *Alloneura* & 1 of *Cephalops*).

Koizumi (1959) described 2 new species of *Eudorylas* and 1 of *Tomosvaryella* from Japan. He again in 1960 described 2 new species of *Dorilaidae* from Japan.

Coe (1966) published a handbook for the identification of British insects included 13 new species of Pipunculidae. He again in 1966 worked on British species of *Chalarus* & *Verallia* with 6 new species. In 1969, he described some Pipunculids from Southern Spain. Whittkar (1969) described life cycle of *Verallia aucta* and *Verallia setosa*.

Hardy (1971) published a review of the Oriental Pipunculids in which 4 species from India i.e. *T. oryzaeatora*, *T. subvirescens*, *P. Eudorylas Javaensis* and *P. (E.) orientalis*. He in 1972 published a monograph on the Oriental Pipunculids in which he described 6 species from India i.e. *P. (Cephalops) subaeneus* Brunetti, *P. (E.) fillicornis* Brunetti, *P. (E.) macropygus* de meijere, *P. (E.) uniformis* Brunetti, *T. limpidipennis* (Brunetti) and *T. nitens* (Brunetti). Bankowska (1972) reviewed the Polish species of Pipunculidae of which 13 were new species. Hardy and Delfinado (1975) brought out a catalogue of the Oriental Pipunculidae of which 23 were from India. Lyenborg (1975) worked on the Danish Pipunculidae and described 2 new species.

Kapooret *et al.* (1977) added 2 new species namely *P. (E.) ranikhetiensis* and *P. (E.) kumaonensis* from Kumaon hills of U.P., India. Rald (1977) worked on the Pipunculidae of Denmark and described some rare species along with a checklist.

Albrecht (1979) described 8 new species of *Dorylomorpha* from Finland. Chandra (1979) worked on the taxonomy of Phillipines. Mc. Lean (1979) described a new Pipunculid, *Nephrocerus flavicornis* from Norflox, Britain.

Kozanek (1981) while working on the Pipunculidae of Czechoslovakia, described 5 new species of *Pipunculus* and *Nephrocerus*. Shatalkin (1981) described 1 new species of *P. claraeola* from USSR.

Kapoor *et al.* (1985) again described 1 new species of the genus *Dorylomorpha* Aczel and 1 new sub genus *Clareosphaera* from India. Jervis (1986) added 2 new species of the genus *Chalarus* Walker (*C. trilineatus* and *C. orientalis*) from Burma. Speight (1986) found *Dorylomorpha maculata* as new to Ireland. Rafael (1986) described a new genus *parapipunculus* of Pipunculidae from Nearctic region & Mexico. He again in 1986 described a new genus *Amanzunculus* from Amazon Basin.

Kapooret *et al.* (1987) described 2 new sub genera from India.

Kuznetsov (1990) described new species of Pipunculidae of Palaearctic region. He again in 1991 described a new species of the genus *Pipunculus* from Palaearctic region. In 1994, he described new Palaearctic species of *Tomosvaryella* Aczel.

Morakoteet *et al.* (1990) studied on the Japanese Pipunculidae. Jervis (1992) worked on taxonomic revision of genus *Chalarus* Walker from European fauna. De Meyer (1993) revised Afrotropical species of *Tomosvaryella* Aczel, 1939. Ale Rocha *et al.* (1995) worked on genus *Tomosvaryella* Aczel from Mexico, Central America.

Skevington & Marshall (1998) studied on systematic of New World Pipunculus. Skevington (1999) described *Cephalosphaera* Enderlein, a genus of Pipunculidae new for Australia. Skevington (2001) worked on revision of Australian Pipunculidae. Skevington (2002) studied on phylogenetic revision of Australian members of the *Allomethus* genus. Skevington (2003) worked on revision of Australian *Eudorylas* Aczel. Moya - Raygoza *et al.* (2004) reported *Eudorylas spinosus* (Hardy) parasitoids of corn leaf

hoppers from Mexico. Koenig & Young (2007) reported genus *Nephrocerus* attack adult crane flies (Tipulidae). Virlaet *et al.* (2009) reported *Eudorylasschreiteri* (Shannon) as a parasitoid of corn leaf hopper.

3. Conclusion

From above review on Pipunculids it is clear that there is little work done in India in comparison to rest of the world. Brunetti (1912) was the first to work on the Indian Pipunculids and described 12 new species from India. Again in 1915, he described 1 new species *Pipunculusuniformis* from Shimla, India. He published his comprehensive work (1923) on these flies in "Fauna of British India" including 24 species. Out of these 24 species 19 are from India. Since 1923 no work was done on the Indian Pipunculids till 1971.

Hardy (1971) published a review of the Oriental Pipunculids included 4 species from India. In 1972, he published a Monograph on the Oriental Pipunculids included 6 species from India. Hardy *et al.* (1975) published a catalogue of the Oriental Pipunculidae including 131 species of which 23 were from India.

Kapooret *et al.* (1977, 1985) described 3 new species from Kumaon hills of U.P., India. Later, Kapooret *et al.* (1987) published A Comprehensive Monograph on Indian Pipunculids including 91 species from India. Taxonomy of these flies are largely neglected in India. So, there is a need for detailed systematic work on these flies in India.

References

- [1] Aczel, M. 1939a. Das system der family Dorylaidae. Zool. Anz., 125: 15-23.
- [2] Aczel, M. 1939b. Die Untergutting Dorylomorpha m. von. Tomosvaryella m. Zool. Anz., 125(3/4): 49-69
- [3] Aczel, M. 1939c. Beckeriaspannonicus, eine neue Gattung und Art des Dorylaiden (Diptera). Dorylaidenstudien 111; Zool. Anz., 126 (718): 191-192.
- [4] Aczel, M. 1940. Vorarbeiten zu einer Monographie der Dorylaiden. Dorylaidenstudien. V. Zool. Anz.; 132 (7/8): 149-169.
- [5] Aczel, M. 1944. Die Gattung Tomosvaryella Aczel (Diptera) Dorylaidenstudien V. Ann. Nat. Hist. Mus. Natl. Hung., 37: 75-130.
- [6] Aczel, M. 1948. Grundlagen einer Monographie der Dorylaiden, Acta Zool. Lilloana del Inst. Mequel Lillo., 6: 5-168.
- [7] Aczel, M. 1952. Catalogo de la familia Dorylaidae (Pipunculidae) de la Region Neotropical. Rev. Soc. Ent. Argentina, 15: 237 - 251.
- [8] Albrecht, A. 1979a. Description of new Dorylomorpha species from Europe (Diptera: Pipunculidae). Entomol. Scand., 10(3): 211-218.
- [9] Albrecht, A. 1979b. Dorylomorphafennica sp. n., a new pipunculid species (Diptera) from Finland. Notulae Entomol., 59: 15-17.
- [10] Alae Rocha, R & Rafael, J. A. 1995. Tomosvaryella Aczel do Mexico, America Central e Antilhas (Diptera: Pipunculidae). Revista Brasileira de Zoologia, 12: 407-427.
- [11] Bankowska, R. 1972. A review of polish species of the family Pipunculidae (Diptera). Fragm. Fean. Wers., 8(14): 257-273.
- [12] Banks, N. 1915. Notes and descriptions of Pipunculidae. Psychae. Boston, 22(2): 166-170.
- [13] Becker, T. 1884. Pipunculidae, Characters of Egyptian species. Mt. Bus. Berlin, 2: 93-98.
- [14] Becker, T. 1900. Studien- V. Pipunculidae. Berlin. Ent. Zeitschr., 45: 215-252.
- [15] Becker, T. 1907. Diptera: Brachycera. Berl. Ent. Zeit 52: 163.
- [16] Becker, T. 1908. The Diptera due nova ex. Hungaria. Ann. Hist. Nat. Mus. Hung. Budapest, 6: 320.
- [17] Becker, T. 1910. Dipterologischae Sammelrensenach Corsica. Berl. Ent. Zeit., 55: 635.
- [18] Becker, T. 1921. Neue Dipteren Fleiner Sammlung. Pipunculidae. Wein., Ent. Zeit., 38: 123-132.
- [19] Becker, T. 1924. H. Sauter's Formosa- Ausbeute. Pipunculidae (Dorylaidae: Diptera). Ent. Mitt., 13 (1): 4-18.
- [20] Brooks, K. and Jagges, H. E. 1941. Preliminary list of Pipunculidae of Iowa. Proc. Iowa Acad. Sci., 48: 437-445.
- [21] Brunetti, E. 1912. New Oriental Diptera. Rec. Indian Mus., 7(5): 445-513.
- [22] Brunetti, E. 1915. Diptera of Shimla District. Rec. Indian Mus., VIII. 13: 82.
- [23] Brunetti, E. 1922. A new Pipunculidae parasitic on leaf hoppers in India. Bull. Ent. Res., 12: 469.
- [24] Brunetti, E. 1923. Pipunculidae, Syrphidae, Conopidae, Oestridae, Fauna of British India, including Ceylon & Burma. Diptera. Taylor & Francis London, 13: 1-123.
- [25] Brunetti, E. 1927. J. Fed. Malay St. Mus., 13: 302.
- [26] Chandra, G. 1979. Taxonomy and the bionomics of the insect parasites of rice leaf hoppers and plant hoppers in the Philippines and their importance in natural biological control. Phillip. Ent., 4(3): 119-139.
- [27] Coe, R. L. 1966a. Handbook for the identification of British insects (Diptera: Pipunculidae). R. Ent. Soc. London., 10 (2C): 1-83.
- [28] Some British species of Chalarus and Verrallia (Diptera: Pipunculidae). Proc. R. Ent. Soc. London (B), 35 (11-12): 149-160.
- [29] Coe, R. L. 1969. Some Pipunculidae Diptera from Southern Spain with description of a New species. Ent. Meddel., 37 (11): 1-8.
- [30] Collin, J. E. 1920. The British species of Sylvanticus group of Pipunculids. Ent. Mon. Mag., 56: 270-275.
- [31] Collin, J. E. 1938. A new Pipunculus (Diptera) Sicily. Proc. R. Ent. Soc. London (B), 7(1): 81-83.
- [32] Collin, J. E. 1941. Some Pipunculidae and Empididae from Ussuri region of Far Eastern border of the USSR. Proc. R. Ent. Soc. (B), 10: 218-248.
- [33] Collin, J. E. 1956. Scandinavian Pipunculidae. Opusc. Ent., 21: 149-161.
- [34] Cresson, E. T. 1911. Studies in North American Dipterology. Pipunculidae. Trans. Amer. Ent. Soc., 36: 291-329.

- [35] **Cression, E. T.** 1912. Studies of some Pipunculidae from the Eastern United States. Ent. News, Philadelphia, 23 : 452-456.
- [36] **Curran, C. H.** 1927. Some New North American Diptera. Cannad. Ent. Orillia., 59 : 290-303.
- [37] **Curran, C. H.** 1927. The Diptera of Kentabo, Bartica District, British Guiana with description of new species. Bull. Amer. Nat. Hist. Mus., 66(3) : 287-532.
- [38] **De Meijere, J. C. H.** 1907. Studienubersudostasiatische Dipteran-1. Tijds. Ent., 50 : 259-264.
- [39] **De Meyer, M.** 1993. A revision of Afrotropical species of Tomosvaryella Aczel. Ama. Nat. Mus. 34 : 43-101.
- [40] **Duda,** O. 1940. Neueoderungen genugendbekannete Zweiflugler der Palearktischen Region. Folio Zool. Hydrobiol. Riga., 10 : 214-236.
- [41] **Enderlein, G.** 1936. Zweifluger, Diptera in Brohmer, P. G. Ehrmann & G. Ulmer : Die Tierwelt Mitteleuropas 6, Insecten, 3 : 128-129.
- [42] **Fallen, C. F.** 1810. Specimen entomologicum novam Dipterosum disponendum methodo mexhibens, 10 : 26 pp.
- [43] **Frey, R.** 1932. Neue Diptera Brachycera aus Finnland und argenzenden Landern. II Notnl. Ent., 12 : 81-85.
- [44] **Giard, A.** 1889. Sur la castration parasitaire des Typhlocybapernnalarvid hymenopterae (Aphelopus melaleucus Dalm) at par une larve de Diptera (Atelenerraspuria Meing.) Comptes Rendus Hebdomadaires de L. Academie Des. Seances. Paris, 109 : 708-710.
- [45] **Hardy, D. E.** 1939. New Nearctic Pipunculidae (Diptera). J. Kansas. Ent. Soc., 12(1) : 16-25.
- [46] **Hardy, D. E.** 1940. Dorylaidae Notes and descriptions (Pipunculidae : Diptera). J. Kansas. Ent. Soc., 13 : 101-114.
- [47] **Hardy, D. E.** 1943. A revision of Nearctic Dorilaidae (Pipunculidae). Univ. of Kansas Sci. Bull., 29 : 1-231.
- [48] **Hardy, D. E.** 1971. Pipunculidae (Diptera) parasitic on rice leaf hoppers in the Oriental Region. Proc. Hawaii Entomol. Soc., 21 (1) : 79-91.
- [49] **Hardy, D. E.** 1972b. Studies on Oriental Pipunculidae (Diptera). Oriental Insect Supplement, nr-2 : 1-76.
- [50] **Hardy, D. E.** 1975. Family Pipunculidae, In : A Catalogue of Diptera of the Oriental Region II. Delfinado, M. D. & D. E. Hardy eds. Hawaii Univ. Press, 296-306.
- [51] **Hardy, D. E.** 1980. Family Pipunculidae, In : A Catalogue of Diptera of the Afrotropical Region. Publication Br. Mus. Nat. Hist. nr-821, 37 : 483-487.
- [52] **Hardy, D. E. & Knowton, G. F.** 1939a. New & little known Pipunculidae (Diptera). Cannad. Ent., 71 : 87-91.
- [53] **Hardy, D. E. & Knowton, G. F.** 1939b. New & little known Western Pipunculidae (Diptera). Ann. Ent. Soc. Amer. Columbus, 32 : 113-124.
- [54] **Hough, G.** 1899. Pipunculidae of North America. Proc. Bost. Soc. Nat. Hist. 29 : 77-86.
- [55] **Jervis, M. A.** 1986. Two new species of Chalarus Walker (Diptera : Pipunculidae) from Burma. Mitt. Schweiz. Entomol. Ges., 58 (3-4) : 435-440.
- [56] **Jervis, M. A.** 1992. A taxonomic revision of the Pipunculid fly genus Chalarus Walker, with particular reference to the European Fauna. Zool. J. of Linn. Soc. 105 : 243-352.
- [57] **Johanneson, D. A.** 1884. Two new species of family Pipunculidae (Diptera). Ent. News. Philadelphia, 14 : 107-108.
- [58] **Johnson, C. W.** 1915. A new species of the genus Nephrocerus, Cannad. Ent., 47 : 54-56.
- [59] **Johnson, C. W.** 1919. A revised list of the Diptera of Jamaica. Bull. Amer. Mus. Nat. Hist., 41(8) : 421-449.
- [60] **Kapoor, V. C. Agarwal, M. L. and Grewal, J. S.** 1977. Two new species of Pipunculidae (Diptera : Pipunculidae) from Ranikhet, Kumaon hills, India. Bull. Ent., 18 : 74-77.
- [61] **Kapoor, V. C., Grewal, J. S. and Sharma, S. K.** 1987. Indian Pipunculids (Diptera : Pipunculidae). A Comprehensive Monograph. Atlantic Publ. & Distr. New Delhi, 201 pp.
- [62] **Kertesz, K.** 1903. Die gattung Pipunculus. Arten Sudasiens, Ann. Hist. Nat. Mus., Natl. Hung., 1 : 465-471.
- [63] **Kertesz, K.** 1907. Vier neue Pipunculus Arten. Ann. Hist. Nat. Mus. Natl. Hung. Budapest., 5 : 579-583.
- [64] **Kertesz, K.** 1912. Sauter's Formosa-Ausbeute Dorylaidae (Diptera). Ann. Hist. Nat. Mus. Natl. Hung., 10 : 285-299.
- [65] **Kertesz, K.** 1915. Contribution to the knowledge of Dorylaidae. Ann. Mus. Nat. Hung., 13 : 386-392.
- [66] **Knob, F.** 1915. Two new species of Pipunculus from Washington. D. C. Proc. Biol. Soc. Wash., 28 : 83-85.
- [67] **Koenig, D. P. & Young, C. W.** 2007. First observation of parasitic relations between big headed flies, Nephrocerus Zetterstedt (Diptera : Pipunculidae) and Crane flies, Tupila Linnaeus (Diptera : Tipulidae : Tipulinae), with larval and puparial description for the genus Nephrocerus. Proc. Entomol. Soc. Wash., 109 : 52-65.
- [68] **Koizumi, K.** 1959. On four Dorylaidae parasites of green rice leaf hoppers Nephrotettix cincticeps Uhler (Diptera). Scient. Rep. Fac. Agric. Okayama Univ., 13 : 37-45.
- [69] **Koizumi, K.** 1960. A new Dorylaeid parasite of zigzag striped leaf hopper Inazumadorsalis & notes on other paddy field inhabiting Dorilaidae. Sci. Rep. Fac. Agric. Okayama Univ., 16 : 33-42.
- [70] **Kowarz, G.** 1887. Beitrage zu einem Verzeichnisse der Dipteren Bohmens 6. Conopidae, Pipunculidae. Wlen. Ent. Zertg., 6 : 146-154.
- [71] **Kozanek, M.** 1981a. Genus Pipunculus (Dip. : Pip.) in Czechoslovakia. Ann. Zool. Bot., 142 : 1-16.
- [72] **Kozanek, M.** 1981b. Description of Pipunculus lichtwardti new sp. (Dip. : Pip.) and redescription of P. omissinervis. Biologia. (Bratisl.), 36(8) : 605-610.
- [73] **Kozanek, M.** 1981c. The genus Nephrocerus (Dip. : Pip.) in Czechoslovakia. Biologia (Bratisl.), 36(5) : 395-396.
- [74] **Lamb, C. G.** 1922. Diptera, Asilidae, Scenopinidae, Dolichopodidae, Pipunculidae, Syrphidae. Trans. Lin. Soc. Lond., 18 (1) : 36-416. Pipunculidae 372-378.
- [75] **Latreille, P. A.** 1902. Histoire naturelle generale et particulière des crustaces et des Insectes. 3 : 1-468.

- [76] **Loew, H.** 1865. Diptera Americae septentrionalis indigena Centuria sexta. Berl. Ent. Zeitschr., 9 : 127-176.
- [77] **Loew, H.** 1872. Diptera Americae septentrionalis indigena Centuria sexta. Berl. Ent. Zeitschr., 16 : 49-124.
- [78] **Lyenborg, L.** 1975. Notes on Danish Pipunculidae (Dipt. : Pip.). Ent. Medd., 43 (3) : 136.
- [79] **Macquart, J.** 1834. Recueil des travaux de la societé d'amateurs des Sciences de l' Agriculture et des Arts de Litte. Insectes du Nord de France (1830) : 356.
- [80] **Malloch, J. R.** 1912. Three new spp of Pipunculidae (Dipt.) from Panama, Washington, D. C. Smithsonian Inst. Misc. Collect., 60 (1) : 1-4, pb nr-2141.
- [81] **Malloch, J. R.** 1913. New American Dipterous insects of the fam. Pipunculidae. Proc. U. S. N. M., 43 : 291.
- [82] **Mc Lean, I. F. G.** 1979. Notes on Diptera taken in 1976. Entomol. Rec. J. Var., 9 (2-3) : 71-72.
- [83] **Meigen, J. W.** 1800. Nouvelle classification des mouches à deuxarten (Diptera) dapresun plan. tout Nouveau. pp. 40.
- [84] **Meigen, J.** 1803. VersucheinerneuenGattungsEintheilung des europaischenZweiflugeligenInsekten. Mag. Insectkten., 2 : 259-286.
- [85] **Moya-Raygoza, G., Kathirithamby, J. and Larsen, K. J.** 2004. Dry season parasitoids of adult corn leaf hoppers (Hemiptera : Cicadellidae) on irrigated maize in Mexico. Can. Entomol. 136 : 119-127.
- [86] **Perkins, R. C. L.** 1905. Leaf hoppers and their natural enemies. Pt. 4, Pipunculidae. Bull. Div. Ent. Hawaiian Sug. Plrs. Ass. Exp. Stn., 1 (4) : 123-157.
- [87] **Rafael, J. A.** 1986a. Parapipunculus, a new genus of Pipunculidae (Diptera) from the Nearctic Region and Mexico. Revista Brasileira de Entomologia, 30 : 421-424.
- [88] **Rafael, J. A.** 1986b. Amazunculus, a new genus of Pipunculid from the Amazon Basin (Diptera : Pipunculidae). Amazoniana, 10 : 15-19.
- [89] **Rald, E.** 1977. New & rare Diptera Brachycera from Denmark. Entomol. Medd., 44 (1) : 23-27, 1976.
- [90] **Rapp, W. F.** 1943. Some new North American Pipunculidae (Diptera). Ent. News, Philadelphia, 54 : 222-224.
- [91] **Rapp, W. F. & Cooper, J. L.** 1943. Preliminary catalogue of African Pipunculidae. Ann. Mag. Nat. Hist., 12 (11) : 128-129.
- [92] **Rondani, C.** 1856. Genera Italicaordinesdipterosumordinatimdispositaetdistincta et in familias et stirpes aggregata. Dipterologiae Italica Prodromus, 1 : 1-228, Parmae.
- [93] **Sack, P.** 1935. Dorylaidae (Pipunculidae). In Lindner Die Fliegen des Palaearktischen Region. 4 (4) : 1-57.
- [94] **Shannon, R. C.** 1927. Some new Diptera from Argentina. Rev. Soc. Ent. Arg., 1 (4) : 31-42.
- [95] **Shatalkin, A. I.** 1981. Claraeolacrassula sp. n. (Pip.) and strongylophthalmyiararicornis sp. n. (Strongylophthalmyiidae), new peculiar Diptera from USSR. Zoologischeskizl., 60 (5) : 791-793.
- [96] **Skevington, J. H.** 1999. Cephalosphaera Enderlein, a genus of Pipunculidae (Diptera) new for Australia, with description of four new species. Australian J. of Entomol., 38 : 247-256.
- [97] **Skevington, J. H.** 2001. Revision of Australian Clistoabdominalis (Dip. : Pipunculidae). Invertebrate Taxonomy., 15 : 695-761.
- [98] **Skevington, J. H.** 2002. Phylogenetic revision of Australian members of the Allomethus genus group (Diptera : Pipunculidae). Insects Systematics & Evolution., 33 : 133-161.
- [99] **Skevington, J. H.** 2003. Revision of Australian Eudorylas Aczel (Diptera : Pipunculidae). Studiadipterologica., 9 (2) (2002) : 621-672.
- [100] **Skevington, J. H. and Marshall, S. A.** 1998. Systematics of New World Pipunculus (Diptera : Pipunculidae). Thomas Say Publication in Entomology Monographs Entomological Society of America, Lanhan, Maryland. 201p.
- [101] **Speight, M. C. D.** 1986. Asafhidon curtum, Dorylomorphamaculate, Selatosomus melencholicus & Syntormon miki, insects new to Ireland. Ir. Nat. J., 22 (1) : 20-23.
- [102] **Strobl.** 1909. Die Diptera Von Steimark Naehtrag. Craz. Mitt. Natn. Ver. Steimark., 45 : 45-293.
- [103] **Stubbs, A. E.** 1980. The largest Pipunculid in the land : Nephrocerus Scutellata (Macquart, 1834) (Dipt. : Pip.) new to Britain, with observation on its behavior in Greece. Proc. Trans. Br. Entomol. Natl. Hist. Soc., 13 (1-2) : 46-48.
- [104] **Tonnoir, A. L.** 1925. New Zealand Pipunculidae (Diptera). Rec. Canterbury Mus., 2 : 313-316.
- [105] **Verall, G. H.** 1901. British flies of Platopezidae, Pipunculidae and Syrphidae, 3 : 1-691.
- [106] **Virla, E. G., Moya - Raygoza, G. & Rafael, J. A.** 2009. First record of Eudorylasschreiteri (Shannon) (Diptera : Pipunculidae) as a parasitoid of corn leaf hopper (Hemiptera : Cicadellidae) in Argentina, with a table of Pipunculid – host associations in the Neotropical Region. Neotropical Entomology., 38 : 152-154.
- [107] **Walker, F.** 1834. Some observation on the British species of Pipunculidae. Ent. Mag., 2 : 262-270.
- [108] **William, S. W.** 1893. In Riley, C. W. Report on a small collection of insects made during the Death Valley Expedition. North American Fauna., 7 : 255.
- [109] **Zetterstedt, J. W.** 1838. Insect Lapponica, 8 : pp 578.