

# Comparison of Clinico-Radiological Profile Of New Smear Positive Pulmonary tuberculosis Cases Among Adults and Elderly People

Arun Babu.V<sup>\*1</sup>, Subramanian.S<sup>2</sup>, Meenakshi.N<sup>3</sup>, Ragulan.R<sup>4</sup>, Viswambhar.V<sup>5</sup>, Apar Jindal<sup>6</sup>

<sup>1</sup>Post graduate student, Department of Respiratory Medicine, Chettinad Hospital and Research Institute, Chennai, India

<sup>2</sup>Associate Professor, Department of Respiratory Medicine, Chettinad Hospital and Research Institute, Chennai, India

<sup>3</sup>Head and Professor, Department of Respiratory Medicine, Chettinad Hospital and Research Institute, Chennai, India

<sup>4</sup>Assistant Professor, Department of Respiratory Medicine, Chettinad Hospital and Research Institute, Chennai, India

<sup>5</sup>Assistant Professor, Department of Respiratory Medicine, Chettinad Hospital and Research Institute, Chennai, India

<sup>6</sup>Post graduate student, Department of Respiratory Medicine, Chettinad Hospital and Research Institute, Chennai, India

**Abstract:** *Introduction:* Elderly age group is one of the risk factor for tuberculosis. The presentation of tuberculosis in elderly may vary at times from young adults. Such atypical presentation may delay the diagnosis in elderly patients. *Aim* of the study is to compare the clinico-radiological pattern of smear positive pulmonary tuberculosis in the adults (18-50 years) and elderly (> 50 years). *Methods:* This prospective observational study was carried out on all new smear-positive pulmonary tuberculosis patients in the Department of pulmonary medicine, Chettinad Hospital and Research Institute, Chennai. *Results:* Males are predominantly affected in both the groups. Loss of appetite, loss of weight, breathlessness and chest pain, were more frequently reported in the elderly group (age > 50 years) whereas hemoptysis was found significantly higher in younger patients group. Bacillary load was higher in elderly. Radiologically bilateral involvement, lower zone predominance and far advanced disease were noted in elderly. *Conclusion:* Elderly patients with tuberculosis have more atypical clinico radiological presentation. A high degree of suspicion need be maintained by the physician to aid early diagnosis of tuberculosis in the elderly.

**Keywords:** Pulmonary tuberculosis, Elderly, Adults, Atypical Presentation

## 1. Introduction

Tuberculosis is an endemic disease in India that progresses from a focal infection to systemic infection caused by Mycobacterium tuberculosis. Most commonly, tuberculosis spreads from person to person by airborne transmission of droplet nuclei. Despite a relatively low transmission rate compared with other contagious diseases and the existence of effective control programmes, still tuberculosis remains a growing public health problem. Approximately one-third of the world's population is infected with tuberculosis. Although occurrence of atypical clinical presentation and chest radiographic findings in the geriatric population is known, a large number of studies conducted overseas report similarity in clinical, microbiological and radiological presentation amongst both adults and elderly. [1] Review of literature reveals a serious deficiency of Indian medical literature describing the problems of diagnosis and management of pulmonary tuberculosis in the elderly population.

India is considered as "an ageing nation" with 7.7% of its population being more than 60 years old. Pulmonary tuberculosis presents varyingly amongst adults and the elderly [2] and need to be evaluated differently. Probably the higher morbidity and mortality reported in the elderly population is due to non-recognition of these subtle differences in presentation thereby, delaying establishment of the diagnosis and early institution of

treatment.[3] Tuberculosis constitutes a major disease burden amongst the elderly and is expected to increase continually in the geriatric community,[4] therefore, this study was undertaken to study the clinic-radiological variability of smear positive pulmonary tuberculosis in elderly as compared to young patients.

## 2. Aims and Objectives

To compare the clinico-radiological presentation in adults (18-50 years) and elderly (> 50 years) patients with smear positive pulmonary tuberculosis.

## 3. Methodology

The study was conducted at the Department of Respiratory Medicine, Chettinad Hospital and Research Institute, Chennai in outpatient and inpatients of Tuberculosis. It is a prospective observational study approved by research and ethical committee. 150 smear positive pulmonary tuberculosis patients were included in the study.

### Inclusion criteria

1. Subjects diagnosed as new smear positive pulmonary Tuberculosis.
2. Subjects in age group 18 - 70 years

**Exclusion criteria**

1. Extra pulmonary tuberculosis
2. Seropositive (HIV/AIDS)
3. Chronic liver/renal failure
4. Chronic steroid/immunosuppressive therapy
5. Diagnosed Connective tissue disorders
6. Malignancy
7. Pregnancy

Informed written consent was obtained from all subjects before enrollment.

The Diagnosis of Tuberculosis was based on detailed history, clinical examination, sputum microscopy for acid fast bacilli and chest radiography. The included subjects were divided into two groups-adults (age 18-50yrs) and elderly (age>50yrs).

All chest x-rays were reviewed for presence or absence of cavities, zones involved and extent of disease. Extent of lesions was classified into 3 categories namely "(i) minimal lesion: disease with a combined area less than that of the right upper lobe, (ii) moderate: disease with a combined area less than that of the right lung but more than that of the right upper lobe and (iii) advanced: disease with a combined area more than that of the right lung." [5]

At the end of the study, the data collected was subject to statistical analysis.

**4. Results**

A total of 150 (63 elderly and 87 adult) patients of new smear positive pulmonary tuberculosis were included in the study. The data thus collected was subject to appropriate statistical analysis, and results are presented below.

**Table 1: Gender Distribution**

Gender	Smear positive TB (n=150)	
	Adults (n=87)	Elderly (n=63)
Male	57 (65.5%)	48 (76.2%)
Female	30 (34.5%)	15 (23.8%)

Table 1 compares the gender distribution in both the groups. Male predominance was seen in both the groups.

**Table 1: Symptom Characteristics**

Parameter	Smear positive TB (n=150)		p Value
	Adults (n=87)	Elderly (n=63)	
Cough	79(90.8%)	59(93.7%)	0.526
Expectoration	85(97.7%)	62(98.4%)	0.759
Breathlessness	43(49.4%)	55(87.3%)	0.000
Chest pain	15(17.2%)	20(31.7%)	0.038
Hemoptysis	30(34.5%)	10(15.9%)	0.011
Fever	69(79.3%)	52(82.5%)	0.621
Loss of appetite	66(75.9%)	57(90.5%)	0.021
Loss of weight	62(71.3%)	57(90.5%)	0.004

**Table 2** compares the symptom characteristics among elderly and adults. Breathlessness, chest pain, loss of appetite and loss of weight were found significantly higher in elderly compared to adults, while hemoptysis was found significantly higher in adults ( $p < 0.01$ ). However cough and/or expectoration, and fever did not significantly differ between the groups

**Table 2: Bacillary load**

AFB	Smear positive TB (n=150)		p Value
	Adults (n=87)	Elderly (n=63)	
Scanty	9(10.3%)	5(7.9%)	0.030
1+	32(36.8%)	17(27.0%)	
2+	26(29.9%)	12(19.0%)	
3+	20(23.0%)	20(46.0%)	

**Table 3** shows the comparison of bacillary load between the two groups. It is observed that the high bacillary load (3+) was found to be more in elderly than adults ( $p < 0.03$ ). Within the elderly group 46% reported 3+ smear positivity.

**5. Comparison of Radiology Characteristics**

**Table 4.1: Comparison of side preference**

Side	Smear positive TB (n=150)		p Value
	Adults (n=87)	Elderly (n=63)	
Right	30(34.5%)	24(38.1%)	0.21
Left	29(33.3%)	13(20.6%)	
Bilateral	28(32.2%)	26(41.3%)	

**Table 4.2: Comparison of lesions**

Lesions	Smear positive TB (n=150)		p Value
	Adults (n=87)	Elderly (n=63)	
Fibrosis	14(16.1%)	16(25.4%)	0.160
Cavity	41(47.1%)	26(41.3%)	0.476
Alveolar opacity	67(77.0%)	45(71.4%)	0.438
Consolidation	33(37.9%)	23(36.5%)	0.859

**Table 4.3: Comparison of extent of lesions**

Extent of lesion by chest x ray	Smear positive TB (n=150)		p Value
	Adults (n=87)	Elderly (n=63)	
Minimal	40(46.0%)	10(15.9%)	0.000
Moderately advanced	33(37.9%)	28(44.4%)	
Far advanced	14(16.1%)	25(39.7%)	

**Table 4.1-4.4** compares the radiological characteristics of both the groups. Bilateral involvement was found more in elderly group compared to adults ( $p = 0.21$ ). Cavity, alveolar opacity and consolidation were seen predominantly in adults whereas fibrosis was more common in the elderly group. Lower zone involvement and far advanced disease were significantly higher ( $p < 0.001$ ) in elderly as compared to adults.

**Table 4.4: Comparison of zone preference**

Lung Zone		Smear positive TB (n=150)		p Value
		Adults (n=87)	Elderly (n=63)	
UZ	Present	61(70.1%)	37(58.7%)	0.148
	Absent	26(29.9%)	26(41.3%)	
MZ	Present	62 (71.3%)	47(74.6%)	0.651
	Absent	25(28.7%)	16(25.4%)	
LZ	Present	15(17.2%)	27(42.9%)	0.001
	Absent	72(82.8%)	36(57.1%)	

**6. Discussion**

**1. Sex distribution**

In our study, in both groups males were predominantly affected similar to the observations of Morris et al [6]. Tripathy and Kar [7] in their study reported 78% of their patients to be males. Meta-analysis of 12 studies done by

Carlos Perez-Guzma [8] showed male preponderance. Similar observations were reported by Tan KK et al [9].

## 2. Comparison of symptomatology

In our study, breathlessness, loss of appetite, loss of weight and chest pain was found significantly higher in elderly and haemoptysis was significantly higher in adults. The most common presenting complaint in both the groups was cough with (or) without expectoration. Jagdish Rawat et al [10] compared the clinico-radiological pattern of pulmonary tuberculosis in adults and elderly patients and observed that "hemoptysis (29.5% vs. 6%), fever (95.4% vs. 76%) and night sweats (54.5% vs. 18.0%) were significantly higher in the adults". Anand Patel et al [11] from Gujarat also reported that elderly had less incidence of fever, chest pain, haemoptysis and higher incidence of dyspnea as compared to adults. Umeki [12] and VandenBrande et al [13] found that hemoptysis occurred equally in both groups.

## 3. Comparison of Bacteriology

High bacillary load was reported in 46% elderly as compared to 23% adults possibly making them more infectious that warrants early diagnosis and management. Tan et al [9] also reported that the elderly had significantly more severe disease and high bacteriological burden compared with age group 15-45 years.

## 4. Comparison of Radiology

Bilateral involvement, lower zone predominance and far advanced disease were more prevalent in elderly compared to adults. Similar observations were reported in tuberculous diabetics by Hariprasad et al [14] also observed that lower lung field is involved more commonly in older age group than adults. Jagdish Rawat et al [10] also reported "a higher involvement of lower zone (24.0% vs. 7.9%) and far advanced lesions (32.0% vs. 14.7%) in the elderly". In our study, minimal lesions were predominantly found among young patients as compared to elderly.

## 7. Conclusion

Atypical presentation with advanced disease and high bacterial load at time of diagnosis is commonly seen in the elderly. Hence high index of suspicion is crucial to early diagnosis and management, thereby reducing morbidity and mortality amongst the patients and curtailing the spread of tuberculosis in the community.

**Conflict of Interest:** None

## References

- [1] Rita Sood. The problem of geriatric tuberculosis. Journal of Indian Academy of Clinical Medicine. Vol.5 (2), 2000; 5:156-8.
- [2] Stead WW, Logfren JP, and Warren E, et al. Tuberculosis as an endemic and nosocomial infection among the elderly in nursing homes. New England Journal of Medicine, 1985; 312:1483-1487.
- [3] Rieder HL, Kelly GD, Bloch AD, et al. Tuberculosis diagnosed at death in the United States. Chest 1991; 100:678-68.

- [4] Tocque K, Bellis MA, Tarn CM, Chan SL et al. Long-term trends in tuberculosis. Comparison of age-cohort data between Hong Kong and England and Wales. American Journal of Respiratory and Critical Care Medicine 1998; 158: 484-8.
- [5] National Tuberculosis Association of the USA. Diagnostic Standards and Classification of Tuberculosis. New York. National Tuberculosis Association, 1961.
- [6] Morris JT, Seaworth BJ, McAllister CK. Pulmonary tuberculosis in diabetics. Chest 1992; 102:539-41
- [7] Tripathy SR, Kar KP, Chakraborty DC, Majumdar AK. Diabetes mellitus and pulmonary tuberculosis – A prospective study. Indian Journal of Tuberculosis 1984; 31:122.
- [8] Perez-Guzman C, Vargas MH, Torres-Cruz A, Villarreal-Velarde H. Does aging modify pulmonary tuberculosis? A meta-analytical review. Chest. 1999 Oct; 116(4):961-7.
- [9] Tan KK, Cherian A, and Teo SK. Tuberculosis in the elderly. Singapore medical journal. 1991; 32(6):423-6. Epub 1991/12/01.
- [10] Jagdish Rawat, Girish Sindhwani and Ruchi Juyal. Clinico-radiological profile of new smear positive pulmonary tuberculosis cases among adults and elderly people. Indian Journal of Tuberculosis 2008; 55: 84-90.
- [11] Patel Anand K, Rami Kiran C, Ghanchi Feroz. Clinical profile of sputum positive pulmonary tuberculosis patients with diabetes mellitus in a teaching hospital at Jamnagar, Gujarat. National journal of medical research Volume 2 Issue 3. July – Sept. 2012
- [12] Umeki S. Comparison of younger and elderly patients with pulmonary tuberculosis. Respiration 1989; 55:75-83.
- [13] Van den Brande P, Vijgen J, Demedts M. Clinical spectrum of pulmonary tuberculosis in older patients: comparison with younger patients. Journal of Gerontology. 1991; 46:M204-9.
- [14] Hariprasad, Ramakrishna MR, Trupti R et al, The study of pulmonary tuberculosis in diabetes mellitus patients. International Journal of Pharma and Bio Sciences 2013 Apr; 4(2): (B) 559-571.

## Author Profile



**Dr. Arun Babu** received his M.B.B.S. degree from Rajiv Gandhi University of Health Sciences, Karnataka, India. From 2012 he is pursuing his Post Graduation in the Department of Respiratory Medicine, Chettinad Hospital and Research Institute, Kancheepuram, Tamilnadu, India