

Computational Assessment of Dose due to Primordial Radionuclides in Soils of Coastal Regions of Kanyakumari District in Tamil Nadu, India

V. Sarojini¹, T. Chithambarathanu²

¹Assistant Professor, Lekshmpuram College of Arts and Science, Neyyoor-629802, Tamil Nadu, India

²Associate Professor, S.T.Hindu College, Nagercoil- 629001, Tamil Nadu, India

Abstract: *The natural radioactivity due to uranium, Thorium and potassium in soil contributes to the radiation dose received by human beings significantly. Gamma ray spectrometric measurements carried out for the natural radioactivity levels due to ²²⁶Ra, ²³²Th and ⁴⁰K in beach sand samples around the coast of Kanyakumari district were analyzed. For assessing the environmental radiological impact to public it is essential to evaluate the activity levels of these radionuclides. The absorbed dose rate due to natural radionuclides was also calculated and the results are reported in this paper. The total annual effective dose of the study area founds to vary between 1.86 mSv to 2.67 mSv. This study provides a baseline data of radioactivity background levels in some Naturally High Background Radiation Areas along the coastal regions of Kanyakumari district and will be useful to assess any changes in the radioactive background levels.*

Keywords: Natural radioactivity, radionuclides, dose

1. Introduction

Terrestrial sources contribute about 84% of the total radiation from natural sources to which an individual is exposed. It is an accumulated effect arising from a number of materials around us such as rock, soil, water, food and from within human body. It is obvious that people living in areas with large concentrations of radionuclides might be getting more than the average annual dose of radiation. Such areas are in Brazil, Sweden, France and coastal areas of Kerala and Tamil Nadu⁴. Similarly, there are active regions rich in granite in France, with a high uranium and thorium content and alumshale, and in Sweden, with high uranium and radium concentrations which also provide larger doses of radiation. Radionuclides are found naturally in air, water and soil. Natural background radiation is of terrestrial and extra-terrestrial origin [1]. Natural environmental radioactivity and the associated external exposure due to gamma radiation depend primarily on the geographical and geological conditions and appear at different levels of the soils of each region in the world. There are few regions in the world known to be high background radiation areas due to local geology and geo chemical effects that cause enhanced levels of terrestrial radiation [2]. In the high background areas of the country such as Austria, Brazil, China, France, India and Iran the radiation levels were found to be high varying over an order of magnitude depending upon the site-specific terrestrial radioactivity[3]. In India there are quite a few monazite sand bearing placer deposits causing high background radiation along its long coastal line. Ullal in Karnataka, Kalppakkam in Tamil Nadu, coastal parts of Tamil Nadu and Kerala state and south western coast of India are known to be high back ground radiation areas [4]. One of the areas in the south west coast where high radiation level has been reported was from coastal regions of Kanyakumari district, in Tamil Nadu. Beach sand

in these areas contains heavy minerals like ilmenite, rutile, zircon, monazite and sillimanite. ²³²Th and ²³⁸U are reported from these regions, caused mainly due to the monazite bearing black sands. Combinations of favorable factors like the hinterland geology, geomorphology, sub-tropical climate and intricate network of drainage aided by wind and coastal processes like waves and currents have influenced these formations [5]. Monazite sands are known to contain thorium with some extent of ²³⁸U and ⁴⁰K. Since the radionuclides are not uniformly distributed the knowledge of their distribution in soil plays an important role in radiation protection and measurement. Also the radioactivity of soils is essential for understanding changes in the natural background [6]. In sense of lives, health and environmental pollution determination of the radioactivity concentration in sands is useful. Therefore an attempt is made in the present investigation to studies on the natural radioactivity content in the beach sands of coastal regions of Kanyakumari district.

2. Literature Survey

Menon et al., explains the naturally occurring radionuclides present in soil include ²²⁸Ra, ²²⁶Ra, ²³⁸U, ²³²Th and ⁴⁰K . Khan et al., gives the knowledge of their distribution in soil and rocks plays an important role in radiation protection and measurement. Uosif.M and Tahir.A provides the assessment of radiation doses from natural radioactive sources is of particular importance as it is the largest contributor to the external dose of the world population. The world is naturally radioactive and around 90% of the human radiation exposure arises from natural sources such as cosmic radiation exposure to radon gas and terrestrial radiation. United Nations Scientific Committee on the effects of Atomic Radiation (UNSCEAR), Sources Effects and Risks of Ionizing Radiation Reported to the General Assembly,

Absorbed dose

The absorbed dose rate (D) [8] due to gamma radiations in air at 1m above ground level for the uniform distribution of naturally occurring nuclides was calculated using equation (1):

$$D = 0.462 C_{Ra} + 0.604 C_{Th} + 0.0042 C_k \dots\dots\dots(1)$$

Where D is the absorbed dose rate in nGyh⁻¹ and C_{Ra}, C_{Th} and C_k are the activity concentrations of ²²⁶Ra, ²³²Th and ⁴⁰K in the soil samples in Bq/Kg.

4.1 Annual Effective Dose

To estimate the annual effective dose, two important factors must be taken into account (i) the conversion coefficient from the absorbed dose in air to the effective dose and (ii) the indoor occupancy factor. The former gives the equivalent dose in Sv⁻¹ from absorbed dose in air Gy⁻¹, while the latter gives the fraction of the time the individual is exposed to outdoor radiation. The first factor has been recommended by UNSCEAR, 2000 as 0.7 SvGy⁻¹ and the second factor as 0.2, which suggests that there is variation from absorbed dose in air to effective dose received by adults. The annual effective dose is calculated, taking into consideration that people in India, on an average, spend nearly 20% of their time outdoors. To estimate the annual effective dose rate, their conversion coefficient from absorbed dose in air to effective dose (0.7 SvGy⁻¹) and outdoor occupancy factor of (0.2) proposed by UNSCEAR, 2000 were used. The annual effective dose rate [9] in units of mSv⁻¹ was calculated by using equation (2):

$$\text{Annual effective dose} = D \times (24 \times 365) \times 0.7 \times 0.2 \times 10^{-6} \text{ mSv} \dots\dots\dots(2)$$

Table 2: Absorbed dose and Annual Effective Dose in the study area

Sampling sites	Absorbed Dose(nGyh ⁻¹)	Annual Effective Dose (mSv)
Enayam	1517.92	1.86
Midalam	1864.43	2.28
Melmidalam	1933.80	2.37
Thengapattinam	1831.39	2.24
Puthenthurai	2178.75	2.67

Table 2. gives the absorbed dose and the annual effective dose from soil. The absorbed dose ranges from 1517.92 to 2178.75 nGyh⁻¹. To estimate the annual effective dose, the indoor occupancy factor and the conversion coefficient from the absorbed dose in air to effective dose must be taken into account. The annual effective dose ranges from 1.86 mSv to 2.67mSv.

5. Conclusion

Radiological assessment for the soil samples collected from the study area indicates that the concentration of radionuclides ranges from 269 to 458 Bq/Kg for ²²⁶Ra, 237 to 427 Bq/Kg for ²²⁸Ra, 2302 to 3250 Bq/Kg for ²³²Th and 77 to 108 Bq/Kg for ⁴⁰K. The absorbed dose in air ranges from 1517.92 nGy/h to 2178.75 nGy/h and the Annual effective dose lie between 1.86 mSv to 2.67 mSv. This enhanced level of radioactivity is due to monazite bearing black sand which

is an orthophosphate of thorium, uranium and rare earths, present with a concentration varying from 0.3 to 5 % in the soils of this area. The highest level of dose 2.67 mSv is estimated at Enayam this is due to the fact that beach sands in these areas contain heavy minerals like ilmenite, rutile, garnet, zircon, monazite, magnetite, sillmenite etc., Combination of favorable factors like the hinter land geology, coastal geomorphology, sub-tropical climate and intricate network of drainage aided by wind and coastal processes like waves and currents have influenced this formation.

6. Future Scope

The dose due to drinking water and air can be further estimated and from this the total dose due to air , water and soil to general public in the area can be determined

References

- [1] UNSCEAR, 2000. *United Nations Scientific Committee on the effects of Atomic Radiation (UNSCEAR), Sources Effects and Risks of Ionizing Radiation Report to the General Assembly, United Nations, New York, USA.*
- [2] Iyengar, M.A.R., and Kannan, V., 1994. Natural Radiation aspects in the high background areas at Kalpakkam, *Proc. 3rd National Symposium on Environment, Thiruvananthapuram, March, 1994*, pp. 48-55. Lakshmi, K.S., Selvasekarpandian, S., Kanna, D. and Meenakshisundaram, V., 2005. Primordial radionuclides concentrations in the beach sands of east coast region of Tamil Nadu, India. *International Congress Series, vol.1276*, 323-324.
- [3] Mohanty, A.K., Sengupta, D., Das, S.K., Saha, S.K., and Vijayan, K.V., 2004. Natural radioactivity and radiation exposure in the high background area at Chhatrapur beach placer deposit of Orissa, India. *Journal of Environmental Radioactivity*, 75, 15-33.
- [4] Ramasamy, V., Senthil, S., Meenakshisundaram and Gajendran, V., 2009. Measurement of natural radioactivity in beach sediments collected from north east coast of Tamil Nadu, India. *Research Journal of Applied Scie., Eng. And Technology*, 1(2) 54-58.
- [5] El-Arabi, A.M., 2007. ²²⁶Ra, ²³²Th and ⁴⁰K concentrations in igneous rocks from eastern desert, Egypt and its radiological implications. *Radiation Measurements*, 42, 94-100. A.A Shanbhag, S.J. Sartandel, T.V. Ramachandran and V.D Puranik 2005. Natural Radioactivity content in beach sands of Ratnagiri coast Maharashtra, *Environmental Geo chemistry*, vol.8, no.1&2, PP.304-308.
- [6] Veiga, R., Sanches, N., Anjos, R.M., Macario, K., Bastos, J., Iguatemy, M., Aguiar, J.G., Santos, A.M.A., Mosquera, B., Carvalho, C., Baptista Filho, M. and Umisedo, N.K., 2006. Measurement of natural radioactivity in Brazilian beach sands. *Radiation Measurements*, 41, 189-196.
- [7] Kerur, B.R., Anil Kumar, S. 2012 Dose due to Radioactive nuclides in Soil of Karnataka. *Proceedings of NSRP-19*, Mamallapuram, Tamil Nadu, India. PP.444-446.

- [8] Menon, M.R., Mishra, U.C., Lalit, B.Y., Shukla, V.K., Ramachandran, T.V., 1982. Uranium, thorium and potassium in Indian rocks and ores. *Proceedings of the Indian Academy of Sciences(Earth Planetary Sciences)* 91, 127-136. Khan, H.M., Khan, K., Atta, M.A., Jan, F., 1994. Measurement of gamma activity of soil samples of Charssada district of Pakistan. *Journal of Chemical Society of Pakistan*, 16, 183-188.
- [9] Uosif, M. and Tahir, A., 2008. Radiological assessment of abu tarpur phosphate western desert Egypt, *Radiation Protection Dosimetry*, 7(1-4), 87-89.
- [10] Overstreet, W. C. (1967) The geologic occurrence of Monazite. U.S. *Geological survey professional paper*, 530.
- [11] Mir Azam Ali, Krishnan, S. and Banerjee, D.C., 2001. Beach and heavy mineral sand investigations and deposits in India- An overview, *special issue on beach and inland heavy mineral sand deposits of India. Exploration and Research for Atomic minerals*, 13, pp.1-21.
- [12] Aswathanarayana, U., 1964. Origin of the heavy mineral beach sand deposits of the south west coast of India. In: *Eds. A.P. Subramanian and S. Balakrishna-Advancing Frontiers in Geology and Geophysics, Indian Geophysical union*, Hyderabad. Pp.481-489.
- [13] Sankaran, A.V., Jayaswal, B., Nambi, K.S.V. and Sunta, C.M., 1988. *U, Th and K distributions inferred from regional geology and the terrestrial radiation profiles in India. BARC, Bombay*, pp.1-104.
- [14] Paul AC, Pillai PMB, Haridasan PP, Sujatha Radhakrishnan and Krishnamony S. Population exposure to airborne thorium at the high natural radiation areas in India, *Journal of Environmental Radioactivity*, 40 (3), 251-259, 1998.
- [15] Lakshmi, K.S., Selvasekarandian, S., Kanna, D. and Meenakshisundaram, V., 2005. Primordial radionuclides concentrations in the beach sands of east coast region of TamilNadu, India. *International Congress Series, vol.1276*, 323-324.
- [16] Khanna, D., Malathi, J., Brahmanandhan, G.M. and Selvasekarandian, S., 2004. Measurement of activity concentrations of ⁴⁰K, ²³⁸U and ²³²Th in soil samples of Agastheeswaram Taluk, Kanyakumari District, India., *paper presented in the Sixth International Conference on High Level Radiation Areas, Japan. September 5-9, 2004.*
- [17] Iyengar, M.A.R., and Kannan, V., 1994. Natural Radiation aspects in the high background areas at Kalpakkam, Proc. 3rd National Symposium on Environment, Thiruvananthapuram, March, 1994, pp. 48-55.

Dr. Cithambarathanu has been working as Associate Professor in S.T.Hindu College, Nagercoil for the past 26 years. He received his Ph.D degree in the year 2003 from M.S.University, Tirunelveli. His major research interests are in Molecular Spectroscopy & Environmental Physics. He has successfully completed his research project entitled "A Study of Surface Ozone variation in Nagercoil, Southern Tamil Nadu" funded by University Grants Commission. He has published more than 50 papers and conducted four national level seminars funded by BARC, NPCIL, IREL and UGC. Sp

Author Profile

Mrs. Sarojini has been working as Assistant Professor in Physics at Lekshmipuram College of Arts & Science, Neyyoor for the past 11 years and doing Research in Radiation Physics. The author is a life member in Indian Science Congress, Indian Society for Radiation Physics, Indian Society for Radiation Biology and in Kanyakumari Academy of Arts & Science. She has also published 8 International papers in the field of Radiation Physics. She received her Ph.D degree in the year 2013.