Traditional Phytotherapy among the Tribes of District Shahdol, Madhya Pradesh, Central India

Manoj Kujur¹, Ramesh Kumar Ahirwar²

Abstract: Present paper deals with 30ethnomedicinal plants have been identified for the treatment of various disease. Harbarium has been prepared which contains information pertaining to botanical name, local name, plants used, their dose and process of administration. A survey of ethnomedicinal plants of shahdol district has been carried out with co-operation of Tribal villagers.

Keywords: Traditional, Ethnomedicinal plants, Tribes, Shahdol, Phytotheraphy

1. Introduction

District Shahdolis situated north eastern part of Madhya Pradesh state. It is lyingbetween 23°17'47" N latitude and 81°21'21" E longitude. Total geographical area sums up to 5671 sq./Km. and has a population of 908148. Shahdol is richedin vast resources of forest and minerals. It is bounded in the north bySatna and Sidhi district, in the east by Korea district, in the south byAnuppur district, in the west by Umaria district. The area is full of watersprings which come out on the top hill slopes. The Shahdol district is average rainfall is 85.11% and above temperature 13.6 °C. The Kol tribes living in some villages situated in and around Shahdol district. They depend solely on their surrounding forest for most of their requirement for food to ethnomedicinal plants.

2. Material and Methods

The area is floristically very rich and harbours a wide range of biodiversity.People living in these forests and adjacent

rural areas depend upon plants for their daily needs and food. Kol tribal villagers have good knowledge about medicinal plants and their uses in different types of diseases. A persusal of literature shows that no systematic study and survey from ethnobotanical and ethnomedicinal point of view has been carried out in the present area of investigation. The present communication deals with 30 ethnomedicinal plant used by the tribal communities in habiting the region of Shahdol district forvarious disease treatment by the Tribals. The plants are arranged alphabetically each by botanical name, family, local name and use of plants. Plants were collected during flowering period with the help of tribals and vaidys. Parts of ethnomedicinal plants used in the treatments of many types disease and other informations were recorded as stated by the tribals and vaidyas. The specimens were identified with the help of published floraand relevant authorities.Some other workers who have contributed in the field of ethnobotany as Jain(1963), Ahirwar (2010), Khan et al (2008) and Verma(1995).




Figure 1: Location Map of Madhya Pradesh in DistrictShahdol.

3. Results and Discussion

We have identified 30 Ethnomedicinal plants, which are uses by tribals of Shahdol district in their daily life for the

treatment of various diseases. Most of these plants are wild and some plants are cultivated. The ethnobotanical plants have been listed alphabetically (Table-1).

Table 1: Shows Traditional Ethnomedicinal plants uses of Tribes in District Shahdol, Madhya Pradesh

Volume 4 Issue 12, December 2015

^{1, 2}Department of Botany, Govt. Nehru P.G. College Burhar, District Shahdol, Madhya Pradesh - 484110 (India)

International Journal of Science and Research (IJSR)

ISSN (Online): 2319-7064

Index Copernicus Value (2013): 6.14 | Impact Factor (2014): 5.611

| S. No. | Name of Family | Botanical Name | Local Name | Ethnomedicinal Used |
|--------|------------------|---------------------------------------|--------------|--|
| 1 | Apiaceae | Centellaasiatica(L). Urban | Bramhi | Entire plant used as nervine tonic, and skin diseases as |
| | | | | weak sedative, cardio depressant, hypotensive and in |
| | | | | leprosy. |
| 2 | Asteraceae | EcliptaprostrataL. | Kala Bhegraj | Entire plant tonic given in spleen enlargement and leaf |
| 2 | | D 10 (7) | G 11 | juice given in eczema and as hair tonic. |
| 3 | Apocynaceae | Rauvolfia serpentine (L.) | Sarapgandha | Leaf juice is used as remedy for opacity of cornea. Root |
| 4 | A1:1 | Benth. | Madar | is sedative, reduces hypertension. Flowers are used in cold, cough and Asthma Powdered |
| 4 | Asclepiadaceae | CalotropisproceraBr. | Madai | Root bark used in dysentery. Fresh leaves in ulcer and as |
| | | | | wormicidal. |
| 5 | Acantahceae | AdhatodaVasicaNees | Adusa | lecoction is used in cough bronchitis and rheumatism. |
| 6 | Acantahceae | Hygrophilaauticulata(Schum) | Talmakhana | Leaves, seeds and roots are used as diuretics and also for |
| | | , | | jaundice, dropsy, rheumatism and urinogenital disorder. |
| 7 | Amaranthaceae | AchyranthesasperaL. | Chirchitta | Used in cough and its decoction is given in renal dropsy |
| | | | | and bronchial infection treatment of |
| | | | | phyrrhoeafever,enlargement of liver and spleen. |
| 8 | Bombacaceae | Bombaxceiba L | Semur | Root is stimulant, tonic and chiefly given in impotency. |
| 9 | Combretaceae | Terminalia arjuna(Roxb.) | Arjun | Bark infusion used in heart troubles and leaf juice in |
| | | Wt. & Am. | | earache, hypertension and as diuretic and has tonic effect |
| 10 | Cucurbitaceae | Cocciniagrandis(L). Voigt | Kundru | in cirrhosis of lever. Juice from leaves and roots is used in diabetes. |
| 11 | Cactaceae | Opundtiaelatior. Mil. | Nagphani | Baked fruit is given in whooping cough and a syrup of |
| 11 | Cactaccac | Opinanaeianor. wiii. | Nagphani | the fruit increases secretion of bile. |
| 12 | Caesalpiniaceae | Cassia fistula L. | Amaltash | Root with black pepper and leaves of kathal applied to |
| | • | J | | cattle for giving relief to swollen throat. |
| 13 | Caesalpiniaceae | Cassia tora L. | Chakwda | Leaf and seeds used in skin disease for ring worm and |
| | | | | itch. |
| 14 | Convolvulaceae | CuscutareflexaRoxb. | Amarbel | Seeds used in ulcer and liver disorders. |
| 15 | Dipterocarpaceae | Shorearobusta Gaertn f. | Sal | Resin is used in skin diseases, diarrhea anddysentery. |
| 16 | Fabaceae | Butea monosperma(Lamk) | Palas | Seeds are internally administerd as an anthelmintic in |
| | | Tamb | | treatment of round worms. Gum is used in treatment of diarrhoea. |
| 17 | Fabaceae | DalbergiasissooRoxb | Shisham | Decoction of leaf Useful in gonorrhoea. |
| 18 | Fabaceae | Mucunapruriens(L) DC | Kemmach | Root used in paralysis seeds are used as nervine tonic, |
| | | (=) = 0 | | menses troubles and as vermifuge, strong infusion of |
| | | | | roots mixed with honey is given in cholera. |
| 19 | Lythraceae | LawsoniainerdmisL. | Mehndi | Lea paste or decoction used as gargle in sore throat, |
| | | | | burns, bruises and skin inflammation, Seeds useful in |
| 20 | | , , , , , , , , , , , , , , , , , , , | | urinary troubles. |
| 20 | Lamiaceae | Leucascephalotes(Wild.) | Gumi | Leaf used as digestive antiseptic and earache and fever. |
| 21 | Lamiaceae | (Roth) Spr. Ocimumamericanum L. | Kali tulsi | Flowers are used in cough and cold. Leaf decoction is given in Maleria and abdominal |
| 21 | Lamaccac | Ocimamamericanum L. | Kan tuisi | diseases. |
| 22 | Menispermaceae | Tinosporacordifolia(L.)Merr | Gurch | Root extract is fever, cold cough, as blood purifier, in |
| | 1 1 | | | acidity and jaundice. |
| 23 | Meliaceae | Azadirachtaindica A. Juss | Neem | Bark, leaf and fruit decoction is antiseptic and used in |
| | | | | ulcer, eczema and skin diseases. |
| 24 | Mimosaceae | Acacia nilotica (L.) Del. | Babul | Bark used in diarrhea, dysentery and diabetes. |
| 25 | Mimosaceae | Albizdialebbeck (L.)Willd | Siris | Bark and seeds given in piles and diarrhea. |
| 26 | Nyctaginaceae | Boerhaaviadiffusa L. | Punarnava | Root is laxative and used in asthma, jaundice and dropsy. |
| 27 | Papaveraceae | ArgemonemexicanaL | Pilikateli | Stem latex used in dropsy, jaundice and conjunctivitis. |
| 28 | Papaveraceae | Papaver somniferumL | Afeem | Flower, fruits and Seeds have pain releasing and sleeping effects and useful in irritating cough, pneumonia, ulcers, |
| | | | | gastritis and influenza |
| 29 | Rutaceae | Aegle marmelos (L)Corr | Bel | Fruits is given in diarrhea dysentery irriation of |
| 1 - | | 2,001 | _ •• | alimentary canal, fever and as tonic and cooling agent. |
| 30 | Sapotaceae | Madhucalongifolia(Koen) | Mahua | Decoction of bark is used in incurring bleeding gums and |
| | - | Mac Br. | | ulcers. Flowers are used in cough and bronchitis. |

4. Acknowledgements

The authors are thankful to their respective college authorities for granting permission to carry out of this work. They are also thankful to Tribals and rural people who co-operated in sharing their knowledge on Ethnomedicinal uses.

References

- [1] Agrawal, D.P. (1997) Traditional Knowledge Systems and Western Science *current science***73**, 731-733.
- [2] Ahirwar, R.K. (2010) A Survey of Medicinal Plants used by tribals of Anuppur District, central India.*Ind. J. Applied &Pure Bio.* Vol.**25** (II) 227-230.

Volume 4 Issue 12, December 2015

International Journal of Science and Research (IJSR)

ISSN (Online): 2319-7064

Index Copernicus Value (2013): 6.14 | Impact Factor (2014): 5.611

- [3] Ahirwar, R.K. (2011) Ethnomedicinal Plants Studies in Jaitpur Forest range of Shahdol, District, Central India. *Ad. PlantSci. Vol.* **24** (II) 681-684.
- [4] Ahirwar,R.K. (2014) Utilization of Medicinal Plants by the Tribes of Bhatiya, District Shahdol, Madhya Pradesh. *Int.J.Sci.andRes*.**3**(9).149-151.
- [5] Ahirwar, Ramesh Kumar (2010) Ethnomedicinal uses of plant roots from Shadol district of M.P. India. *Ind. J.Appl. Pure Bio.* **25** (1) 71-76.
- [6] Ahirwar, Ramesh Kumar and Singh,G.K. (2011) Some anti diabetic plants from Dindori District of Madhya Pradesh (India). *Ind. J.Appl. pure Bio.* **26** (2) 269-271.
- [7] Ahirwar, R.K. (2015) Diversity of Ethnomedicinal Plants in Boridand Forest of District Korea, Chhattisgarh,India. *American Journal of Plant Sciences*, **6**, 413-425. http://dx.doi.org/10.4236/ajps.2015.62047
- [8] Biswas, T.K. and Mukherjee, B. (2003) Plant Medicines of Indian Origin for Wound Healing Activity: A Review. *In-ternational Journal of Lower Extremity Wounds*, 2, 25-39.
- [9] Busia, K. (2005) Medical Provision in Africa-Past and Present. *Phytotherapy Research*, 19, 919-923. http://dx.doi.org/10.1002/ptr.1775
- [10] Chopra R.N, Nyar SL & Chopra I.C.(1956) Glossary of Indian Medicinal Plants ,Council of scientific and Industrial Research, New Delhi.
- [11] Hemadri, Koppala&Rao.S.S.(1989) Folk Medicine of Bastar, *J. Ethnobotany* **1**, 61-66.
- [12] Jain S.K. (1963) Observation on the Tribals of Madhya Pradesh *Vanyajati*. **11**, 177-183.
- [13] Jain, S.K. (1991) Dictionary of folk Medicines and Ethnobotany, Deep Publications, New delhi.
- [14] Jain, S.K. (1981) Glimpses of Ethnobotany, Oxford & IBH. NewDelhi.
- [15] Jain, S.K.(1965) Medicinal Plant lore of tribals of Bastar Econ. Bot, 19, 236-250.
- [16] Jain, S.P. Singh, S.C. (1994) Ethno-medico-botanical survey of Ambikapur, Madhya Pradesh, IV ICE, Lucknow, Nov.17-21.
- [17] Khan.A.A, Agnihotri, S.K, Singh M.K & Ahirwar, R.K. (2008) Enumaration of certain Angiospermic plants used by Baiga tribe for Conservations of Plants Species. *Plant Archives***8**, (I) 289-291.
- [18] Kirtikar, K.R. &Basu,B.D.(1999) Indian Medicinal Plants (International book Distributors Book sellers and Publishers, Deheradun.) Vol. 3.
- [19] Kumar, K. (2007) Working Plan of North Surguja, Ambikapur Forest Department Govt.of Chhattisgarh, Raipur.
- [20] Khan.A.A, Agnihotri, S.K, Singh M.K & Ahirwar, R.K. (2008) Observation of Certain Plants used in skin diseases by Baiga Tribes of Mandala district. *Plant Archives* 8. (I) 283-284.
- [21] Kala, C.P. (2007) Local Preferences of Ethno-Botanical Species in the Indian Himalaya: Implications for Environmental Conservation. *Current Science*, **93**, 1828-1834.
- [22] Kala, C.P. (2005) Indigenous Uses, Population Density and Conservation of Threatened Medicinal Plants in Protected Areas of the Indian Himalayas. *Conservation Biology*, 19, 368-378. http://dx.doi.org/10.1111/j.1523-1739.2005.00602.x

- [23] Maheswari, J.K. (1964) A Contribution to the Flora Kanha National Park, Madhya Pradesh *Bull. Bot. surv.India.* **5**(2), 177-140.
- [24] Masih, V., Sahu, P.K. and Singh, M. (2013) Observation on Ethno-Medicinal Herbs of Dantewada, Chhattisgarh India. *Interna- tional Journal of Drug Discovery and Herbal Research*, 3, 644-648.
- [25] Miller, N.N., (1990) Traditional medicine in East Africa, America Universities field staff report, 22, pp 1-15.
- [26] Oommachan M, Masih S.K. (1989) Ethnobotanical observations on certain forest plants of Jabalpur (M.P.) *Ind. J. Appl. pure Bio.***4** (2) 73-78