

Solving Hybrid Fuzzy Fractional Differential Equations by Runge Kutta 4th Order Method

S. Ruban Raj¹, M. Saradha²

¹Associate Professor, Department of Mathematics, St. Joseph College, Trichirappalli, Tamilnadu, India

²Research Scholar, Department of Mathematics, St. Joseph College, Trichirappalli, Tamilnadu, India

Abstract: In this paper we study numerical methods for hybrid fuzzy fractional differential equation, Degree of Sub element hood and the iteration method is used to solve the hybrid fuzzy fractional differential equations with a fuzzy initial condition.

Keywords: hybrid fuzzy fractional differential equations, Degree of Sub Element Hood

Mathematical Subject Classification (MSC) Code: 65L06

1. Introduction

Fuzzy initial value problems for fractional differential equations have been considered by some authors recently [2, 3]. To study some dynamical processes, it is necessary to take into account imprecision, randomness or uncertainty. The objective of the present paper is to extend the application of the iteration method, to provide approximate solutions for fuzzy initial value problems of differential equations of fractional order, and to make comparison with that obtained by an exact fuzzy solution.

2. Analytical solution of Hybrid Fuzzy Fractional Differential Equations

Let us consider the following fractional differential equation

$${}_c D_a^\beta x(t) = f(t, x(t), \lambda_k(x_k)), t \in [t_k, t_{k+1}] \quad (1)$$

$$x(t_k) = x_k$$

Where, $0 \leq t_0 \leq t_1 \leq \dots \leq t_k \rightarrow \infty$

3. The Fourth Order Runge Kutta Method With Harmonic Mean

For a hybrid fuzzy fractional differential equation we develop the fourth order Runge Kutta method with harmonic mean when f and λ_k in (1) can be obtained via the Zadeh extension principle from:

$$f \in [R^+ \times R \times R, R] \text{ and } \lambda_k \in C [R, R]$$

we assume that the existence and uniqueness of solutions of (1) hold for each $[t_k, t_{k+1}]$. For a fixed r , to integrate the system in (3) $[t_0, t_1], [t_1, t_2], \dots, [t_k, t_{k+1}], \dots$ we replace each interval by a set of N_{k+1} discrete equally spaced grid points (including the end points) at which the exact solution $x(t; r) = (\underline{x}(t; r), \overline{x}(t; r))$ is approximated by some $(\underline{y}(t; r), \overline{y}(t; r))$. For the chosen grid points on $[t_k, t_{k+1}]$ at

$$t_{k,n} = t_k + nh_k, \quad h_k = \frac{t_{k+1} - t_k}{N_k}, \quad 0 \leq n \leq N_k.$$

Let $(\underline{Y}_k(t; r), \overline{Y}_k(t; r)) \equiv (\underline{x}_k(t; r), \overline{x}_k(t; r))$, $(\underline{y}_k(t; r), \overline{y}_k(t; r))$ and $(\underline{y}_{k,n}(t; r), \overline{y}_{k,n}(t; r))$ may be denoted respectively by $(\underline{Y}_{k,n}(t; r), \overline{Y}_{k,n}(t; r))$ and $(\underline{y}_{k,n}(t; r), \overline{y}_{k,n}(t; r))$.

We allow N_k 's to vary over the $[t_k, t_{k+1}]$'s so that the h_k 's may be comparable. The Fourth Order Runge Kutta method for (1) is given by:

$$(\underline{Y}_k(t; r), \overline{Y}_k(t; r)) \equiv (\underline{x}_k(t; r), \overline{x}_k(t; r)),$$

$$(\underline{y}_k(t; r), \overline{y}_k(t; r))$$

Where

$$k_1(t_{k,n}; y_{k,n}(r)) = \min \left\{ \begin{array}{l} h_k f(t_{k,n}, u, \lambda_k(u_k)) \\ u \in [\underline{y}_{k,n}(r), \overline{y}_{k,n}(r)] \\ u_k \in [\underline{y}_{k,n}(r), \overline{y}_{k,n}(r)] \end{array} \right\},$$

$$\overline{k}_1(t_{k,n}; y_{k,n}(r)) = \max \left\{ \begin{array}{l} h_k f(t_{k,n}, u, \lambda_k(u_k)) \\ u \in [\underline{y}_{k,n}(r), \overline{y}_{k,n}(r)] \\ u_k \in [\underline{y}_{k,n}(r), \overline{y}_{k,n}(r)] \end{array} \right\},$$

$$k_2(t_{k,n}; y_{k,n}(r)) = \min \left\{ \begin{array}{l} h_k f(t_{k,n} + \frac{1}{2}(h_k), u, \lambda_k(u_k)) \\ u \in \left[\frac{\Phi_{k_1}(t_{k,n}, y_{k,n})}{\overline{\Phi}_{k_1}(t_{k,n}, y_{k,n})} \right] \\ u_k \in [\underline{y}_{k,0}(r), \overline{y}_{k,0}(r)] \end{array} \right\},$$

$$\bar{k}_2(t_{k,n}; y_{k,n}(r)) = \max \left\{ \begin{array}{l} h_k f(t_{k,n} + \frac{1}{2}(h_k), u, \lambda_k(u_k)) \\ u \in \left[\frac{\Phi_{k_1}(t_{k,n}, y_{k,n})}{\bar{\Phi}_{k_1}(t_{k,n}, y_{k,n})} \right] \\ u_k \in [\underline{y}_{k,0}(r), \bar{y}_{k,0}(r)] \end{array} \right\}, \quad \underline{k}_4(t_{k,n}; y_{k,n}(r)) = \min \left\{ \begin{array}{l} h_k f(t_{k,n} + h_k, u, \lambda_k(u_k)) \\ u \in \left[\frac{\Phi_{k_3}(t_{k,n}, y_{k,n})}{\bar{\Phi}_{k_3}(t_{k,n}, y_{k,n})} \right] \\ u_k \in [\underline{y}_{k,0}(r), \bar{y}_{k,0}(r)] \end{array} \right\}$$

$$\underline{k}_3(t_{k,n}; y_{k,n}(r)) = \min \left\{ \begin{array}{l} h_k f(t_{k,n} + \frac{1}{2}(h_k), u, \lambda_k(u_k)) \\ u \in \left[\frac{\Phi_{k_2}(t_{k,n}, y_{k,n})}{\bar{\Phi}_{k_2}(t_{k,n}, y_{k,n})} \right] \\ u_k \in [\underline{y}_{k,0}(r), \bar{y}_{k,0}(r)] \end{array} \right\}, \quad \bar{k}_4(t_{k,n}; y_{k,n}(r)) = \max \left\{ \begin{array}{l} h_k f(t_{k,n} + h_k, u, \lambda_k(u_k)) \\ u \in \left[\frac{\Phi_{k_3}(t_{k,n}, y_{k,n})}{\bar{\Phi}_{k_3}(t_{k,n}, y_{k,n})} \right] \\ u_k \in [\underline{y}_{k,0}(r), \bar{y}_{k,0}(r)] \end{array} \right\}$$

Where

$$\begin{aligned} \underline{\Phi}_{k_1}(t_{k,n}, y_{k,n}(r)) &= \underline{y}_{k,n}(r) + \frac{1}{2} \underline{k}_1(t_{k,n}, y_{k,n}(r)) \\ \bar{\Phi}_{k_1}(t_{k,n}, y_{k,n}(r)) &= \bar{y}_{k,n}(r) + \frac{1}{2} \bar{k}_1(t_{k,n}, y_{k,n}(r)) \\ \underline{\Phi}_{k_2}(t_{k,n}, y_{k,n}(r)) &= \underline{y}_{k,n}(r) + \frac{1}{2} \underline{k}_2(t_{k,n}, y_{k,n}(r)) \\ \bar{\Phi}_{k_2}(t_{k,n}, y_{k,n}(r)) &= \bar{y}_{k,n}(r) + \frac{1}{2} \bar{k}_2(t_{k,n}, y_{k,n}(r)) \\ \underline{\Phi}_{k_3}(t_{k,n}, y_{k,n}(r)) &= \underline{y}_{k,n}(r) + \underline{k}_3(t_{k,n}, y_{k,n}(r)) \\ \bar{\Phi}_{k_3}(t_{k,n}, y_{k,n}(r)) &= \bar{y}_{k,n}(r) + \bar{k}_3(t_{k,n}, y_{k,n}(r)) \end{aligned}$$

$$\begin{aligned} S_k[t_{k,n}, \underline{y}_{k,n}(r), \bar{y}_{k,n}(r)] &= \frac{1}{6} \{ \underline{k}_1(t_{k,n}; y_{k,n}(r)) + 2[\underline{k}_2(t_{k,n}; y_{k,n}(r)) + \underline{k}_3(t_{k,n}; y_{k,n}(r))] + \underline{k}_4(t_{k,n}; y_{k,n}(r)) \} \\ T_k[t_{k,n}, \underline{y}_{k,n}(r), \bar{y}_{k,n}(r)] &= \frac{1}{6} \{ \bar{k}_1(t_{k,n}; y_{k,n}(r)) + 2[\bar{k}_2(t_{k,n}; y_{k,n}(r)) + \bar{k}_3(t_{k,n}; y_{k,n}(r))] + \bar{k}_4(t_{k,n}; y_{k,n}(r)) \} \end{aligned}$$

The exact solution at $t_{k,n+1}$ is given by:

$$\begin{cases} F_{k,n+1}(r) = \underline{Y}_{k,n}(r) + S_k[t_{k,n}, \underline{y}_{k,n}(r), \bar{y}_{k,n}(r)], \\ G_{k,n+1}(r) = \bar{Y}_{k,n}(r) + T_k[t_{k,n}, \underline{y}_{k,n}(r), \bar{y}_{k,n}(r)]. \end{cases} \dots\dots(4)$$

Where $|F_{k,n+1}| = \sum_{\epsilon_j \in A} \exp(F_{k,n+1})$
 and $\mathfrak{S}(G_{k,n+1}, F_{k,n+1}) = \frac{1}{|G_{k,n+1}|} \{ |G_{k,n+1}| - \sum \max\{0, (G_{k,n+1}) - (F_{k,n+1})\} \}$.

4. Degree of Sub Element hood

Let X be a Universal, U be a set of parameters and let $(F_{k,n+1})$ and $(G_{k,n+1})$ are two fuzzy elements of X. Then the degree of sub element hood denoted by $\mathfrak{S}(F_{k,n+1}, G_{k,n+1})$ is defined as,

$$\mathfrak{S}(F_{k,n+1}, G_{k,n+1}) = \frac{1}{|F_{k,n+1}|} \{ |(F_{k,n+1})| - \sum \max\{0, (F_{k,n+1}) - (G_{k,n+1})\} \}$$

5. Numerical Examples

In this section, we present the examples for solving hybrid fuzzy fractional differential equations.

Example: 1
 Consider the following linear hybrid fuzzy fractional differential equation:
 ${}_c D_a^\beta X(t) = 1 + X^2 \dots\dots (5)$
 $X(0) = X_0,$

where $\beta \in (0,1]$, $t > 0$, and X_0 is any triangular fuzzy number. This problem is a generalization of the following hybrid fuzzy fractional differential equation:

$${}_c D_a^\beta x(t) = 1+x^2(t) = 1 + [\underline{x}(t;r), \bar{x}(t;r)]^2 \quad \dots(6)$$

$$x(t) = x_0,$$

Where $\beta \in (0,1]$, $t > 0$, α is the step size and x_0 is a real number.

We can find the solution of the hybrid fractional fuzzy differential equation, by the method of Runge kutta 4th order Method. We compared & generalized the hybrid fractional fuzzy differential equation solution with the exact solution in the following table; also we illustrated the figure for this generalization by using Matlab.

Table 1: Numerical Solution of Example 1

α	$Y_{k,n+1}$	$F_{k,n+1}$	$G_{k,n+1}$
0.1	1.187298247867520	0.787298247867520	.587298247867520
0.2	1.362772659265217	0.962772659265217	1.762772659265216
0.3	1.533606860771214	1.133606860771214	1.933606860771214
0.4	1.702940459085539	1.302940459085540	2.102940459085539
0.5	1.872820986069762	1.472820986069763	2.272820986069762
0.6	2.044858431782411	1.644858431782411	2.444858431782411
0.7	2.220494863603316	1.820494863603317	2.620494863603316
0.8	2.401149486106885	2.001149486106885	2.801149486106885
0.9	2.588315276211985	2.188315276211985	2.988315276211985
1.0	2.783636789711434	2.383636789711434	3.183636789711434

$$|F_{k,n+1}| = 15.69789406$$

$$|G_{k,n+1}| = 23.69789406$$

$$\mathfrak{S}(F_{k,n+1}, G_{k,n+1}) \cong 1$$

$$\mathfrak{S}(G_{k,n+1}, F_{k,n+1}) = 0.66241726 \cong 0.66$$




Figure 1: Comparison of exact and approximated solution of Example 1

6. Conclusion

In this paper, we have studied a hybrid fuzzy fractional differential equation. Final results showed that the solution of hybrid fuzzy fractional differential equations approaches the solution of hybrid fuzzy differential equations as the fractional order approaches the integer order. The results of the study reveal that the proposed method with fuzzy fractional derivatives is efficient, accurate, and convenient for solving the hybrid fuzzy fractional differential equations.

References

- [1] Jafari, H, Saeidy, M, Baleanu, D: The variational iteration method for solving n-th order fuzzy differential equations. Cent. Eur. J. Phys. 10(1), 76-85 (2012)
- [2] Arshad, S, Lupulescu, V: Fractional differential equation with the fuzzy initial condition. Electron. J. Differ. Equ. 34, 1-8 (2011)
- [3] Agarwal, RP, Lakshmikanthama, V, Nieto, JJ: On the concept of solution for fractional differential equations with uncertainty. Nonlinear Anal. 72, 2859-2862 (2010)
- [4] G. Badowski and G. Yin, Stability of hybrid dynamic systems containing singularly perturbed random processes, IEEE Trans. Automat. Control 47 (2002), 2021–2032.
- [5] L. Debnath, Recent applications of fractional calculus to science and engineering, Int. J. Math. Sci. 2003 (2003), 3413–3442.
- [6] N. J. Ford and J. A. Connolly, Comparison of numerical methods for fractional differential equations, Commun. Pure Appl. Anal. 5 (2006), 289–306.
- [7] V. Lakshmikantham and X. Z. Liu, Impulsive hybrid systems and stability theory, Intern. J. Nonl. Diff. Eqns. 5 (1999), 9–17.
- [8] K. Miller and B. Ross, An introduction to the fractional calculus and fractional differential equations, Wiley, New York, 1993.
- [9] K. Oldham and J. Spanier, The fractional calculus, Academic Press, New York-London, 1974.
- [10] S. Ruban Raj, M. Saradha, Solving Hybrid Fuzzy Fractional Differential Equations by Runge Kutta 4th Order Method, International Journal Science and Research(IJSR), ISSN:2319-7064,volume 4, Issue 1, January 2015.
- [11] S. Pederson and M. Sambandham, Numerical solution to hybrid fuzzy systems, Mathematical and Computer Modelling 45 (2007), 1133–1144.
- [12] I. Podlubny, Fractional differential equations- An introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications, Academic Press, San Diego, CA, 1999.
- [13] G. Yin and Q. Zhang, Stability of nonlinear hybrid systems, in: New trends in nonlinear dynamics and control, and their applications, Lecture Notes in Control and Inform. Sci., 295.
- [14] S. Ruban Raj, M. Saradha, Properties of Fuzzy Soft Set, International Journal for Basic Sciences and Social Sciences (IJBS), ISSN:2319-2968, 2(1) (2013), pp.112-118.
- [15] S. Pederson and M. Sambandham, Numerical Solution Of Hybrid Fractional Differential Equations, Communications In Applied Analysis 12 (2008), No. 4, 429–440.