

Integral Solutions of the Homogeneous Biquadratic Diophantine Equations with Five Unknowns

$$(X^2 - Y^2)(3X^2 + 3Y^2 - 2XY) = 12(Z^2 - W^2)T^2$$

Dr. P. Jayakumar¹, G. Shankarakalidoss²

¹Department of Mathematics, A.V.V.M Sri Pushpam College (Autonomous), Poondi, Thanjavur-613503, India

²Department of Mathematics, Kings College of Engineering, Punalkulam, Pudukkottai (Dist) -613303, India

Abstract: Four different patterns are used to find non-zero distinct integral solutions for the homogeneous biquadratic Diophantine equations $(X^2 - Y^2)(3X^2 + 3Y^2 - 2XY) = 12(Z^2 - W^2)T^2$. Different types of properties are exposed in every pattern with polygonal, nasty, square and cubic numbers.

Keywords: Homogeneous biquadratic, integral solutions, special numbers

2010 Mathematics subject classification: 11D09.

Notations used

$T_{m,n}$ - Polygonal number of rank n with size m
 P_n^m - Pyramidal number of rank n with size m
 g_n - Gnomonic number of rank n
 Pr_n - Pronic number of rank n
 CP_n^6 - Centered hexagonal pyramidal number of rank n
 OH_n - Octahedral number of rank n
 SO_n - Stella octangular number of rank n

1. Introduction

The number theory is queen of Mathematics. In particular, the Diophantine equations have a blend of attracted interesting problems. For an extensive review of variety of problems, one may refer to [3-12]. In this work, we are observed another interesting four different methods of the non-zero integral solutions of the homogeneous Biquadratic Diophantine equations with five unknowns $(X^2 - Y^2)(3X^2 + 3Y^2 - 2XY) = 12(Z^2 - W^2)T^2$. Further, some elegant properties among the special numbers and the solutions are observed.

2. Method of Analysis

The homogeneous Biquadratic

Diophantine equations to be solved is $(X^2 - Y^2)(3X^2 + 3Y^2 - 2XY) = 12(Z^2 - W^2)T^2$ (1)

Introducing the linear transformations

$X = u + v, Y = u - v, Z = 2u + v, W = 2u - v$, where $u \neq v \neq 0$ (2)

in (1), we get $u^2 + 2v^2 = 6T^2$ (3)

Assume $T(a, b) = a^2 + 2b^2$, where $(a, b \neq 0)$ (4)

We present below four different patterns of non-zero distinct integer solutions to (1).

2.1 Pattern 1:

Take 6 as

$$6 = (2 + i\sqrt{2})(2 - i\sqrt{2}) \quad (5)$$

Substituting (4) and (5) in (3) and applying the method of factorization, we get

$$(u + i\sqrt{2}v) = (2 + i\sqrt{2})(a + i\sqrt{2}b)^2$$

Equating real and imaginary parts, we get

$$u = 2a^2 - 4b^2 - 4ab \quad (6)$$

$$v = a^2 - 2b^2 + 4ab \quad (7)$$

Putting (6) and (7) in (2), we get non-zero distinct integer valued for x, y, z, w and satisfying (1) are given below

$$X = X(a, b) = 3a^2 - 6b^2 \quad (8)$$

$$Y = Y(a, b) = a^2 - 2b^2 - 8ab \quad (9)$$

$$Z = Z(a, b) = 5a^2 - 10b^2 - 4ab \quad (10)$$

$$W = W(a, b) = 3a^2 - 6b^2 - 12ab \quad (11)$$

The equations (8) to (11) and (4) give non-zero distinct integral solutions of (1) in two parameters.

Properties:

- $X(A, A(A+1)) + 24T_{3,A}^2 = 3T_{4,A}$
- $Y(A(A+1), A+2) - 4T_{3,A}^2 + T_{6,A} + 48P_A^3 \equiv -8 \pmod{9}$
- $X(A, A+1) - 3Y(A, A+1) = 8Pr_A$
- $5Y(A, 1) - Z(A, 1) + g_{15A+1} = 0$
- $8T(A, A)$ is a Nasty number
- $-6Y(A, A)$ is a Nasty number
- $-2\{w(A, A)\}$ is a Nasty number

2.2 Pattern:2

Instead of (5), Take 6 as

$$6 = (-2 + i\sqrt{2})(-2 - i\sqrt{2}) \quad (12)$$

We do the same procedure as in pattern:1 and from (2) we get

$$X = X(a, b) = -a^2 + 2b^2 - 8ab \quad (13)$$

$$Y = Y(a, b) = -3a^2 + 6b^2 \quad (14)$$

$$Z = Z(a, b) = -3a^2 + 6b^2 - 12ab \quad (15)$$

$$W=W(a, b) = -5a^2 + 10b^2 - 4ab \quad (16)$$

The equations (13) to (16) and (4) give non-zero distinct integral solutions of (1) in two parameters.

Properties:

1. $X(A, A(A+1)) + T_{4,A} + 16P_A^5 = 8T_{3,A}^2$
 2. $X(A, 1) - 3Y(A, 1) + T(A, 1) - T_{4,A} \equiv 2 \pmod{24}$
 3. $5Y(A, 1) - Z(A, 1) + T(A, 1) - T_{4,A} \equiv 2 \pmod{36}$
 4. $X(A, 1) - W(A, 1) + T(A, 1) - 12Pr_A \equiv 2 \pmod{11}$
 5. $\{W(A, A)\}$ is a Square number
- Each of the following represents a Nasty number
6. $8\{Y(A, A)\}$
 7. $\frac{8}{3}\{Z(A, A)\}$
 8. $-12\{X(A, A)\}$

2.3 Pattern:3

We write 6 as

$$6 = \frac{(22 + i\sqrt{2})(22 - i\sqrt{2})}{81} \quad (17)$$

Using (17) and (4) in (3), and applying method of factorization, we get

$$(u + i\sqrt{2}v) = \frac{(22 + i\sqrt{2})}{9} (a + i\sqrt{2}b)^2 \quad (18)$$

Equating real and imaginary part, we get

$$u = \frac{1}{9} [22a^2 - 44b^2 - 4ab]$$

$$v = \frac{1}{9} [a^2 - 2b^2 + 44ab]$$

In view of (2), we get the values of X, Y, Z, and W as

$$X = X(a, b) = \frac{1}{9} [23a^2 - 46b^2 + 40ab] \quad (19)$$

$$Y = Y(a, b) = \frac{1}{9} [21a^2 - 42b^2 - 48ab] \quad (20)$$

$$Z = Z(a, b) = \frac{1}{9} [45a^2 - 90b^2 + 36ab] \quad (21)$$

$$W = W(a, b) = \frac{1}{9} [43a^2 - 86b^2 - 52ab] \quad (22)$$

Choose a and b so that the values of X, Y, Z, and W are in integer,

Putting $a = 3A, b = 3B$ in (19) to (22) and (4), we get

$$X = X(a, b) = 23a^2 - 46b^2 + 40ab \quad (23)$$

$$Y = Y(a, b) = 21a^2 - 42b^2 - 48ab \quad (24)$$

$$Z = Z(a, b) = 45a^2 - 90b^2 + 36ab \quad (25)$$

$$W = W(a, b) = 43a^2 - 86b^2 - 52ab \quad (26)$$

$$T = T(a, b) = 9A^2 + 18B^2 \quad (27)$$

The equations (23) to (27) give non-zero distinct integral solutions of (1) in two parameters.

Properties:

1. $9X(A, A+1) - 23T(A, A+1) + T_{1658,A} - 36Pr_A \equiv -828 \pmod{2483}$
2. $Z(A, A(A+1)) - 5T(A, A(A+1)) = 72P_A^5$
3. $Y(1, B) + T_{86,B} \equiv 21 \pmod{89}$
4. $X(A, A^2) + 40CP_n^6 + 69T_{4,A}^2 = 23Pr_{A^2}$

$$5. W(A, 2A^2+1) + 344Pr_{A^2} + 43T_{4,A} + 156OH_A \equiv 0 \pmod{2}$$

$$6. Z(A, 2A^2-1) + 360Pr_{A^2} - 45T_{4,A} + 36SO_A \equiv 0 \pmod{5}$$

$$7. -\frac{1}{13} \{X(A, A) + W(A, A)\} \text{ is a Nasty number}$$

$$8. -\frac{3}{13} \{Z(A, A) + W(A, A)\} \text{ is a Nasty number}$$

2.4 Pattern:4

Rewrite (1) as

$$6T^2 - u^2 = 2v^2 \quad (28)$$

Consider 2 as

$$2 = (\sqrt{6} + 2)(\sqrt{6} - 2) \quad (29)$$

$$\text{Assume } v(a, b) = 6a^2 - b^2; a, b \neq 0 \quad (30)$$

Substituting (29), (30) in (28) and applying method of factorization, we get

$$(\sqrt{6}T + u) = (\sqrt{6} + 2)(\sqrt{6}a + b)^2 \quad (31)$$

Equating real and imaginary part, we get

$$T = T(a, b) = 6a^2 + b^2 + 4ab \quad (32)$$

$$u = u(a, b) = 12a^2 + 2b^2 + 12ab \quad (33)$$

Applying (30) and (33) in (2), we get

$$X = X(a, b) = 18a^2 - b^2 + 12ab \quad (34)$$

$$Y = Y(a, b) = 6a^2 + 3b^2 + 12ab \quad (34)$$

$$Z = Z(a, b) = 30a^2 + 3b^2 + 24ab \quad (35)$$

$$W = W(a, b) = 18a^2 + 5b^2 + 24ab \quad (36)$$

$$T = T(a, b) = 6a^2 + b^2 + 4ab \quad (37)$$

The equation (34) to (37) give the non-zero distinct integral solutions of (1) in two parameters.

Properties:

1. $X(A, A+1) - T_{36,A} - g_{7A} = 12Pr_A$
2. $5Y(1, B) - Z(1, B) - 12Pr_B - g_{12B} = 1$
3. $X(1, B) - W(1, B) + T_{14,B} \equiv 0 \pmod{19}$
4. $Y(B, B) - T(B, B) - T_{6,B} \equiv 0 \pmod{3}$
5. $\frac{1}{2} \{X(A, A) + Y(A, A)\}$ is a Square number
6. $\{Y(A, A^2) - T(A, A^2) - 2T_{4,A}^2\}$ is a Cubic number
7. $\frac{3}{2} \{Y(A, A) + T(A, A)\}$ is a Nasty number

3. Conclusion

In this work, we have observed four different patterns of the non-zero integer solutions of the homogeneous Biquadratic Diophantine equation $(X^2 - Y^2)(3X^2 + 3Y^2 - 2XY) = 12(Z^2 - W^2)T^2$ and relations between solutions and special numbers are also obtained. One may research for any other patterns of this equation and their corresponding properties.

References

- [1] Dickson, L.E., History of Theory of Numbers, Vol.2, Chelsea Publishing company, New York, 1952
- [2] Mordell, L.J., Diophantine equations, Academic press, New York, 1969
- [3] Jayakumar. P, Sangeetha. K "Lattice points on the cone $x^2 + 9y^2 = 50z^2$ " International Journal of Science and Research, Vol (3), Issue 12, 20-2 (December - 2014)

- [4] Jayakumar. P, Kanaga Dhurga. C, “ On Quadratic Diophantine equation $x^2 + 16y^2 = 20z^2$ ” Galois J. Maths ,1(1)(2014), 17-23
- [5] Jayakumar. P, Kanaga Dhurga. C, “Lattice points on the cone $x^2 + 9y^2 = 50z^2$ ” Diophantus J. Math., 3(2)(2014), 61-71
- [6] Jayakumar,P, Prabha. S “On Ternary Quadratic Diophantine equation $x^2 + 15y^2 = 14z^2$ ”Archimedes J. Math., 4(3) (2014), 159-164
- [7] Jayakumar. P, Meena. J ‘Integral solutions of the Ternary Quadratic Diophantine equation: $x^2 + 7y^2 = 16z^2$ ” International Journal of Science and Technology, Vol.4, Issue 4, 1-4, Decembe2014.
- [8] Jayakumar. P, Shankarakalidoss. G ““Lattice points on Homogenous cone $x^2 + 9y^2 = 50z^2$ ” International Journal of Science and Research, Vol (4), Issue 1 , 2053-2055, January – 2015
- [9] Jayakumar. P, Shankarakalidoss. G “Integral points on the Homogenous cone $x^2 + y^2 = 10z^2$ ”International Journal for Scientific Research and Development, Vol (2), Issue 11 , 234-235, January - 2015
- [10] Jayakumar.P, Prabha.S “Integral points on the cone $x^2 + 25y^2 = 17z^2$ ” International Journal of Science and Research,Vol (4), Issue 1, 2050-2052, January-2015
- [11] Jayakumar.P, Prabha.S “Lattice points on the cone $x^2 + 9y^2 = 26z^2$ ” International Journal of Science and Research, Vol (4), Issue 1 , 2050-2052, January – 2015
- [12] Jayakumar. P, Sangeetha. K, “Integral solutions of the Homogeneous Biquadratic Diophantine equation with six unknowns: $(x^3 - y^3)z = (W^2 - P^2)R^2$ ”International Journal of Science and Research, Vol (3), Issue 12, 1021- 1023 (December - 2014)

Author Profile

P. Jayakumar received the B. Sc , M.Sc degrees in Mathematics from Madras University in 1980 and 1983 and the M. Phil, Ph.D degrees in Mathematics from Bharathidasan University , Thiruchirappalli in 1988 and 2010. Who is now working as Associate Professor of Mathematics, A.V.V.M Sri Pushpam College Poondi (Autonomous), Thanjavur (District) – 613 503, Tamil Nadu, India.

G. Shankarakalidoss received the B. Sc , M.Sc, MPhil degrees in Mathematics from Bharathidasan University, Thiruchirappalli in 2002, 2004 and 2007. Who is now working as Assistant Professor of Mathematics, Kings College of Engineering, Punalkulam, Pudukkottai (Dist) Pin- 613303, India