

- [7] J.E. Randall, J.L. Justine, “*Cephalopholis aurantia* x *C. spiloparaea*, a hybrid serranid fish from New Caledonia,” *The Raffles Bulletin of Zoology*, LVI (1), pp. 157-159, 2008
- [8] J.E. Randall, A. Ben-Tuvia, “A review of the groupers (Pisces: Serranidae: Epinephelinae) of the Red Sea, with description of a new species of *Cephalopholis*,” *Bulletin of Marine Science*, XXXIII (2), pp. 373-426, 1983.
- [9] M. Shpigel, L. Fishelson, “Habitat partitioning between species of the genus *Cephalopholis* (Pisces, Serranidae) across the fringing reef of the Gulf of Aqaba (Red Sea). *Marine Ecology Program Series*, CDLVIII, pp. 17-25, 1989.
- [10] T.J. Donaldson, Partitioning behaviour and intra and interspecific interactions: a comparison between male and female groupers, *Cephalopholis spiloparaea* (Pisces: Serranidae: Epinephelinae), *Marine Biology*, CXXI (4), pp. 581-584, 1995.
- [11] T.J. Donaldson, Habitat association and depth distribution of two sympatric groupers of the genus *Cephalopholis* (Serranidae: Epinephelinae), *Ichthyological Research*, LIX, 191-193, 2002.
- [12] A.M. Abied, M.A. Abu El Regal, A.H. Khalil, “RAPD-PCR analysis of four coral reef fish species, genus *Cephalopholis* (Family: Serranidae) in the Red Sea,” *Egyptian Journal of Aquatic Biology and Fisheries*, XVIII (2), pp. 125-141, 2014.
- [13] K. Sujatha, “Taxonomic studies on three families of fishes represented in the fishery off Visakhapatnam,” Ph.D Thesis, Andhra University, India, 1981.
- [14] K. Sujatha, “Groupers off Visakhapatnam, north east coast of India,” *Journal of Marine Biological Association of India*, XLVI (1), pp. 87-92, 2004.
- [15] S. Rajapackiam, S. Mohan, P. Jaiganesh, S. Rajan, R. Vasu, “Heavy landing of tomato hind *Cephalopholis sonnerati* (Valenciennes, 1828) at Chennai”, *Marine Fisheries Information Service T&E Series*, CCX, 2011.
- [16] V. Sachithanandam, P.M. Mohan, “New distributional record of *Cephalopholis aurantia* Valenciennes, 1828 (Pisces: Serranidae) golden hind to Indian waters from Andaman and Nicobar Islands,” *Indian Journal of GeoMarine Sciences*, XLIII (12), pp. 30, 2014.
- [17] T.S.C. Roy, A. Gopalakrishnan, P.M.A. Muneer, L. John, K.K. Musammilu, V.S. Basheer, “Resolution of taxonomic ambiguity in groupers (Pisces: Serranidae) by the random amplified polymorphic DNA (RAPD) technique,” *Indian Journal of Fisheries*, LXI (2), pp. 28-34, 2014.
- [18] V.S. Basheer, N. Vineesh, K.K. Bineesh, R.G. Kumar, C. Mohitha, T.R. Swaminathan, C.P.R. Shanis, A. Gopalakrishnan, J.K. Jena, “Barcoding and phylogenetic inferences in grouper fish species from Indian waters,” in Abstracts, 10th Indian Fisheries and Aquaculture Forum, Lucknow, 2014.
- [19] R. Froese, D. Pauly, FishBase, World Wide Web electronic publication. www.fishbase.org (Accessed February 2014).
- [20] L.G.L. Lemus, “Biochemical systematic analysis of evolutionary relationships of groupers from the Gulf of California,” *Biochemical Systematics and Ecology*, XVI (1), pp. 79-87, 1988.
- [21] E. Mayer, P.D. Ashlock, *Principles of Systematic Zoology*, 2nd McGraw-Hill Incorporation, 1991.
- [22] R.S. Thorpe, “Review of the numerical methods for recognizing and analyzing racial variation,” in *Numerical Taxonomy*, J. Felsenstein (eds.), Springer-Verlag, Berlin, 1983.
- [23] J. Sambrook, E.F. Fritsch, T. Maniatus, *Molecular Cloning: A Laboratory Manual*, second edition, Cold Spring Harbor Laboratory Press, New York, 1989.
- [24] A. Gopalakrishnan, K.K. Lal, A.G. Ponniah, “Esterases in Indian major carps-rohu (*Labeo rohita*) and mrigal (*Cirrhinus mrigala*) (Teleostei, Cyprinidae),” *Indian Journal of Fisheries*, XLIV, pp. 361-368, 1997.
- [25] D.H. Whitmore, “Electrophoretic and Isoelectric Focusing Techniques in Fisheries Management”, CRC Press, Inc., Florida, USA, 1990.
- [26] J.B. Shaklee, F.W. Allendorf, D.C. Morison, G.S. Whitt, “Gene nomenclature for protein coding loci in fish,” *Trans American Fisheries Society*, CXIX, pp. 2-15, 1990.
- [27] M. Raymond, F. Rousset, GENEPop (Ver 1.2), A population genetics software for exact test and ecumenicism, *Journal of Heredity*, LVIII, pp. 248-249, <http://www.cefe.cnrs.mop.fr/genepop.html>
- [28] M. Nei, “Estimation of average heterozygosity and genetic distance from the small number of individual,” *Genetics*, LXXXIV, pp. 583-590, 1978.
- [29] F.C. Yeh, R.C. Yang, T.J.B. Boyle, Z.H. Ye, J.X. Mao, POPGENE, the user friendly software for population analysis, University of Alberta, Canada. Available at <http://www.ualberta.ca/~fyeh/index.html>.
- [30] I.A.A. Mekki, A.S. Mohammad, “Morphometrics and meristics of the three Epinepheline species: *Cephalopholis argus* (Bloch and Schneider, 1801), *Cephalopholis miniata* (Forsskal, 1775) and *Variola louti* (Forsskal, 1775) from the Red Sea, Egypt,” *Journal of Biological Sciences*, XI (1), pp. 10-21, 2011.
- [31] F.W. Allendorf, F.M. Utter, *Population Genetics, Fish Physiology*, VIII, Academic Press Incorporation, 1979.
- [32] T.T.C. Chan, Y. Sadovy. 2002. Reproductive biology, age and growth in the chocolate hind *Cephalopholis boenak* (Bloch, 1790), in *Hong Kong. Mar. and Fresh. Res.*, **53**, 791-803.
- [33] D.V. Rao, Kamala Devi, P.T. Rajan, “An account of Ichthyofauna of Andaman and Nicobar Islands, Bay of Bengal”, *Records of Zoological Survey of India Occasional Paper no CLXXVIII*, 2000.
- [34] P.T. Rajan, A field Guide to grouper and snapper fishes of Andaman and Nicobar Islands, *Zoological Survey of India, Kolkata*, pp. 1-102, 2002.
- [35] D.V. Rao, Guide to Reef fishes of Andaman and Nicobar Islands, *Zoological Survey of India, Kolkata*, 2004.
- [36] R.D. Ward, M. Woodwark, D.O.F. Skibinski, “A comparison of genetic diversity levels in marine, freshwater and anadromous fishes”, *Journal of Fish Biology*, XLIV, pp. 213-232, 1994.