

A Study to Assess the Knowledge and Practices Regarding Prevention of Anaemia among Antenatal Women Attending a Tertiary Level Hospital in Pune

Maj Sivapriya S¹, Lt Col Laxmipriya Parida²

¹Clinical Instructor, College of Nursing, CH (CC), Lucknow

²Associate Professor (Maternity Health Nursing), College of Nursing, A F M C, Pune

Abstract: *Background:* Anaemia is one of the potentially lethal complications of pregnancy leading to large number of maternal and foetal losses but it is preventable and curable disease. Antenatal women should equip themselves with adequate knowledge about the prevention of anaemia and routine beneficial diet, hygiene and treatment practices to prevent the consequence of anaemia in pregnancy. The aim of this study was to find out the level of knowledge and practices regarding prevention of anaemia antenatal women and the correlation of knowledge and practice level with selected variables such as age, educational qualification, occupation, obstetrical score, no of antenatal visits and haemoglobin at first visit. *Method:* Descriptive study was conducted on randomly selected 200 antenatal women of a selected tertiary level hospital of Pune. Knowledge and practice were analysed by structured interview schedule which included questionnaire on knowledge and self reported rating scale on practice. *Result:* 69% had good knowledge about prevention of anaemia. 59.5% antenatal women were following good practices to prevent anaemia in pregnancy. There is a significant positive correlation between knowledge and practice. There is a significant relation between knowledge and practice with age, educational qualification and family income of study participants. *Conclusion:* The current emphasis on health for all demands that every individual should be self sufficient and self reliant. Assessments of knowledge and practice and health education are essential step towards prevention of anaemia in pregnancy.

Keywords: Antenatal women, Knowledge, Practice, Prevention of anaemia, Tertiary care hospital.

1. Introduction

Anaemia in pregnancy is one of the leading causes responsible for maternal and perinatal morbidity and mortality. WHO has estimated that prevalence of anaemia among pregnant women is 14 % in developed and 51 % in developing countries. 65 -75% of population in India is affected with anaemia.

In India, anaemia is directly or indirectly responsible for 40 per cent of maternal deaths. There is 8 to 10 fold increase in MMR when the Hb falls below 5 g/dl. Early detection and effective management of anaemia in pregnancy can contribute substantially to reduction in maternal mortality. Maternal anaemia is associated with poor intrauterine growth and increased risk of preterm births and low birth weight rates.

The present study was descriptive in nature. The objectives of the study was to assess the knowledge and practice regarding prevention of anaemia among antenatal women, find the relationship between knowledge and practices regarding prevention of anaemia among antenatal women and assess the relationship between knowledge and practice with selected demographic variables.

2. Materials and Methods

Non experimental cross sectional descriptive study conducted in antenatal clinics of a selected tertiary care hospital in Pune. Hospital ethical committee permission was obtained. After taking informed consent 200 antenatal women were selected by systematic random sampling

method. Knowledge and practice was assessed using structured interview schedule which included questionnaire on knowledge and self reported rating scale to assess the practice regarding prevention of anaemia. Descriptive and inferential statistics was used to analyze the data

3. Results

The results were computed using descriptive statistics such as mean, median, standard deviation, and inferential statistics such as Z test, ANOVA and Man Whitney 'U' test and co – relation coefficient to find out the correlation between knowledge and practice regarding prevention of anaemia

(i) Sample distribution as per demographic and baseline variables

Majority of the subjects belonged to 21 – 25yrs, educated up to X standard and house wives. 96.5% antenatal women have done registration in the first trimester. 75% of the study participants had more than 4 antenatal visits. Only 47.5% had Hb more than 11 gm% at the time of registration. Only 1 .5% of the study participants has done de-worming within 6 months before the present pregnancy.

(ii) Knowledge regarding anaemia and its prevention

62(31%) antenatal women had average and 138 (69%) had good knowledge about prevention of anaemia. Most of the antenatal women, 128 (64%) had good knowledge and 71 (35.5%) had average knowledge on disease aspect of anaemia. 99% study participants had average or good knowledge on diet aspect of anaemia. All the antenatal

women had average or above average knowledge regarding prophylactic treatment of anaemia.




Figure 1: Distribution of antenatal women based on knowledge score

Table 1: Distribution of women based on knowledge about various important aspects of prevention of anaemia N =200

Various aspects of anaemia	No of antenatal women (f)	%
Anaemia is reduced level of Haemoglobin	122	61
Minimum Hb required during pregnancy is 11 gm/dl	97	48.5
Birth spacing should be minimum 2 yrs	189	94.5
Anaemia can be prevented by iron rich food and iron tablets	199	99.5
Lemon increases iron absorption	10	5
Jaggery is the richest source of iron	150	75
Iron tablets prevent anaemia	189	94.5
Continue iron tablets till 6 weeks after delivery	33	16.5
Antenatal registration has to be done within 3 months	199	99.5

(iii) Practice regarding anaemia and its prevention

All the antenatal women were in the category of average or good practice. 119 (59.5%) antenatal women were following good practices to prevent anaemia in pregnancy. Average practices were found in 81 (40.5%) antenatal women.




Figure 2: Distribution of practice score among antenatal women in study group

Detailed analysis revealed that majority of the women followed beneficial practices such as including green leafy vegetables and jaggery in diet, using iron utensils, hand washing before meals and after defecation, following regular antenatal visits and drug compliance. The data showed that

most of women were taking iron and calcium tablets together or with milk products. Only 1.5% women have done de-worming.

Table 2: Distribution of antenatal mothers according to practices regarding diet, n = 200

Diet Practices	Beneficial practices		Non beneficial practices	
	Frequency	Percentage	Frequency	Percentage
Made improvement in meal pattern in pregnancy	156	78	44	22
Use iron utensils for cooking	110	55	90	45
Includes jaggery in diet	142	71	58	29
Includes green leafy vegetables in diet	196	98	4	2
Skip meals	182	91	18	9
Habit of eating mud and charcoal	185	92.5	15	7.5
Avoid iron rich food	157	78.5	43	21.5

Table 3: Distribution of antenatal women according to practices regarding treatment, N =200

Treatment	No of antenatal women	Percentage
Regular antenatal visits	197	98.5
Intake of iron tablets	169	84.5
Take Vitamin C tablets along with iron tablets	3	1.5
Take iron tablets with lemon juice	1	0.5





Figure 3: Distribution of antenatal women according to non-beneficial treatment practices

67 (33.5%) of antenatal women were taking iron and calcium tablets together. 58 (29%) of women were taking iron tablets with milk products that inhibits absorption of iron.

(iv) Association between knowledge & practice regarding prevention of anaemia

There was significant positive correlation between knowledge and practice score among the subjects ($r = +0.41$) with $p < 0.01$ implying that as knowledge increased, the practice is also proportionally increased.



(v) Association between knowledge and practice with selected variables

There is significant relation between knowledge and practice with age, educational qualification, family income i.e., $p < 0.05$. But there was no statistically significant association between knowledge and practice with obstetrical score, number of antenatal visits and haemoglobin on registration as p value > 0.05

4. Interpretation and Conclusion

This was a cross sectional descriptive study to assess the knowledge and practice regarding prevention of anaemia among antenatal women. All the participants were having average and above knowledge and practice regarding diet and disease aspect. There is positive correlation between knowledge and practices among study subjects. Knowledge and practice comparison with selected demographic data strongly emphasize that there is increase in knowledge and practice according to age, education qualification and family income. No correlation of knowledge and practice was found with selected variables such as obstetrical score, no of antenatal visit, Hb during first visit in study group.

The findings were supported by a study conducted by Massawe S, Urass E, Lindmark G, Nystrom L in Dar-es-Salaam strongly support the present study findings that 88% of the subjects considered anaemia as a major health problem and 85% were aware of the causes and ways of preventing anaemia. 90% were aware of the advantages of early booking for antenatal care. The most frequently mentioned causes of anaemia were related to nutrition while intestinal parasites were mentioned by a few women.

The present study highlighted the importance of providing information to all the antenatal women in the clinics regarding beneficial and non beneficial practices to prevent anaemia in pregnancy. Even though drug compliance was good, faulty drug intake habits such as taking iron and calcium tablets together or with milk decreased the absorption and ultimately the desired outcome of the treatment. Only less number of the antenatal women has done de-worming before or during pregnancy. There should be mandatory preconception and antenatal counselling sessions of structured teachings to the antenatal women to identify the risk factors in pregnancy to reduce the enormous burden of anaemia in pregnancy.

The present study revealed that most of the antenatal women had average and good knowledge regarding diet and hygiene practices but they lag behind in knowledge and practice in prophylactic treatment to prevent anaemia. Hence the nurses must increase their efforts to impart more knowledge and create awareness on treatment to prevent anaemia in pregnancy. The current emphasis on health for all demands that every individual should be self sufficient and self reliant. There should be mandatory preconception and antenatal counselling sessions for the women to identify the risk factors in pregnancy and structured teachings to reduce the enormous burden of anaemia in pregnancy.

References

- [1] WHO. Worldwide prevalence of anaemia 1993 – 2005. (Cited 2014 Jan 20). Available from: http://apps.who.int/iris/bitstream/10665/43894/1/9789241596657_eng.pdf.
- [2] Dr Samar K Basu. Anaemia in pregnancy. Health matters (document on the Internet) (cited 2014 Jan 10).
- [3] Available from delhimedicalcouncil.nic.in/anemiainpregnancy.pdf
- [4] United Nations Children's Fund, United Nations University, WHO. Iron deficiency anaemia, assessment, prevention and control a guide for programme managers. 2001.
- [5] World Health Organisation. Antenatal care. (Cited 2014 Jan 13). Available from <http://www.who.int/pmnch/media/publications/aonsectionIII.2pd>
- [6] Park K. Preventive and social medicine. 22nd edition. Jabalpur: M/s Banarsidas Bhanot publishers; 2013. Page no 840
- [7] Planning Commission. Tenth Five-Year Plan. Sectoral Policies and Programmes. Nutrition. Government of India. New Delhi. 2003-2007. [cited 2013 Mar 15] Available at: http://planningcommission.nic.in/plans/planrel/fiveyr/10th/volume2/v2_ch3_3.pd f.
- [8] Sharma Anshu, Patnaik Rita, Garg Suman, Ramachandran Prema. Detection and management of anaemia in pregnancy in urban primary health care institution. Indian J Med Res. 2008; 128: 45 -51.
- [9] Agarwal K N, Agarwal D K, Sharma A, Sharma K, Prasad K, Kalita MC, et al Prevalence of anaemia in pregnant & lactating women in India . Indian J Med Res. 2004; 124 (2): 173 – 184.

- [10] National Family Health Survey-India 1998-99. 2003. Bombay: International Institute of Population Studies
- [11] Survey on Maternal Mortality Ratio. Report of Registrar General of India Sample Registration System (RGI –SRS) for 2007 -2009. Available from : [censusindia.gov.in/vital statistics/SRS_Bulletins/MMR](http://censusindia.gov.in/vital_statistics/SRS_Bulletins/MMR)
- [12] Global burden of disease report WHO 2002. Available from: www.who.int/healthinfo/globalburden_disease/estimates_regional_2002.
- [13] Bruno Banoist, McLean Erin, Ines Egli, Cogswell Mary. WHO global database on anaemia. Worldwide prevalence of anaemia 1993 – 2005.
- [14] Kalaivani K. Prevalence and consequences of anaemia in pregnancy. Indian J Med Res. 2009; 130: 627 -633
- [15] National Family Health Survey of India – 2.
- [16] Available from: www.rchiips.org/nfhs/nfhs2.s
- [17] Nugraheni S A, Dasuk D, Ismail D. Knowledge, attitude and practice of pregnant women in correlation with anaemia. 2004 (unpublished).
- [18] George Julia B. Nursing theories-the base for professional nursing practice. 6th edn. South Asia: Dorling Kindersley (India) Pvt Ltd; 2011.
- [19] Marriner TA, Raile AM. Nursing theorists and their work. 5th ed. Sakraida T.Nola J. Pender. The Health Promotion Model. St Louis: Mosby; 2005
- [20] Suresh K Sharma. Nursing research and statistics. I edition. Elsevier publishers:2010.
- [21] Polit D F, Beck C T. Nursing research: generating and assessing evidence for nursing practice. 8th edition. New Delhi. Wolters Kluwer: 2008.
- [22] The safe motherhood Initiative and beyond, WHO 2007. Available from: www.who.int/bulletin/volumes/85/10/07-045963/en.
- [23] The global burden of disease: 2004. www.who.int/healthinfo/global_burden_of_disease/GBD_report_2004update.
- [24] K N Agarwal, D K Agarwal, Anshu Sharma. Anaemia in pregnancy interstate differences project funded by dept of Family welfare, GOI, Nutrition foundation of India
- [25] Kumar Pankaj, Pore Prasad, Patil Usha. Maternal anaemia and its impact on perinatal outcome in a tertiary care hospital of Pune, Maharashtra. Indian Journal of Basic & Applied Medical Research. 2012; 2(1): 111 -119.
- [26] Ghimire N, Pandey N. Knowledge and practice of mothers regarding the prevention of anaemia during pregnancy in teaching hospital, Kathmandu. Journal of Chitwan Medical College 2013; 3(5):14 -17.
- [27] Rae Galloway, Erin Dusch, Lesilie Elder, Endang Achadi, Ruben Grajeda, Jolly Rajratnam, et al. women's perception of iron deficiency anaemia and prevention and control in eight developing countries. Mother care project. 1991 -1998
- [28] Howyida S. Abd ElHameed1, Aziza I. Mohammed and Lamiaa T. Abd El Hameed. Effect of Nutritional Educational Guideline among pregnant women with Iron deficiency anaemia at Rural Area in Kalyobia. Life Science Journal. 2012; 9(2):1212-1216.
- [29] Sarin Jyoi, Podder Lily, Kumari Right, N Sujata. A study to determine the prevalence of anemia among registered antenatal mothers in MMIMS & R antenatal OPD with a view to develop and evaluate a planned health education programme on prevention and management of anaemia in pregnancy in terms of knowledge and practices of antenatal mothers. Nursing and Midwifery Research Journal.2009.4 (2).39-50.
- [30] Ailinger L Rita, Jean B Moore, Lisa Pawloski, Lidya Ruth Zamora. Concepts of anaemia among low income Nicaraguan women. Revista Latino- American de Enfermagem.2009.17 (2).
- [31] Massawe S, Urassa E, Lindmark G, Nystrom L. Anaemia in pregnancy: perception of patients in Dar-es-Salaam. East Afr Med J. 1995 Aug 72(8); 498 -503
- [32] Myrada & St John's Medical College Bangalore.NRHM. Study of anaemia prevention and control among adolescent girls and pregnant women in three districts of karnataka: how sustainable ? how effective?. 2010. (unpublished)
- [33] Kalimbira A A, Mtimuni B M, Chilima DM. maternal knowledge and practices related to anaemia and iron supplementation in Rural Malawi : A cross sectional study. African Journal of food agriculture nutrition and development. 2009;9(1): 550 -564.
- [34] Mbule M A, Byaruhanga Y B, Kabahenda M, Lubowa A. Determinants of anaemia among pregnant women in rural Uganda. The international electronic journal of rural health and education. 2013; 13 (2259). Available at [http:// www.rrh.org.au](http://www.rrh.org.au).
- [35] Marwan Shehda Salama Mosleh. Awareness of anaemia among pregnant women at unrwa clinics in gaza strip. M Sc in Public health (Dissertation) Umeda international school of public health Sweden: Umea university 2010 (unpublished study)
- [36] Derbyshire E. Iron deficiency anaemia in pregnancy – what are the solutions?. Nutrition and Food Science. 2010; 40 (6): 5.
- [37] Kaur Manmeet , Singh Kamaljeet. Effect of health education on knowledge, attitude and practices about anaemia among rural women in Chandigarh. Indian journal of community medicine. 2001; 26 (3):
- [38] Luzine Mirzoyan. Iron deficiency anaemia in pregnancy: Assessment of knowledge, attitudes and practices of pregnant women. Department of public health, American university of Armenia (Unpublished)
- [39] Manna Prabir kumar, De Debasis, Ghosh Ddebidas. Knowledge attitude and practices for antenatal care and delivery of the mothers of tea garden in Jalpaiguri and Darjeeling districts, West Bengal. National journal of community medicine. 2011 ; 2(1): 4 -8
- [40] Gautam Virender, Bansal Yogesh, Taneja D K, Saha Renuka. Prevalence of anaemia amongst pregnant women and its socio demographic associates in a rural area of Delhi. Indian Journal of Community Medicine. 2002; 27 (4): 157 -160.
- [41] Viveki1 R G, Halappanavar A B, Viveki P R, Halki1 S B, Maled V S, Deshpande P . Prevalence of anaemia and its epidemiological determinants in pregnant women. Al Ame en J Med S c i. 2 012; 5 (3):216 -223.
- [42] Kerlinger N Fred. Foundations of behavioural research. 3rd edition. New York. Orlando Harcourt Brace and Company:1986; 15-25
- [43] Abdellah F G. Better patient care through nursing research. 2nd edition. New York. Macmillan company: 1979: 208 -209.

- [44] Joint statement by WHO & UNICEF. Focusing on anaemia. Towards an integrated approach for effective control of anaemia. Available from: WHO home page: <http://www.who.in>.
- [45] K S Saraswathi, Farhana Aljabri, R Shyamala. Prevalence of anaemia among antenatal women in a tertiary care hospital, South India. *Der Pharmacia Lettre*. 2013. 5(1): 146 -148.
- [46] Ahmad N, Kalakoti P, Bano R, Syed. The prevalence of anaemia and associated factors in pregnant women in a rural Indian community. *Australian Medical Journal*, 2010,3,5,276-280
- [47] V Raghuram , Anil Manjula, S Jayaram. Prevalence of anaemia amongst women in the reproductive age group in a rural area in south India. *Int J Biol Med Res*. 2012;3(2):1482 -1484.
- [48] Amer Mariam, Faten Khayarat, Nawal Mahmoud Soliman. Effect of nutritional interventions on anaemic pregnant women's health using health promotion model. Ain Shams University, Palastine. (Unpublished).