

# Study of Plasma Homocysteine Level in Coronary Heart Disease Patients and Compare with Control

Sharma Hemlata<sup>1</sup>, Vyas Shalini<sup>2</sup>, Dr. Vyas R. K.<sup>3</sup>, Dr. Chhparwal Amit<sup>4</sup>

<sup>1,2</sup>M.Sc.(medicine) Biochemistry, Biochemist, Department of Biochemistry, S.P. Medical college, Bikaner, India

<sup>3</sup>PhD (medicine) Biochemistry, Professor & head, Department of Biochemistry, S.P. Medical College, Bikaner, India

<sup>4</sup>M.D.S., Assistant Professor, Geetanjali dental research institute, Geetanjali medical college, Udaipur, India

**Abstract:** The study was conducted on 50 patients of coronary heart disease and 50 persons are healthy subjects between the age group 25-70 years of both sex. Homocysteine may increase cellular oxidative stress in which mitochondrial thioredoxin, and peroxidation are decreased and NADH oxidase activity is increased and generating peroxynitrite and nitrotyrosine in contractile protein which causes vascular dysfunction. Plasma homocysteine were measured by hplc grade kit with the help of HPLC. plasma homocysteine level of coronary heart disease patients showed a highly significant ( $p < 0.0001$ ) relationship. Estimation of plasma homocysteine is reliable, economic and sensitive and it can be used in the proper management of chronic complications of coronary heart disease.

**Keywords:** Coronary heart disease, Homocysteine, HPLC, oxidative stress

## 1. Introduction

The cardiovascular disease (CVD) burden of India is expected to double in the next two decades, making it the single largest cause of death and the second largest cause of disability by the year 2020<sup>[1]</sup>

This will be characterized by an enormous burden of CVD among urban communities. Further, the prevalence of CVD in rural and semi-urban areas is expected to increase substantially<sup>[2,3]</sup>. Coronary heart disease also known as coronary artery disease or Ischemic heart disease) is the most common form of heart disease and results from atherosclerosis or the accumulation of fatty plaques in artery walls that causes narrowing of artery lumen<sup>[4]</sup>.

Hyperhomocysteinemia has been strongly associated with the pathogenesis of coronary vascular disease, and correspondingly has been identified as a contributing factor in four main disease mechanisms including thrombosis, vascular oxidative stress, apoptosis and cellular proliferation.<sup>[5-7]</sup>

## 2. Material & Methods

The present study was conducted on 50 patients of coronary heart disease of HRMC, Bikaner and 50 persons are healthy subjects between the age group 25-70 years of both sexes.

### Determination of plasma homocysteine

For the determination of homocysteine, the sample is reduced and derivatized in one step. The albumin bound and the oxidized homocysteine is reduced. During a precipitation step high molecular substances are removed and analysed by high performance liquid chromatography with uv detector.

## 3. Results

The plasma homocysteine level was found to be  $30.42 \pm 10.10 \mu\text{mol/L}$  with a range of 16.04-46.66  $\mu\text{mol/L}$  in CHD. The increase was statistically highly significant as compared to control group with  $14.95 \pm 14.95 \mu\text{mol/L}$ ; while it ranged from 6.80 to 21.51  $\mu\text{mol/L}$  as evident by P-value ( $P < 0.0001$ )

### Plasma Homocysteine concentration ( $\mu\text{mol/L}$ ) in CHD subjects with that of control.

S.No	Values	Control group	CHD group
1	Mean	14.955	30.426
2	Range	6.80-21.51	16.04-46.66
3	SD	4.16	10.10
4	SE	0.587	1.428
5	DF		98
6	T		10.014
7	P-value		0.0001***

\* Significant

\*\*\*Highly Significant

Df = Degree of Freedom

## 4. Discussion & Conclusion

Plasma homocysteine level varied from 6.8 to 18.7  $\mu\text{mol/L}$  with mean as  $12.92 \pm 3.61 \mu\text{mol/L}$  in normal control male subject as shows in the table-X the values of homocysteine obtained in the present series of study in normal control subjects were in close agreement with the findings of coldea et al (2011)<sup>[7]</sup>

In this study, a highly significant correlation was observed when when plasma homocysteine of CHD subjects was compared with healthy subjects. The increase Plasma homocysteine concentration in CHD patients was statistically significant as compared to normal control more subjects with age difference as evident by P-Value ( $P < 0.001$ ) These values are in close collaboration with the findings of Klerk et al (2002)<sup>[8]</sup>

Estimation of plasma homocysteine is simple, reliable, economic and sensitive and it can be used in the proper management of chronic complications coronary heart disease.

## References

- [1] Murray CJL, Lopez AD. Global health statistics: Global burden of disease and injury series. Vols. I and II. Boston:Harvard School of Public Health; 1992.
- [2] Gupta R, Gupta VP. Meta-analysis of coronary heart disease prevalence in India. Indian Heart J 1996;48:241–5
- [3] Krishnaswami S, Joseph G, Richard J. Demands on tertiary care for cardiovascular diseases in India: Analysis of data for 1960–89. Bull World Health Organ 1991;69:325–30.
- [4] Cheek D. What's the difference about heart disease in women? Nursing,2003;33:8:36-42.
- [5] Al-Obaidi- M K, P J Stubbs, R Amersey, M I M Noble. (2001), Acute and convalescent changes in plasma homocysteine concentrations in acute coronary syndromes, Heart 2001;85:380.384.
- [6] Lubos, E., Lasclazo, J., & Handy, D. : Homocysteine and glutathione peroxidase-1. Antioxid. Redox. Signal , 2007;19:23-1940
- [7] Coldea Agoston - Lucia, Teodora Mocan, Marc Gafosse, Silvia Lupu, Dan L. Dumitrascu Plasma homocysteine and the severity of heart failure in patients with previous myocardial infarction Cardiol J; 2011;18, 1: 55.62.
- [8] Klerk M, Verhoef P, Clarke R et al MTHFR studies collaboration group MTHFR 677CT poly morphism and risk of coronary heart disease meta analysis. JAMA 2002;288:2023-2031.

## Author Profile

**Hemlata Sharma** is biochemist, Department of Biochemistry in S.P. Medical College, Bikaner (Rajasthan), India. she had also worked as a senior demonstrator Department of Biochemistry in Dr. S.N. Medical College, Jodhpur (Rajasthan), India.