Common Fixed Point Theorems Weak Compatible in Cone Metric Spaces

Raghu Nandan Patel¹, Damyanti Patel²

¹Department of Mathematics, Government Naveen Colleeg, Balrampur (C. G.), India

² Department of Mathematics, Government Engineering College, Bilaspur (C.G.), India

Abstract: In this paper we established common fixed point theorems for weakly compatible mappings in cone metric spaces.

AMS Subject Classification: 47H10, 54H25

Keywords: fixed point, cone metric spaces and weakly compatible mappings.

1. Introduction

The Banach contraction mapping principle is widely recognized as the source of metric fixed point theory. This contraction principle has further several generalizations in metric spaces as well as in cone metric spaces. Huang and Zhang [1] introduced the concept of cone metric space, where every pair of elements is assigned to an element of a Banach space and defined a partial order on the Banach space with the help of a subset of the Banach space called cone which satisfy certain properties.

2. Preliminary Notes

First, we recall some standard definitions and other results that will be needed in the sequel.

Definition 2.1. Let E be a real Banach space and P be a subset of E. P is called a cone if

(1) P is closed, nonempty and $P \neq \{0\}$;

(2) $a, b \in R, a, b \ge 0, x, y \in P \Longrightarrow ax + by \in P$;

(3) $x \in P$ and $-x \in P \Longrightarrow x = 0$.

Given a cone $P \subseteq E$, we define a partial ordering " \leq " in E by $x \leq y$ if $x - y \in P$. We write x < y to denote $x \leq y$ but $x \neq y$ and $x \ll y$ to denote $y - x \in P^0$, where P^0 stands for the interior of P. We assume cone is solid i. e. that $P^0 \neq \phi$.

Proposition 2.2 [7] : Let P be a cone in a real Banach space E.

(1) If $a \in P$ and $a \le ka$, for some $k \in [0, 1)$ then a = 0. (2) If $a \in P$ and $a \ll c$, for all $c \in P^0$ then a = 0.

A cone P is called normal if there is constant K > 0 such that, for all $x, y \in E$. $0 \le x \le y \Rightarrow ||x|| \le K||y||$. The least value of constant K satisfying this inequality is called the normal constant of P.

Definition 2.3 [1]: Let X be a nonempty set and E be a real Banach space. Suppose that the mapping d: X x $X \rightarrow E$ satisfies

(1) $0 \le d(x, y)$, for all $x, y \in X$ and d(x, y) = 0 if and only if x = y;

(2) d(x, y) = d(y, x), for all $x, y \in X$;

(3) $d(x, y) \le d(x, z) + d(z, y)$, for all $x, y, z \in X$

Then d is called a cone metric on X and (X, d) is called a cone metric space.

For examples of cone metric spaces we refer [1, 4]. Henceforth unless otherwise indicated, P is a normal cone in real Banach space E and " \leq " is partial ordering with respect to P.

Definition 2.4 [1]: Let (X, d) be a cone metric space. Let $\{xn\}$ be a sequence in X and $x \in X$.

(a) If for every $c \in E$ with $0 \ll c$ (or equivalently $c \in P^0$) there is positive integer n_0 such that for all $n > n_0$, $d(x_n, x) \ll c$ then the sequence $\{x_n\}$ converges to x. We denote this by $x_n \to x$, as $n \to \infty$ or $\lim_{n\to\infty} x_n = x$.

(b) If for every $c \in E$ with $0 \ll c$ there is positive integer n_0 such that for all $n, m > n_0, d(x_n, x_m) \ll c$ then the sequence $\{x_n\}$ is called a Cauchy sequence in X.

(X, d) is called a complete cone metric space, if every Cauchy sequence in X is convergent in X.

Lemma 2.5 [1]: Let (X, d) be a cone metric space, P be a normal cone with normal constant K. Let $\{x_n\}$ be a sequence in X, then $\{x_n\}$ converges to x if and only if $d(x_n, x) \rightarrow 0$ (n $\rightarrow \infty$).

Lemma 2.6 [1]: Let (X, d) be a cone metric space, P be a normal cone with normal constant K. Let $\{x_n\}$ be a sequence in X, then $\{x_n\}$ is a Cauchy sequence if and only if $d(x_n, x_m) \rightarrow 0$ $(n, m \rightarrow \infty)$.

Definition 2.7: A function $f : P \to P$ is called subadditive if for all $x, y \in P$, $f(x + y) \le f(x) + f(y)$.

Definition 2.8: If Y be any partially ordered set with relation " \leq " and Ψ : Y \rightarrow Y , we say that Ψ is non decreasing if x, y \in Y, x \leq y \Rightarrow Ψ (x) \leq Ψ (y).

Definition 2.9 [2]: Let $\Psi : P \rightarrow P$ be a vector valued function then Ψ is called MS-Altering function if

(a) Ψ is non decreasing, subadditive, continuous and sequentially convergent;

(b) $\Psi(a) = 0$ if and only if a = 0.

We replace conditions (a) and (b) by weaker conditions and define cone altering function as follows

Definition 2.10: Let $\Psi: P \to P$ be a vector valued function then Ψ is called cone altering function if

(a) Ψ is non decreasing, subadditive;

(b) $\Psi(a_n) \to 0$ if and only if $a_n \to 0$, for any sequence $\{a_n\}$ in P.

Note that for cone altering function Ψ on normal cone P, $\Psi(a) = 0$ if and only if a = 0.

Definition 2.11: Let X be any nonempty set, f, $g : X \to X$ be mappings. A point $w \in X$ is called point of coincidence of f and g if there is $x \in X$ such that fx = gx = w.

Definition 2.12: Let X be any nonempty set, f, $g : X \to X$ be mappings. Pair (f, g) is called weakly compatible if $x \in X$, $fx = gx \Rightarrow fgx = gfx$.

Lemma 2.13: Let (X, d) be a cone metric space and P be a normal cone in a real Banach space E, Ψ is a cone altering function and k_1 , k_2 , k > 0. If $x_n \to x$, $y_n \to y$ in X and ka $\leq k_1 \Psi [d(x_n, x)] + k_2 \Psi [d(y_n, y)]$, then a = 0.

Lemma 2.14: Let X be any nonempty set and f, $g : X \to X$ be mappings. If (f, g) is weakly compatible pair and have a unique point of coincidence then it is unique common fixed point of f and g.

3. Main Results

Theorem 3.1: Let (X, d) be a cone metric space with normal cone P and let A, S and T be self mappings of X, Ψ : $P \rightarrow P$ is cone altering function such that

 $(3.1.1) A(X) \subseteq S(X) \cap T(X)$

(3.1.2) the pairs { A, S} and {A, T} be weakly compatible.

(3.1.3i) there exist for all $x, y \in X$

 $\begin{array}{ll} \Psi[d(Ax, \quad Ay)] &\leq \quad a_1 \Psi[d(Sx, \quad Ty)] + a_2 \Psi[d(Ax, \\ Sx)] + a_3 \Psi[d(Ay, Ty)] + a_4 \Psi[d(Ax, Ty)] \end{array}$

 $+a_5\Psi[d(Ay, Sx)]$

where a_i , i = 1, 2, 3, 4, 5 are nonnegative constant such that $a_1+a_2+a_3+a_4+a_5 < 1$. Then there exists a unique point $w \in X$ such that Aw = Sw = w and a unique point $z \in X$ such that Az = Tz = z. Moreover z = w, so that there is a unique common fixed point of A, S and T.

Proof: Let the pairs $\{A, S\}$ and $\{A, T\}$ be weakly compatible so there are points $x, y \in X$ such that Ax = Sx and Ay = Ty. We claim that Ax = Ay. If not, by inequality (3.1.3)

$$\begin{split} \Psi[d(Ax, Ay)] &\leq a_1 \Psi[d(Sx, Ty)] + a_2 \Psi[d(Ax, Sx)] + a_3 \Psi[d(Ay, Ty)] + a_4 \Psi[d(Ax, Ty)] \end{split}$$

$$\begin{split} +a_5\Psi[d(Ay,\,Sx)]\\ &=a_1\Psi[d(Ax,\,Ay)]+a_2\Psi[d(Ax,\,Ax)]+a_3\Psi[d(Ay,\,Ay)]+a_4\Psi[d(Ax,\,Ay)] \end{split}$$

$$+a_5\Psi[d(Ay, Ax)]$$

= (a₁+a₄+a₅) $\Psi[d(Ay, Ax)]$

since $(a_1 + a_4 + a_5) < 1$ hence by proposition 2.2, we have $\Psi[d(Ax, Ay)] = 0$ i.e. d(Ax, Ay) = 0 or Ax = Ay.

Therefore Ax = Sx = Ay = Ty. Suppose that there is a another point z such that Az = Sz then by (3.1.3) we have Az = Sz = Ay = Ty, so Ax = Az and w = Ax = Sx is the unique point of coincidence of A and S. By Lemma 2.14 w is the only common fixed point of A and S. Similarly there is a unique point $z \in X$ such that z = Az = Tz.

$$\begin{split} & \text{Assume that } w \neq z. \text{ We have} \\ & \Psi[d(w, z)] = \Psi[d(Aw, Az)] \\ & \leq a_1 \Psi[d(Sw, Tz)] + a_2 \Psi[d(Aw, Sw)] + a_3 \Psi[d(Az, Tz)] + a_4 \Psi[d(Aw, Tz)] \end{split}$$

$$+a_5\Psi[d(Az, Sw)]$$

 $= a_1 \Psi[d(w, z)] + a_2 \Psi[d(w, w)] + a_3 \Psi[d(z, z)] + a_4 \Psi[d(w, z)]$

$$+a_5\Psi[d(z, w)]$$

 $= (a_1 + a_4 + a_5) \Psi[d(z, w)]$

since $(a_1 + a_4 + a_5) < 1$ hence by proposition 2.2, we have $\Psi[d(w, z)] = 0$ i.e. d(w, z) = 0 or w = z by Lemma 2.14 and z is a unique common fixed point of A, S and T.

Theorem 3.2: Let (X, d) be a cone metric space with normal cone P and let A, B, S and T be self mappings of X, $\Psi : P \rightarrow P$ is cone altering function such that

(3.2.1) $A(X) \subseteq S(X)$ and $B(X) \subseteq T(X)$

(3.2.2) the pairs { A, S} and {B, T} be weakly compatible. (3.2.3) there exist for all $x, y \in X$

 $\begin{array}{rl} \Psi[d(Ax, \ By)] &\leq a_1 \Psi[d(Sx, \ Ty)] + a_2 \Psi[d(Ax, Sx)] + a_3 \Psi[d(By, Ty)] + a_4 \Psi[d(Ax, Ty)] \\ &\quad + a_5 \Psi[d(By, Sx)] \end{array}$

where a_i , i = 1, 2, 3, 4, 5 are nonnegative constant such that $a_1+a_2+a_3+a_4+a_5 < 1$. Then there exists a unique point $w \in X$ such that Aw = Sw = w and a unique point $z \in X$ such that Bz = Tz = z. Moreover z = w, so that there is a unique common fixed point of A, B, S and T.

Proof: Let the pairs $\{A, S\}$ and $\{B, T\}$ be weakly compatible so there are points $x, y \in X$ such that Ax = Sx and By = Ty. We claim that Ax = By. If not, by inequality (3.2.3)

$$\begin{split} \Psi[d(Ax, By)] &\leq a_1 \Psi[d(Sx, Ty)] + a_2 \Psi[d(Ax, Sx)] + a_3 \Psi[d(By, Ty)] + a_4 \Psi[d(Ax, Ty)] \end{split}$$

$$+a_5\Psi[d(By, Sx)]$$

 $=a_1\Psi[d(Ax,\ By)]+a_2\Psi[d(Ax,\ Ax)]+a_3\Psi[d(By,\ By)]+a_4\Psi[d(Ax,\ By)]$

$$+a_5\Psi[d(Av, Bx)]$$

$$= (a_1 + a_4 + a_5) \Psi[d(Bv, Ax)]$$

since $(a_1 + a_4 + a_5) < 1$ hence by proposition 2.2, we have $\Psi[d(Ax, By)] = 0$ i.e. d(Ax, By) = 0 or Ax = By.

Therefore Ax = Sx = By = Ty. Suppose that there is a another point z such that Az = Sz then by (3.2.3) we have Az = Sz = By = Ty, so Ax = Az and w = Ax = Sx is the unique point of coincidence of A and S. By Lemma 2.14 w is the only common fixed point of A and S. Similarly there is a unique point $z \in X$ such that z = Bz = Tz.

Assume that $w \neq z$. We have

 $\Psi[d(w, z)] = \Psi[d(Aw, Bz)]$

$$\leq a_1 \Psi[d(Sw, Tz)] + a_2 \Psi[d(Aw, Sw)] + a_3 \Psi[d(Bz, Fz)] + a_4 \Psi[d(Aw, Tz)] + a_5 \Psi[d(Bz, Sw)]$$

 $= a_1 \Psi[d(w, z)] + a_2 \Psi[d(w, w)] + a_3 \Psi[d(z, z)] + a_4 \Psi[d(w, z)]$

 $+a_5\Psi[d(z, w)]$

 $= (a_1 + a_4 + a_5) \Psi[d(z, w)]$

since $(a_1 + a_4 + a_5) < 1$ hence by proposition 2.2, we have $\Psi[d(w, z)] = 0$ i.e. d(w, z) = 0 or w = z by Lemma 2.14 and z is a unique common fixed point of A, B, S and T.

References

- L.G. Huang, X.Zhang, Cone Metric Spaces and Fixed Point Theorems of contractive mappings, J. Math.Anal. Appl., 332, (2007), 1468-1476.
- [2] S. K. Malhotra, S Shukla and R. Sen Common Fixed Point Theorems in Cone Metric Space by Altering Distances; International Mathematical Forum, 54(6) (2011), 2665 – 2671.
- [3] S. K. Malhotra, S Shukla and R. Sen Common Fixed Point Theorems in Cone Metric Space by Altering Distances; Mathematica Aeterna, 1(6) (2011), 347 – 352.
- [4] Sh. Rezapour, R. Hamlbarani, Some notes on the paper Cone Metric spaces and fixed [point theorems of contractive mappings, J. Math. Anal. Appl. 345(2008), 719-724.
- [5] A.K.Dubey and A. Narayan, Cone Metric Spaces and Fixed Point Theorems for Pair of contractive maps, Mathematica Aeterna, Vol.2,2012,no.10,839-845.
- [6] A.K Dubey Rita Shukla, R.P. Dubey, An Extension of the Paper "Cone Metric Spaces and Fixed Point Theorems of Contractive Mappings", Int. J. of Applied Mathematical Research, 2(1), (2013), 84-90.
- [7] D. Ilic, V Rakocevic, Qusai-contraction on a cone metric space, applied Mathematics Letter, 22(5) (2009), 728 – 731.

Author Profile



Dr Raghu Nandan Patel is Assistant Professor, Department of Mathematics, Govt, Naveen College Balrampur (C. G.) India



Dr Damyanti Patel is Lecture in Department of Mathematics, Government Engineering College, Bilaspur (C. G.) India