

Life form Composition and Biological Spectrum of Ramnagar Wildlife Sanctuary, J&K, India

Sehrish Gazal¹, Anil K. Raina²

^{1,2}University of Jammu, Department of Environmental Sciences, Babasaheb Ambedkar Road, Jammu- 180 006, India

Abstract: *The present study has been carried out to understand the life form composition and Biological spectrum of Ramnagar Wildlife Sanctuary, Jammu and Kashmir. A total of 259 tracheophytes have been listed and grouped into various life-form classes. Therophytes (34.36%), Macrophanerophytes (23.16%), Nanophanerophytes (11.58%) and Chamaephytes (10.81%) are the major life-form classes present in the area. Hemicryptophytes (8.88), Lianas (8.11%), Cryptophytes (2.31%) and Epiphytes (0.77%) were observed to have low occurrence. The biological spectrum of the area was prepared and compared to Raunkiaer's Normal biological spectrum as well the spectra of the adjacent regions. Phytoclimate of the area was determined as Thero-phanerophytic, as per the Raunkiaer's terminology, with the preponderance of therophytes and phanerophytes. The study reflects the impact of various biotic factors on overall vegetation structure and composition of the area.*

Keywords: Biological spectrum, Life form, Raunkiaer, Phytoclimate, Ramnagar wildlife sanctuary.

1. Introduction

The vegetation may be classified according to growth-form [1]–[3] or life-form which refers to the sum total of adaptations to a specific climate manifesting a specific form and structure [4] and is reflected in the physiognomy or general appearance of the vegetation. Expression of the percentage distribution or relative proportion of different life-forms in the flora of the region is known as “the spectrum of life-forms” [5] or “biological spectrum”. Raunkiaer [3] defined life-forms as „the sum of the adaptation of a plant to climate” and classified the plant species into various life-forms based upon the principle of position and degree of protection of the buds during adverse conditions. This system for the description and classification of plant life forms is the most accepted from a number of different systems devised by many ecologists [6]–[8]. According to Raunkiaer's system [3], plant species can be grouped into five main classes, arranged according to increased protection of the renewing buds: phanerophytes, chamaephytes, hemicryptophytes, cryptophytes and therophytes. Raunkiaer [3] proposed the “biological spectrum” to express both the life-form distribution in a flora and the phytoclimate under which the prevailing life-forms evolve. The biological spectrum is the percent representation of the number of species belonging to each life-form in a given flora. Raunkiaer [3] constructed a „normal spectrum” which could act as a null model against which different life form spectra could be compared. Raunkiaer's normal spectrum indicates a phanerophytic community for the world and deviation from it determines the phytoclimate of the habitats. The occurrence of similar biological spectra in different regions indicates similar climatic conditions. Differences in the life-form distribution between the normal spectrum and a biological spectrum would point out which life-form characterizes the phytoclimate or the vegetation understudy [9].

Biological spectrum can be used to calculate numerical results quickly [10] to indicate the stratification and layering pattern of a community [11], to indicate the prevailing environment [12], its aridity or humidity [13] to monitor the impact of ambient stress factors on climate [4] and to

determine the nature of bioclimate or phytoclimate. Biological spectra are useful in comparing geographically widely separated plant communities and are also regarded as indicators of biotic interaction, climate and habitat deterioration [14].

Life-form studies are important to find phytoclimate of the area as structure and rate of change of composition are sensitive indicators of whole environment. The biological system is thus useful as an index of the health status of a forest. When worked out at periodic intervals, biological spectrum may set the guidelines for eco-restoration and optimization of a community. In view of this, the present work was under taken in the Ramnagar Wildlife Sanctuary, located in lower Shiwalik range, to study the assemblage of different life-forms and to prepare the biological spectrum to infer the existing phytoclimate of the area.

2. Materials and Methods

The Wildlife Sanctuary is situated between 32°44' to 32°47' North latitude and 74°50' to 74°53' East longitude with altitude varying from 270 m to above 520 m asl. The total area of the Sanctuary is about 706 hectare with an undulating topography which is crisscrossed by a number of seasonal streams (locally called as khads). The Sanctuary is traversed by Jammu-Srinagar National highway which enters in south and leaves in north-east, dividing it into two unequal segments. A trail road also passes through the sanctuary and is being used by morning walkers. The area experiences great extremities of temperature with June recorded as hottest and January as coldest month with average maximum and minimum temperature of 45.3°C and 2.9°C, respectively. Average rainfall of area is 115 cm, of which 80% occurs during monsoon. The life-form classification and construction of the biological spectra of the area was carried out after detailed floristic investigation of the area during 2011 to 2013. Only seed plants and ferns (Angiosperms, Gymnosperms and Pteridophytes) were taken into account. For the purpose of identification, plants were photographed on the spot with due care taken to avoid any untoward loss or perturbation to the habitat. Various local and other floras viz. Flora of Jammu [15] and herbarium of department of Botany,

University of Jammu were used besides consulting taxonomic experts of the region. The form and nature of perennating buds of plant species were recorded and worked out according to the life-forms system of Raunkiaer [3]. The percentage distribution of these species in different life-forms was calculated for construction of biological spectrum of the area. The values, thus determined, were compared with the normal spectrum given by Raunkiaer [3]. The percentage life-form was calculated as follows;

$$\% \text{ Lifeform} = \frac{\text{Number of species in any life form}}{\text{Total number of species of all life forms}} \times 100$$

3. Results and Discussion

The forest of the Ramnagar wildlife Sanctuary represents typical subtropical vegetation, dominated mainly by broad leaved trees inter-spread with few scattered patches of shrubs. The forest has been classified as Northern dry-mixed deciduous forest (Type 5B/C2) as described by Champion and Seth [16]. *Acacia modesta* has been observed to be the most prominent tree species. Other dominant tree species include *Grewia optiva*, *Mallotus philipensis*, *Cassia fistula*, *Dalbergia sissoo*, *Flacourtia indica*, *Lannea coromendalica*, *Crataeva adansonii*, *Aegle marmelos*, *Phyllanthus emblica*, *Acacia catechu*, *Leucaena leucocephala* and *Mitragyna parviflora*. *Justicia adhatoda* and *Lantana camara* have found to be the most conspicuous shrub species covering large area of ground in certain places. Other important shrub species include *Carissa opaca*, *Woodfordia fruticosa*, *Murraya koiengii*, *Capparis sepriaria*, *Solanum erianthum*, *Gymnosporia royleana*, *Helicteres isora* and *Abutilon ramosum*. *Fluggea virosa*, *Ricinus communis* and *Ziziphus nummularia* were found abundant along the road passing through the Sanctuary. The herbaceous layer, structurally and numerically most prominent during monsoon season, is represented by *Anagalis arvensis*, *Argemone mexicana*, *Oxalis corniculata*, *Stellaria media*, *Malvastrum coromendellianum*, *Bidens bipinnata*, *Achyranthes aspera*, *Sonchus asper*, *Euphorbia hirta*, *Martynia annua*, *Cleome viscosa*, *Medicago denticulate*, *Chenopodium album*, *Pupalia lappacea*, *Cynodon dactylon*, etc. Climbers, typical of these hills and slopes have been represented by *Tinospora cordifolia*, *Abrus precatorius*, *Cuscuta reflexa*, *Ipomoea carica*, *Diplocyclos palmatus*, *Dioscorea belophylla*, *Macfadyena unguis-cati*, *Merremia aegyptia*, *Luffa acutangula*, *Pergularia extensa*, *Vallisneria spiralis* and *Trichosanthes cucumerina*.

A total of 259 tracheophytes (belonging to 72 families) collected from the study area have been classified into various life-forms as per the classification proposed by Raunkiaer [3]. Number of species and percentage of different life-form have been depicted in Table [1].

Table 1: Total number of species and percentage of different life form classes

Life form classes	Abbreviation	Number	Percentage (%)
Macro-phanerophytes	Mp	60	23.16
Nano-phanerophytes	Np	30	11.58
Chamaephytes	Ch	28	10.81
Hemi-cryptophytes	H	23	8.88
Therophytes	Th	89	34.36
Cryptophytes	Cr	6	2.31

Lianas	L	21	8.11
Epiphytes	E	2	0.77
Hydrophytes	Hh	0	0
Total		259	100

Perusal of the table revealed therophytes (89 spp., 34.36%) to be the most dominating life-form followed by Macrophanerophytes (60 spp., 23.16%), Nanophanerophytes (30 spp., 11.58%), Chamaephytes (28 spp., 10.81%), Hemicryptophytes (23 spp., 8.88), Lianas (21 spp., 8.11%), Cryptophytes (6 spp., 2.31%) and Epiphytes (2 spp., 0.77%). The biological spectrum of the present study showed variation from the normal biological spectrum of Raunkiaer [3] (Fig. 1).




Figure 1: Comparison of Biological spectrum of present area with Normal biological spectrum

From the analysis of the relative proportion of various life-forms present in the study area, the phyto-climate of the study area has been classified as Thero-Phanerophytic. The dominance of therophytes (89 spp., 34.36%) is the characteristic of tropics and often related to soil conditions and climate [17]. Moreover, the predominance of therophytes is also attributed to introduction of annual weeds and biotic influences [18] and indicates a disturbed environmental condition [19]. Biotic influences can be a major factor for increased therophytic percentage owing to the fact that the sanctuary is located in vicinity of Jammu city. Peaking developmental activities and expanding urban sprawl has been a major cause of increased biotic interferences in the area. Since the Sanctuary is intersected by national highway the effect of biotic influences is further magnified.

Higher percentage of Macrophanerophytes (60 spp., 23.16%) reveals the predominance of trees as the area falls in Shivaliks which provides congenial edaphic and climatic conditions for growth of over-story [14]. The climate of the study area is warm in general and dry during summers and warm and moist during rainy season, thus confirming the preponderance of Therophytes (plants of warm and dry climate) and Phanerophytes (plants of warm and moist climate). The comparison of life forms of study area with the

Normal biological spectrum [3] and with adjoining areas like Jammu [14], Trikuta Hills [20], Kathua [21], Mahamaya [22] and Renuka wildlife sanctuary [17] having similar climatic conditions in North western Himalayas is represented in Table [2]. Perusal of the table reveals a similar type of phytoclimate in most of the areas except for Trikuta Hills

where a Thero-chamaephytic phytoclimate has been reported which may be because of the varied amount of disturbances and latitudinal and longitudinal difference. Occurrence of similar biological spectrum in different regions indicates similar climatic conditions [23].

Table 2: Comparison of life forms of study area with different areas of North-west Himalayas

Region	Mp	Np	Ch	H	Th	E	Cr	L	Phytoclimate
NBS	28	15	9	26	13	3	4	-	Phanerophytic
Trikuta Hills	9.44	14.45	22.94	13.87	31.02	0.19	4.62	1.55	Thero-chamaephytic
Kathua district	16.63	12.79	12.36	15.77	32.89	0.63	2.55	5.11	Thero-phanerophytic
Mahamaya	19.5	10.99	13.82	7.44	38.29	0.70	1.06	7.08	Thero-phanerophytic
RenukaWLS	29.4	19.16	7.61	5.51	31.24	0.78	2.10	0.79	Thero-phanerophytic
Jammu	19.49	9.37	11.9	8.35	38.23	5.06	1.78	3.8	Thero-phanerophytic
Present study	23.16	11.58	10.81	8.88	34.36	0.77	2.31	8.11	Thero-phanerophytic

NBS= Normal Biological Spectrum, WLS= Wildlife Sanctuary, Mp= Macrophanerophyte, Np= Nanophanerophyte, Ch= Chamaephytes, Cr=Criptophyte, H=Hemicriptophyte, Th=Therophyte, E= Epiphytes, L=Linanas.

4. Conclusion

Present study revealed the vegetation to be predominantly sub-tropical in nature having a higher percentage of therophytes followed by phanerophytes as compared to normal biological spectrum. On the basis of this study the phytoclimate of the area, as per Raunkiaer's terminology has been described as thero-phanerophytic. This indicates influence of anthropogenic activities in the area which favours the chances of growth of short lived annuals. The study is also significant because information related to life-forms gives an account of prevailing phytoclimatic conditions which can further be used to infer microclimatic conditions in an area. The study further gains significance as the life-forms suggest increased biotic interference in the wildlife sanctuary which in real terms is meant for protection and conservation of natural environment.

References

- [1] H.J. Oosting, The Study of Plant Communities (2d ed.). W.H. Freeman and Company. San Francisco. pp. 440, 1956.
- [2] T.L. Steiger, "Structure of prairie vegetation," Ecology, 11, pp. 170-238, 1930.
- [3] C. Raunkiaer, The life forms of plants and statistical plant geography, Clarendon Press, Oxford. 1934.
- [4] D. Palit, G. Ganguly, A. Mukherjee, "Ecological studies in Ramnabagan Wildlife Sanctuary, Burdwan: I. Biological spectrum," Science and Culture, 68(5-6), pp. 147-149, 2002.
- [5] L. Milne, M. Milne, "The Arena of Life: The Dynamics of Ecology," Doubleday, Natural History Press, Garden City, New York, pp. 240, 1971.
- [6] E. Warming, Ecology of Plant. An introduction to the study of communities, Clarendon Press, Oxford, pp. 422, 1909.
- [7] P. Dansereau, "Description and recording of vegetation upon a structural basis," Ecology, 32, pp. 172-229, 1957.
- [8] E. O. Box, "Predicting physiognomic vegetation types with climate variables," Vegetatio, 45, pp. 127-139, 1981.
- [9] C.R. Reddy, P.H. Krishna, S.L. Meena, R. Bhardwaj, K.C. Sharma, "Composition of life forms and biological spectrum along climatic gradient in Rajasthan, India," International Journal of Environmental Sciences, 1(7), pp. 1632-1639, 2001.
- [10] P.C. Lemon, "Field and Laboratory Guide for Ecology," Burgess, Minn, pp. 154, 1962.
- [11] C.J. Krebs, Ecology: The Experimental Analysis of Distribution and Abundance, (4th eds), Harper Collins, California, pp. 467. 1994.
- [12] O. S. Kotiwar, B.R. Pandit, A.J. Paturkar, "Forms and biological spectrum of the flora of Gir forest, Gujarat," Geobios new Reports, 15, pp. 17-20, 1996.
- [13] V.M. Meher-Homaji, "Life-forms and biological spectra as epharmonic criteria of aridity and humidity in tropics," Journ. India Bot. Soc., 43(3), pp. 424-430, 1964.
- [14] A.K. Raina, N. Sharma, "Floristic Composition, Lifeform Classification and Biological Spectrum of District Jammu, Jammu and Kashmir, North-Western Himalayas," Environment Conservation Journal, 11(3), pp. 49-57, 2010.
- [15] B.M. Sharma, P. Kachroo, Flora of Jammu and Plants of Neighbourhood, Volume I, Bishan Pal Singh Mahindra Pal Singh, Connaught Place, Dehradun, India. 1981.
- [16] H.G. Champion, S.K. Seth, "A Revised Survey of the Forest Types of India," Department of Forest, Published by Govt. of India Press, 1968.
- [17] S.P. Subramani, V. Jishtu, R.K. Verma, K.S. Kapoor, "Floristic Composition, Life-forms and Biological spectrum of Renuka Wildlife Sanctuary, Himachal Pradesh," The Indian Forester, 133(1), pp. 79-91, 2007.
- [18] Z. Sher, Z. D. Khan, "Floristic Composition, Life Form and Leaf Spectra of the Vegetation of Chagharzai Valley, District Buner," Pakistan Journal of Plant Sciences, 13(1), pp. 57-66, 2007.
- [19] M. Al-Yemni, H. Sher, "Biological spectrum with some other ecological attributes of the flora and vegetation of the Asir Mountain of South west, Saudi Arabia," African Journal of Biotechnology, 9(34), pp. 5550-5559. 2010.
- [20] S.K. Kapur, "Phytosociological studies in Trikuta Hills with special reference to the distribution pattern of economic plant products," Ph.D. thesis, University of Jammu, Jammu (J&K), India, 1982.

- [21] M. Jhangir, "Phytodiversity Characterization of Distict Kathua using Remote Sensing and GIS," Ph.D. thesis, University of Jammu, Jammu (J&K), India, 2004.
- [22] J. Sudan, "Phytodiversity and Socio-economic studies of Mahamaya Chatchment, Jammu, J&K," Ph.D. thesis, University of Jammu, Jammu (J&K), India. 2007.
- [23] M. Khan, F. Hussain, S. Musharaf, Imdadullah, "Floristic composition, life form and leaf size spectra of the coal mine area vegetation of darra adam khel, Khyber pakhtonkhwa, Pakistan," Journal of Biodiversity and Environmental Sciences, 1(3), pp. 1-6, 2011.

