

Tissue Damage Induces *in vivo* Production of Asiaticoside in *Centella asiatica* (Linn.) Urban

Kumari Nisha¹, Dr. Rimjhim Sheel², Jainendra Kumar³

¹Research Scholar, Department of Biotechnology, College of Commerce, Patna, India

²Department of Botany, Ganga Devi Mahila Mahavidyalaya, Kankerbagh, Patna, 800 020, India

³Professor, Department of Botany and Biotechnology, College of Commerce, Patna (Bihar), India 800 020


Abstract: The sub-erect plant *Centella asiatica* (Linn.) Urban was collected from two different areas. 7.5gm of leaves were collected from protected area where no trampling and grazing occurs and same amount of leaves from the heavily grazed ruderal area. The production of triterpene, Asiaticoside increases in response to grazing and trampling by hoofed animals. The production of Asiaticoside is also induced by tissue damage (artificially) in leaves and petioles of potted plants.

Keywords: *Centella asiatica*, Asiaticoside, Tissue damage, Grazing, Trampling, Ruderal

1. Introduction

Centella asiatica (Linn.) Urban is a sub-erect plant from family Apiaceae (previously known as Umbelliferae) Known as Mandukparniin Ayurveda in India. Occurring in swampy areas, comprises some 50 species (James and Dubery 2009), inhabiting tropical and sub-tropical regions. Small medicinal herb yielding a number of preparations used for different ailments (Dey, 1980), and the plant leaves have also been used as powder for improving Memory (Kirtikar and Basu 1934). Active compounds asiaticoside and madecassoside (Kartnig, 1988; Farnsworth and Bunyaphatsara 1992) are the ester glycoside derivatives of triterpenic acid. Asiatic acid and madecassic acid have been isolated and purified (Kumar et al. a, 2012). Schaneberg et al. (2003) reported six triterpenes - asiaticoside, madecassoside, asiatic acid, madecassic acid, terminolic acid and asiaticoside – B.

Structure of Asiaticoside



Asiaticoside

Molecular Formula C₄₂H₇₂O₁₃

Molecular Weight 959.12

Pentacyclic triterpenes are synthesized from the C₅ compound isopentenyl pyrophosphate (IPP) in cytoplasm (Isvett et al. 2002). In plants, IPP is produced from Mevalonic acid (Brown 1998), Deoxyxylulose (Rohmer et al. 1993) and Aminoacids (Koops et al. 1991). However, Mevalonate (MVA) is the favoured precursor for IPP. When

pentacyclic triterpenes are synthesized (Akashi et al. 1994), phenotypic plasticity of some degree have been observed (Dutta and Maiti 1968 and Singh 1991) and reported in this plant without any change in chromosome number. Chemical variability is also indicated by variable colour intensity of the separated straw pink layer developed by HCL (Kumar et al., 2012).

2. Material and Methods

Plant materials were collected from protected area where no trampling and grazing occurred and from heavily grazed ruderal area. Total triterpene estimation was achieved by adding 10 drops of acetic anhydride into chloroform extract of leaves and petioles. Two drops of concentrated HCL was mixed to develop three partitioned coloured phases in which middle straw pink phase represented terpenes.

7.5 g of leaves and petioles, air-dried for 15 hours, were powdered in mortar and pestle. MeOH-H₂O gradient protocol was followed to isolate asiaticoside from the crude methanolic extract of the plant leaves and petioles through fractionation (Kumar et al. 2012a). Fraction 3 eluted with 60:40 MeOH-H₂O was used to obtain crystals of Asiaticoside at 40⁰ C in a vacuum oven. Same protocol was also followed in artificially damaged leaves of *Centella asiatica* grown in the pots.

Plate 1



Figure 1: Heavily grazed (Ruderal habitat)

Volume 4 Issue 4, April 2015

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY




Figure 2: Protected Area



Figure 3: Normal Specimen (In pots)



Figure 4: Artificially Tissue

3. Result and Discussion

Investigation indicated that the Asiaticoside (Kumar *et.al.*, 2012) are produced in high amounts in leaves and petioles in heavily grazed ruderal area damage and also in artificially induced tissue damage condition. Table-1 shows the level of Asiaticoside was 1.5 mg/gm where the tissue damaged occurred due to heavy grazing of animals whereas in the protected areas where no trampling or grazing occurred it was 0.65mg/g. Hence, there was a rise of 0.36 mg./gm in the production of Asiaticoside in heavily grazed area.

Table 1

Sample	Conditions	Amount of asiaticoside (mg/gm of leaf & petiole)
1.	Protected habitat where practically no trampling occurs	0.65
2.	heavily grazed area	1.5


The impact of tissue damage conditions on the productivity of Asiaticoside in *Centella asiatica*

Table 2

Sample <i>Centella asiatica</i> (Potted Plant) (mg/gm of dried leaf & petiole).	Normal Specimen	Tissue damaged specimen
	0.75	2.05

Content of the Asiaticoside in plants of *Centella asiatica* which was subjected to artificially induced tissue damage.

Table 3



Histogram shows the production of Terpene, Asiaticoside increases during tissue damage due to heavily grazing and trampling or damaging artificially

The increasing trend in the production of Asiaticoside was also noticed in the potted plants, where the tissues of the leaves were damaged artificially by scrubbing with sand paper. 2.05 mg/gm of the terpene Asiaticoside was produced

by the potted plant in response to the artificial tissue damage. This was 1.3 mg/gm higher than the asiaticoside produced in the potted plant (without tissue damage). Higher production of defensive compound has been reported for several plants such as for *Trifolium repens* (white clover) where HCN is produced in response to herbivory grazers (Hayden and Parker, 2002).

4. Conclusion

It was discovered that the plant produced more useful terpene the Asiaticoside under conditions of tissue damage by herbivory animals in natural habitat. This was also noticed in the potted plants where artificially. induction of tissue damage also causes higher production of these useful compounds.

References

- [1] James, J. T. Meyer, M. & Dubery, I. A. 2008. Characterisation of two phenotypes of *Centella asiatica* in Southern Africa through the composition of four triterpenoids in callus, cell suspension and leaves. *Plant cell tissue organ culture* 94: 91-99.
- [2] James, J.T. & Dubery, I.A. 2009. Pentacyclic triterpenoids from the medicinal herb, *Centella asiatica* (L.) Urban. *Molecules* 14: 3922-3941
- [3] Dey, A.C. 1980. *Indian Medicinal Plant* Bishen Singh Mahendra Pal Singh, Dehradun India.
- [4] Kirtikar, K. R. & Basu, B. D. 1934. (Reprint 1988) *Indian Medicinal Plants*. Vol. 2 Bishen Singh and Mahendra Pal Singh, Dehradun, India.
- [5] Karting, T. 1988. Clinical applications of *Centella asiatica* (L) Urb. In *herbs, Spices and Medicinal Plants*; Cracker, L.E., Simon, J.E., Eds.; Oxyx Press: Phonix, AZ, USA, 1988; pp. 145-173.
- [6] Farnsworth, N.R and Bunyapraphatsara N.(eds.) 1992. *Thai medicinal plants*. Bangkok Prachachon.
- [7] Hayden, K.J and Parker, I.M. 2002. Plasticity in cynogenesis of *Trifolium repens*: inducibility fitness costs and variable expression. *Evolutionary Ecology Research*, 4:155 – 168.
- [8] Kumar, J., Kumar, M., Sheel, R. and Nisha, K. 2012 a. *Centella asiatica* (Linn.) Urb. <http://www.harmonybihar.org/Centella.pdf>. Retrieved on May 15 2013.
- [9] Kumar J, Nisha K, Kumar M and Sheel R. 2012. (a) Isolation of biologically active triterpene ester glycosides (asiaticoside and madecassoside) from *Centella asiatica* (Linn.) Urban. *Biospectra* 7 (3) : 115 – 120.
- [10] Schaneberg, B.T., Milkil, J.R., Bedir, E. & Khan, L.A. 2003. An improved HPLC method for quantitative determination of six triterpenes in *Centella asiatica* extracts and commercial products. *Pharmazie*. 58(6): 381 – 4.
- [11] Isvett, J.F.S., Jaime, O.L., Maria, D.E.L., Carmen, M. H. & Ana, C. R. V. 2002. Biosynthesis of sterols and triterpenes in cell suspension cultures of *Uncaria tomentosa*. *Plant cell Physiol* 43 (12): 1502 – 1509.
- [12] Brown, G. D. 1998. The biosynthesis of steroids and triterpenoids. *Nat Prod Rep* 15: 653 – 696.
- [13] Kooops, A.J., Italiaander, E. & Groeneveld, H. W. 1991. Triterpenoids biosynthesis in the seedling of *Euphorbia lathyris* L. From sucrose and amino acids. *Plant. Sci.* 74: 193 – 201.
- [14] Akashi, T., Furuno, T., Takahashi, T. & Ayabe, S. 1994. Biosynthesis of triterpenoids in cultured cells and regenerated and wild plant organs of *Taraxacum officinale*. *Phytochemistry* 36: 303 – 308.
- [15] Dutta, P. C. & Maithi, R.K. 1986. Chromosomal biotypes of *Centella asiatica* (L.) Urban. and *hydrocotyl javanica* Thunb. *The nucleus* Vol I (2): 111 – 117.