

i Sëmundjeve Infektive. pp.41–53.53. Tirana: Mediaprint.

[18] A Sejdini; R Mahmud; Y A L Lim; M Mahdy; F Sejdini; V Gjoni; K Xhaferraj; G Kasmi (2011). Intestinal parasitic infections among children in central Albania. *Ann Trop Med Parasitol*. Apr; 105(3): 241–250.

[19] Bashkim Nezaj; Majlinda Kota; Gjeorgjina Kuli-Lito (2014). *International Journal of Science and Research (IJSR) Volume 3 Issue 11*.

[20] Peruzzi S; Gorrini C; Piccolo G; Calderaro A; Dettori G; Chezzi C. (2006). Prevalence of intestinal parasites in the area of Parma during the year 2005. *Acta Biomed*. Dec; 77(3):147-51.

[21] Tamer GS; Caliskan S; Willke A. (2008). Distribution of intestinal parasites among patients who presented at the parasitology laboratory of the Kocaeli University School of Medicine Hospital. *Turkiye Parazitol. Der*: 32(2):126-9.

[22] Traoré SG, Odermatt P, Bonfoh B, Utzinger J, Aka ND, Adoubryn KD, Assoumou A, Dreyfuss G, Koussémon M. (2011). No *Paragonimus* in high-risk groups in Côte d'Ivoire, but considerable prevalence of helminths and intestinal protozoon infections. *Parasit Vectors* 4:96. [PubMed Abstract](#) | [BioMed Central Full Text](#) | [PubMed Central Full Text](#).

[23] Cifuentes E, Suarez L, Espinosa M, Juarez-Figueroa L, Martinez-Palomo A. (2004). Risk of *Giardia intestinalis* infection in children from an artificially recharged groundwater area in Mexico City. *Am J Trop Med Hyg*, 71:65-70. [PubMed Abstract](#) | [Publisher Full Text](#).

[24] Ish- Horowicz M; Korman S. H; Shapiro M. (1989): Asymptomatic giardiasis in children. *Pediatr. Infect. Dis. J.* 8: 773 9.

[25] Hjelt K., Poerregaard A., Krasilnikoff P. A. (1992). Giardiasis causing diarrhea in suburban Copenhagen; Incidence, physical growth, clinical symptoms and small intestinal abnormality. *nActa Paediatr*; 81: 881 6.

[26] Stephenson L. S., Latham M. C. and Ottesen E. A. (2000). Malnutrition and parasitic helminth infections, *Parasitology* 121, pp. S23–38. Full Text via CrossRef.

[27] Tarleton J. L., Haque R., Mondal D., Shu J., Farr B. M. and Petri W. A. (2006). Cognitive effects of diarrhea, malnutrition, and *Entamoeba histolytica* infection on school age children in Dhaka, Bangladesh, *Am J Trop Med Hyg* 74, pp. 475–481. View Record in Scopus | Cited By in Scopus (8).

[28] Negrao-Correa D. and Teixeira M. M. (2006). The mutual influence of nematode infection and allergy, *Chem Immunol Allerg* 90, pp. 14–28. View Record in Scopus | Cited By in Scopus (6).

[29] El-Nofely A. and Shaalan A. (1999). Effect of *Ascaris* infection on the nutritional status and IQ of children, *Int J Anthropol* 14, pp. 55–59. Full Text via CrossRef | View Record in Scopus | Cited By in Scopus (0).

[30] Berrilli F, di Cave D, d'Orazi C, Orecchia P, Xhelilaj L, Bejko D, Çaça P, Bebeci D, Cenko F, Donia D, Divizia M. (2006). Prevalence and genotyping of human isolates of *Giardia duodenalis* from Albania. *Parasitology International* 55:295–297. [PubMed](#).

Table 1: Distribution of Positivity infection among children in Baltëz according to age groups and sex

Characters	Infected with parasites	Positivity No. (%)
Age groups (years): P>0.05		
2-3	14	10 (71.4%)
4-6	47	27 (57.4%)
7-10	65	42 (64.4%)
11-15	33	30 (91%)
Gender: P>0.05 $\chi^2=2.39$		
Male (-)	20	25.6%
Male (+)	58	74.4%
Female (-)	30	37%
Female (+)	51	63%

(-) negative cases
 (+) positive cases

Table 2: The positivity of parasites according to the data of standardized questioner

Data	Negative	Positive	Total number	Statistics data
Ethnicity				P>0.05 $\chi^2=0.33$
Kosovo	13	28	41	
Roman	13	33	46	
Albanian	24	48	72	
Weight loss				P>0.05 $\chi^2=1.175$
Children without loss of wight	5	18	23	
Children with loss of wight	45	91	136	
Anemia				P<0.001 $\chi^2=13.44$
Without Anemia	33	38	71	
With Anemia	17	71	88	
Mother education status :				P<0.03 $\chi^2=6.6$
Illiterate or incomplete primary	11	47	58	
Complete primary or secondary	35	56	91	
High school	4	6	10	
University	0	0	0	

Table 3: The kind of pathogens and their percentage

<i>Pathogens</i>	<i>Frequency</i>	<i>Percent</i>
Ascaris lumbricoides	1	0.6
Blasocystis hominis	18	11.3
Chilomastic masnili	1	0.6
Endolimax nana	2	1.3
E. nana and B. hominis	1	0.6
Entamoeba spp	6	3.9
Entamoeba spp and B.homis	17	10.6
Entamoeba spp and B.homis and E. nana	3	2
Entamoeba spp and E.nana	1	0.6
Enterobius vermicularis and Entamoeba spp and E. nana and B. hominis	1	0.6
Entrobius vermicularis and Entamoeba spp	1	0.6
Giardia lamblia	20	12.6
G. lamblia and A. lumbricoides	1	0.6
G. lamblia and B.hominis	5	3.1
G. lamblia and Entamoeba spp	4	2.5
G. lamblia and Entamoeba spp and B. hominis	1	0.6
G. lamblia and Trichuris trichiura	3	2
Hymenolepis nana	5	3.2
H. nana and Entamoeba spp	1	0.6
H. nana and Entamoeba spp and B. hominis	1	0.6
H. nana and G. lamblia and Trichuris trichiura	1	0.6
H. nana and Strongyloides stercoralis	1	0.6
H. nana and T. trichiura and B. hominis	2	1.3
S. stercoralis	1	0.6
T. trichiura	6	3.9
T. trichiura and Entamoeba spp and B. hominis	4	2.5
T. trichiura and G. lamblia and Entamoeba spp and B. hominis	1	0.6
Negativ	50	31.4

