









- origin", Annals of Entomological Society of America 75(1), pp. 51-55. 1996.
- [2] H. Baumann, K. J. Wilson, P. S. Chen, R. E. Humbel, "The amino acid sequence of a peptide (PS-1) from *Drosophila funebris*: a paragonial peptide from males which reduces the receptivity of the females", European Journal of Biochemistry, 52, pp. 521-529. 1975.
- [3] H. P. J. Bennett, "Use of ion-exchange Sep-Pac cartridges in the batch fractionation of pituitary peptides", Journal of Chromatography A, 359, pp. 383-390. 1986.
- [4] V. P. Bindokas, M. E. Adams, "Hemolymph composition of the tobacco budworm, *Heliothis virescens* F. (Lepidoptera: Noctuidae)" Comparative Biochemistry and Physiology, 90, pp. 151-155. 1988.
- [5] R. T. Carde, A. K. Minks, "Control of moth pests by mating disruption: success and constraints", Annual Review of Entomology, 40, pp. 559-585. 1995.
- [6] R. T. Carde, K. Trammel, W. L. Roelofs, "Disruption of sex attraction of the redbanded leafroller *Argyrotaenia velutinana* with microencapsulated pheromone components", Environmental Entomology, 4, pp. 448-552. 1975.
- [7] T. Chapman, "Seminal fluid-mediated fitness traits in *Drosophila*", Heredity, 87, pp. 511-521. 2001.
- [8] T. Chapman, L. A. Herndon, Y. Heifetz, L. Parteidge, M. F. Wolfner, "The Acp26Aa seminal fluid protein is a modulator of early egg-hatchability in *Drosophila melanogaster*". Proc Biol Sci 268:1647-1654. 2001
- [9] P. S. Chen, "The accessory gland proteins in male *Drosophila*: structural, reproductive, and evolutionary aspects", Experientia, 52, pp. 503-510. 1996.
- [10] G. B. Craig Jr, "Mosquitoes: female monogamy induced by male accessory gland substance", Science, 156, pp. 1499-1501. 1967.
- [11] S. J. Davies, T. Chapman, "Identification of genes expressed in the accessory glands of male Mediterranean fruit flies (*Ceratitidis capitata*)", Insect Biochemistry and Molecular Biology, 36, pp. 846-856. 2006.
- [12] J. P. Deland, G. J. R. Judd, B. D. Roitberg, "Disruption of pheromone communication in three sympatric leafroller (Lepidoptera: Tortricidae) pests of apple in British Columbia", Environ Entomology, 23, pp. 1084-1090. 1996.
- [13] Y. G. Divakar, N. Medhini, D. Manjulakumari, "Mass rearing of *Spodoptera litura* (fabricus.) under laboratory conditions on artificial diet and studies on their mating and egg laying behavior", in Insect Pest Management, A Current Scenario, Dunston P Ambrose (ed), India, 2011.
- [14] L. A. Herndon, M. F. Wolfner, "A *Drosophila* seminal fluid protein, Acp26Aa, stimulates egg laying in females for 1 day after mating", Proceedings of National Academy of Science, USA, 92, pp. 10114-10118. 1995.
- [15] Isidra Recio, Servaas Visser, "Two-ion exchange chromatographic methods for the isolation of antibacterial peptides from lactoferrin in-situ enzymatic hydrolysis on an ion-exchange membrane, J Chromatography A, 831, pp. 191-201. 1999.
- [16] H. Izadi, B. Subrahmanyam, "Protein profiles of *Spodoptera litura* (F). male accessory reproductive gland and its secretions", Communications in Agricultural and Applied Biological Sciences, 70(4), pp. 857-862. 2005.
- [17] Z. Y. Jin, H. Gong, "Male accessory gland derived factors can stimulate oogenesis and enhance oviposition in *Helicoverpa armigera* (Lepidoptera: Noctuidae)", Archives of Insect Biochemistry and Physiology, 46, pp. 175-185. 2001.
- [18] Jin-Feng Yu, Cong Li, Jin Xu, Jian-Hong Liu, Hui Ye, "Male accessory gland secretions modulate female post-mating behaviour in the moth *Spodoptera litura*", Journal of Insect Behaviour, 27, pp. 105-116. 2014.
- [19] T. G. Kingan, W. M. Bondar, A. K. Raina, J. Shabanowitz, D. F. Hunt, "The loss of female sex pheromone after mating in the corn earworm moth *Helicoverpa zea*: identification of a male pheromonostatic peptide", Proceedings of National Academy of Sciences, USA, 92, pp. 5082-5086. 1995.
- [20] T. G. Kingan, P. A. Thomas-Laemont, A. K. Raina, "Male accessory gland factors elicit change from 'virgin' to 'mated' behaviour in the female corn earworm moth *Helicoverpa zea*", Journal of Experimental Biology, 183, pp. 61-76, 1993.
- [21] E. Kubli, "Sex-peptides: seminal peptides of the *Drosophila* male", Cellular and Molecular Life Sciences, 60, pp. 1689-1704. 2003.
- [22] E. L. Larson, J. A. Andres, R. G. Harrison, "Influence of the male ejaculate on postmating prezygotic barriers in field crickets", PLoS One, 7(10):e46202, 2012.
- [23] R. A. Leopold, "The role of male accessory glands in insect reproduction", Annual Review of Entomology, 21, pp. 199-221. 1976.
- [24] D. Manjulakumari, Geetha Bali, "Mating: A mechanism that regulates oviposition behaviour in the silk moth *Bombyx mori* L.", in Readings in Behavior, Ramamurthy and Geetha Bali (eds.), New Age International Ltd., India, 1996.
- [25] C. Naccarati, N. Audsley J. N. Keen, J. H. Kim, G. J. Howell, Y. J. Kim, R. E. Isaac, "The host-seeking inhibitory peptide, Aea-HP-1, is made in the male accessory gland transferred to the female during copulation", Peptides, 34, pp. 150-157. 2012.
- [26] Y. Y. Ohashi, K. Haino-Fukushima, Y. Fuyama, "Purification and characterization of an ovulation stimulating substance from the male accessory glands of *Drosophila suzukii*", Insect Biochemistry, 21, pp. 413-419. 1991.
- [27] L. Paemen, A. Tips, L. Schoofs, P. Proost, J. Van Damme, A. De Loof, "Lom-AG-myotropin: a novel myotropic peptide from the male accessory glands of *Locusta migratoria*", Peptides, 12, pp. 7-10. 1991.
- [28] H. Qin, Z. Ye, S. Huang, J. Ding, R. Luo, "The correlations of the different host plants with preference level, life duration and survival rate of *Spodoptera litura* Fabricius", Chinese Journal of Eco-Agriculture, 12, pp. 40-42. 2004.
- [29] J. G. Riemann, D. J. Moen, B. J. Thorson, "Female monogamy and its control in houseflies", Journal of Insect Physiology, 13, pp. 407-418, 1967.

- [30] J. Rooney, S. M. Lewis, "Differential allocation of male-derived nutrients in two lampyrid beetles with contrasting life-history characteristics", *Behavioral Ecology*, 10, pp. 97-104. 1999.
- [31] T. Schmidt, Y. Choffat, S. Klauser, E. Kubli, "The *Drosophila melanogaster* sex-peptide: A molecular analysis of structure-function relationship", *Journal of Insect Physiology*, 39, pp. 361-368. 1993.
- [32] A. Shobha, "Characterization of peptides isolated from male accessory glands of *Helicoverpa armigera*", Ph D thesis, Bangalore University, India, 2007.
- [33] A. Shobha, Geetha Bali, D. Manjulakumari, "Purification and partial characterization of a male derived peptide that enhances egg maturation and oviposition in female cotton bollworm, *Helicoverpa armigera* (Lepidoptera: Noctuidae)", *Entomon*, 34(4), pp. 213-221, 2009.
- [34] H. M. Smid, A. B. Koopmanschap, C. A. D. De Kort, H. Schooneveld, "A peptide from the male accessory gland in *Leptinotarsa decemlineata*: purification, characterization and molecular cloning", *Journal of Insect Physiology*, 43, pp. 355-362. 1997.
- [35] A. Suzuki, H. Nagasawa, H. Kataoka, Y. Hori, A. Isogai, S. Tamura, F. Guo, X. Zhong, H. Ishizaki, M. Fujishita, A. Mizoguchi, "Isolation and characterization of prothoracicotropic hormone from silkworm, *Bombyx mori*". *Agricultural and Biological Chemistry*, 46, pp. 1107-1109. 1982.
- [36] J. A. Veenstra, F. G. Noriega, R. Graf, R. Feyereisen, "Identification of three allatostatins and their cDNA from the mosquito *Aedes aegypti*", *Peptides*, 18, pp. 937-942, 1997.
- [37] M. F. Wolfner, "Tokens of love: functions and regulation of *Drosophila* male accessory gland products", *Insect Biochemistry and Molecular Biology*, 27, pp. 179-192. 1997.
- [38] J. Xu, Q. Wang, "Seminal fluid reduces female longevity and stimulates egg production and sperm trigger oviposition in a moth", *Journal of Insect Physiology*, 57, pp. 385-390, 2011.
- [39] S. X. Yi, C. Gillott, "Purification and characterization of an oviposition-stimulating protein of the long hyaline tubules in the male migratory grasshopper, *Melanoplus sanguinipes*", *Journal of Insect Physiology*, 45, pp. 143-150, 1999.