



excised and washed with ethanol (90%) for 1 min then stirred in 20% bleach (5.25% NaOCl) for 30 min. This sterilized explant was used for induction of shoot and then shoot multiplication according to Haq and Dahot (2007a, 2007b, 2007c) and Haq *et al.* (2011, 2014). Briefly, as shoot tips (3 - 4 mm) were divided into two parts and cultured on MS<sub>1</sub> [MS<sub>0</sub> (Murashige and Skoog, 1962) basal salts with B5 vitamins (Gamborget *et al.*, 1968), 10 mg L<sup>-1</sup> BA, 15 mg L<sup>-1</sup> IAA, 30.0 mg L<sup>-1</sup> L-cystein, 3% sucrose and 3.60g L<sup>-1</sup> phytigel] medium for 15-20-days for organogenesis. Shoot induction was carried on MS<sub>1a</sub> (MS<sub>0</sub>, 8mg L<sup>-1</sup> BA and 20 mg L<sup>-1</sup> L-cysteine, 1.0 g L<sup>-1</sup> phytigel), while shoot micropropagation on MS<sub>2</sub> (MS<sub>0</sub>, 8 mg L<sup>-1</sup> BA and 20 mg L<sup>-1</sup> L-cysteine, 2.0 g L<sup>-1</sup> phytigel). Shoot multiplication culture were refreshed after every 3<sup>rd</sup> week, while 3<sup>rd</sup> subculture was subjected for PEG stress (represented as MS<sub>2b</sub>) along with BBTV infected tissue (represented as MS<sub>2a</sub>).

## 2.2 Establishment of PEG-stressed along BBTV infected cultures

On 3<sup>rd</sup> week of 2<sup>nd</sup> subculture, rapidly micro-propagating plantlets from healthy (MS<sub>2</sub>) and BBTV infected (MS<sub>2a</sub>) cultures were refreshed on same culture including one with PEG stress on healthy shoot multiplication [MS<sub>2b</sub> (MS<sub>2</sub> 5% PEG-4000)] medium. These cultures were labelled as MS<sub>2</sub> (control), MS<sub>2b</sub> (BBTV-infected plantlets) and MS<sub>2c</sub> (MS<sub>2</sub> + 5% PEG) and incubated for 6-weeks under 18/6 h lightening with ~2000 lux intensity at 25 ± 1°C. The pH of each culture was adjusted 5.7 to 5.8 and autoclaved at 121°C with 15 lbs for 10 min.

## 2.3 Measurement of Morphological Parameters

At the end of 6<sup>th</sup> week of culture, micro-propagated plantlets taken out from culture glass jars and washed with running tap-water to removed entangled medium. Plantlets were dried on filter-paper and number of plantlets per explant were counted. Shoot height and fresh shoot biomass was taken. Dry shoot biomass was measured by drying shoots at 72°C with electric oven for 72-hrs. Relative water contents (RWC) were expressed [Catsky, 1974; Turner, 1981] by using this formula -  $RWC (\%) = [(FM - DM)/TM] * 100$ .

## 2.4 Measurement of Biochemical Parameters

Fresh shoots were subjected for biochemical analysis like as chlorophyll or photosynthetic pigments were determined by using methods of Arnon (1949) and Lichtenthaler (1987). Glycinebetaine (Grieve and Gratter, 1983) and proline (Bates *et al.*, 1973) contents were determined spectrophotometrically. Total phenol contents were measured as by Ozyigit *et al.* (2007). Dry shoot material was also analyzed for various biochemical contents like as total protein contents (Bradford, 1976), total carbohydrates (Dubois *et al.*, 1956; Cihai and Brun, 1978), reducing sugars (Miller, 1959) and nitrate contents (Morris and Riley, 1963) were analyzed.

## 2.5 Statistical Data Analysis

Data significance of treatment [each treatment with 4-replicates were arranged for one way analysis of variance (ANOVA)] was calculated by using COSTAT computer


statistical software (CoHort software, Berkeley, USA) and DMR test at 5% level of treatment difference.

## 3. Results and Discussion

Banana is a delicate herbaceous plant with false stem and grow above the ground. Stem is comprised on leaf-sheaths full with water. Delay in irrigation, it shows wilting character that is very similar to symptoms developed by plants growing under abiotic stresses like as salinity, cold, high temperature as well as same to biotic stressed symptoms by initial *Banana bunchy top disease* (BBTD) infection (Haq *et al.*, 2014). Both stresses cause growth inhibition and have obviously playing their harmful effects on vegetative growth and yield of banana crop. Diverse soil texture, non-availability of water and pathogen infection have negative impacts on growth of banana crop plants. Such growth retarding factors can be studied in the laboratory for identification of their precise impacts on morpho-physiological aspects of multiplying plantlets. In this experiment, young suckers of BBTV-infected banana plants and healthy banana were cultured under *in-vitro* conditions for multiplication. When micro-propagation cultures ready for 3<sup>rd</sup> sub-culture stage. At this stage, plantlets were also sub-cultured on micro-propagation medium supplemented with 5% PEG-4000 in comparison to healthy and BBTV-infected multiplying plantlets.

In nature, soil exposes its versatile characteristics to the growing plants. Some of them beneficial for plant growth while others especially balance supply of water to growing plants is not always same. When water level down than soil nutrients also have toxic impacts on plant growth both cases water deficit and soil salinity causes growth inhibition. *In-vitro* plant micro-propagation is a correct way for supply of balanced nutrition for optimal plant biomass production. This system is also suitable for study of diverse impacts of different growth retarding factors at cell level. So 3-typed plant micropropagation cultures of banana (*Musa* spp.) cv., Basrai were established like as MS<sub>2</sub> (healthy banana plantlets), MS<sub>2b</sub> (BBTV-infected plantlets) and MS<sub>2c</sub> (healthy plantlets under 5% PEG stress) for 6-weeks. Both PEG stressed and BBTV infected cultures showed a number of variations in their bio-metrics in comparison to healthy plantlets (MS<sub>2</sub>) as shown in table 1.

Plant multiplication rate was observed maximum in MS<sub>2</sub> medium as already suggested by Haq and Dahot (2007b, c), over other two cultures like as MS<sub>2a</sub> and MS<sub>2b</sub>. Culture effect on multiplication rate was non-significant, while order of decrease in number of plantlets per explant was MS<sub>2</sub> > MS<sub>2a</sub> > MS<sub>2b</sub>. The PEG stress have reduced plant multiplication rate greater than BBTV infection, while abnormal plant growth was observed among the multiplying banana plantlets in BBTV infected cultures (Fig 1). Fresh weight and dry weight of plantlets were decreased significantly (p < 0.05). Plant growth rate or fresh biomass has significant relationship with green photosynthetic pigments that also decreased significantly (Haq *et al.*, 2008). Chlorophyll b was decreased in BBTV infected cultures like to chlorophyll a, while it slightly increases in PEG stressed plantlets. Overall decrease in total chlorophyll contents was significant.



**Figure 1:** Comparative morphological appearance of PEG stressed and BBTV-infected plantlets under aseptic micropropagation of Banana (*Musa* spp.) cv., Basrai. a: Control MS<sub>2</sub> culture; b: MS<sub>26</sub> cultures with BBTV infected plantlets; c: MS<sub>2c</sub> -5% PEG-4000 stressed plantlets.

The trend of coloured pigments like as total carotenoids was opposite to green or photosynthetic pigments, they were increased significantly in both PEG stressed as well as in BBTV infected plantlets (Table 1). It can be observed from figure 1, there variation in morphology of micro-propagating culture after 6-weeks. Generally, abiotic-stresses cause chloroplast-injury lead to decrease in green pigmentation. Less pigmentation, low photosynthesis results to decrease in plant vegetative growth. Water deficit stress increases the concentrations of certain free amino acids in the cells for resist against plastid's injury like as proline and glycinebetaine. In actual, both proline and photosynthetic pigments are synthesized in mesophyll cells from same precursor. Under stresses, when breakdown of photosynthetic pigments starts, well there initiation of proline synthesis also starts that could be a best indication in plants are being under stress either biotic or abiotic (Aspinal and Paleg, 1981).

Under environmental stresses, plants express certain molecular bio-markers as indicators. Like as free amino acids and carbohydrates also. Reducing sugars also increased significantly under both PEG as well as BBTV infected banana plants (Table 1). Total proteins and total carbohydrates were decreased ( $p < 0.05$ ) in stressed plantlets. The phenolic components were also increased ( $p < 0.05$ ). Nitrates were decreased among the stressed cultures significantly. Decrease in important plant nutrient substituents under stressed conditions causes inhibition of growth, while increase in certain components in the cell are responsible for prevention of cell injury by stressing factors. These accumulated bio-components can be a source of carbon, hydrogen, nitrogen and energy for the recovery of inhibited physiological processes upon relax of stresses (Blum and Ebercon, 1976; Slamaet *et al.*, 2007). All external stresses are limiting the plant growth efficiency significantly (Hartmann *et al.*, 2005). Especially, abiotic stresses (salinity and drought) increases osmotic potential within the cell (Djibrilet *et al.*, 2005; Vanden and Zeng, 2006). Biotic stresses could also be playing same when infecting the plant tissue externally, like as fungus,

**Table 1:** Comparative morpho-bio-chemical attributes of PEG stress to BBTV infected plantlets of banana (*Musa* spp.) cv., Basrai micro-propagated for 6-weeks at 3<sup>rd</sup> sub-culture stage.

nematodes or carnivores, while viruses also have significant role in plant growth inhibition either by malfunctioning of metabolic biosynthesis or peliotropism in gene expression level of the plant cell.. In this experiment, almost all the parameters were considered to be affected significantly in the micro-propagating plantlets of banana (*Musa* spp.) cultivar Basrai. Both photosynthetic pigments and organic contents developed differential characteristics in the stressed cultures in comparison with the control.

#### 4. Conclusions

Edible banana is a fruit but being one among the important stable food crops, while farmers are replacing banana with other food crops because of its yield production is decreasing to 100% by BBTV infection and now deficiency of irrigation water in the rivers. Both stresses decrease its plant multiplication rate and yields because of increase in osmotic potential of the cell. Meanwhile, lot of free amino acids especially proline and glycinebetaine including other specific groups of reducing sugars and total carotenoids are increased as being stress related bio-markers. Their biosynthesis is activated by osmotic signals developed under external as well as internal stresses. These organic contents could develop specific tolerance against biotic and abiotic stresses but under viral stress is still question mark. Future layout for development resistance against specific factors depends on the expression of specific markers but virus infection based markers are not completing a task to handle related phenomena.

#### 5. Acknowledgement

Author is heartily very thankful to lab assistants as well as students for particular assistance during this work and University of Sindh also for financing this experiment.

#s.	Characteristics	MS <sub>2</sub> - control cultures	MS <sub>2a</sub> - 5% PEG cultures	MS <sub>2b</sub> - BBTV infected cultures	Significance @ 5%
<b>A. Morphological attributes</b>					
a.	# of shoots explant <sup>-1</sup>	<sup>a</sup> 5.75 ± 1.555	<sup>ab</sup> 4.50 ± 0.289	<sup>b</sup> 3.25 ± 0.408	ns
b.	Shoot F.Wt (g)	<sup>a</sup> 1.981 ± 0.002	<sup>b</sup> 1.635 ± 0.001	<sup>c</sup> 1.334 ± 0.001	***
c.	Shoot D.Wt (g)	<sup>a</sup> 0.581 ± 0.004	<sup>b</sup> 0.451 ± 0.005	<sup>c</sup> 0.374 ± 0.005	***
<b>B. Chlorophyll contents (mg g<sup>-1</sup>)</b>					
a.	Chlorophyll a	<sup>a</sup> 0.127 ± 0.117	<sup>b</sup> 0.067 ± 0.025	<sup>c</sup> 0.038 ± 0.011	***
b.	Chlorophyll b	<sup>b</sup> 0.114 ± 0.005	<sup>c</sup> 0.070 ± 0.013	<sup>a</sup> 0.123 ± 0.009	***
c.	Chlorophyll ab	<sup>a</sup> 0.240 ± 0.004	<sup>c</sup> 0.136 ± 0.005	<sup>b</sup> 0.161 ± 0.004	***
d.	Total carotenoids	<sup>c</sup> 1.783 ± 0.002	<sup>b</sup> 1.952 ± 0.006	<sup>a</sup> 2.144 ± 0.002	***
<b>C. Organic contents</b>					
a.	Total carbohydrates (mg g <sup>-1</sup> )	<sup>a</sup> 1.104 ± 0.002	<sup>ab</sup> 1.085 ± 0.001	<sup>c</sup> 1.070 ± 0.001	ns
b.	Reducing sugars (mg g <sup>-1</sup> )	<sup>c</sup> 0.547 ± 0.007	<sup>b</sup> 0.645 ± 0.007	<sup>a</sup> 0.701 ± 0.010	***
c.	Total protein contents (mg g <sup>-1</sup> )	<sup>a</sup> 1.491 ± 0.004	<sup>b</sup> 1.301 ± 0.003	<sup>b</sup> 1.273 ± 0.003	***
d.	Proline contents (mg g <sup>-1</sup> )	<sup>a</sup> 0.844 ± 0.004	<sup>b</sup> 0.867 ± 0.003	<sup>c</sup> 0.944 ± 0.003	***
e.	Glycinebetaine contents (mg g <sup>-1</sup> )	<sup>c</sup> 0.640 ± 0.002	<sup>b</sup> 0.776 ± 0.001	<sup>a</sup> 1.053 ± 0.001	***
f.	Phenolics (mol m <sup>-3</sup> g <sup>-1</sup> )	<sup>c</sup> 0.271 ± 0.004	<sup>b</sup> 0.364 ± 0.003	<sup>a</sup> 0.450 ± 0.003	***
<b>D. in-organic contents</b>					
a.	RWC (%)	<sup>a</sup> 222.25 ± 0.002	<sup>b</sup> 180.98 ± 0.003	<sup>c</sup> 142.74 ± 0.002	***
b.	Nitrates -NO <sub>3</sub> <sup>-1</sup> (μmol g <sup>-1</sup> )	<sup>a</sup> 0.162 ± 0.007	<sup>b</sup> 0.099 ± 0.008	<sup>b</sup> 0.102 ± 0.010	***

## References

- [1] Arnon, D.I., 1949, Copper enzymes in isolated chloroplasts. Polyphenoloxidase in *Beta vulgaris*. *Plant Physiol.*24: 1-15.
- [2] Aspinall, D. and Paleg, L.G., 1981. Proline accumulation. In: *Physiology and Biochemistry of Drought Resistance in Plants (Physiology Aspects)*, Academic Press, New York, pp. 205-240.
- [3] Bates, L.S., Waldren, R.P. and Tear, I.D., 1973. Rapid determination of free proline for water stress studies. *Plant Soil*39: 205-207.
- [4] Blum, A. and Ebercon, F., 1976. Genotype responses in sorghum to drought stress. III. Free proline accumulation and drought resistance. *Crop Sci.* 16: 379-386.
- [5] Bradford, M.N., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 72: 248-254.
- [6] Catsky, J., 1974, Direct methods of water content determination. In: Slavik, B., (Ed.) *Methods of studying plant water relations*, Springer-Verlag, Berlin, pp.121-123.
- [7] Chaum, S., Siringam, K., Juntawong, N. and Kirdmanee, C., 2010. Water relations, pigment stabilization, photosynthetic abilities and growth improvement in salt stressed rice plants treated with exogenous potassium nitrate application. *Intl. J. Plant Prod.* 4: 187-197.
- [8] Ciha, A.J. and Brun, W.A., 1978. Effect of pod removal on nonstructural carbohydrate concentration in soybean tissue. *Crop Sci.* 18: 773-776.
- [9] Djibril, S., Mohamed, O.K., Diaga, D., Diégane, D., Abaye, B.F., Maurice, S. and Alain, B., 2005. Growth and development of date palm (*Phoenix dactylifera*L.) seedlings under drought and salinity stresses. *Afr. J. Biotechnol.* 4: 968-972.
- [10] Dubois, M., Gilles, K.A., Hamilton, J.K., Rebers, P.A. and Smith, F., 1956. Colorimetric method for determination of sugars and related substances. *Anal. Chem.* 28: 350-356.
- [11] FAO, 2005. Global network on integrated soil management for sustainable use of salt affected Soils. Rome, Italy: FAO Land and Plant Nutrition Management Service.
- [12] Frison, E.A., Orjeda, G. and Sharrock, S.L., 1997. PROMUSA: A global programme for *Musa* improvement. In: *Proceedings of a meeting held in Gosier Guadeloupe*, (Eds.) Frison EA, Orjeda G, Sharrock SL, pp. 8-11; International Network for the Improvement of Banana and Plantain, Montpellier, France, The World Bank, Washington, USA.
- [13] Gamborg, O.L., Miller, R.A. and Ojima, K., 1968. Nutrient requirements of suspension cultures of soybean root cells. *Exp. Cell Res.* 50: 151-158.
- [14] Grieve, C.M. and Grattan, S.R., 1983. Rapid assay for determination of watersoluble quaternary-amino compounds. *Plant Soil*70: 303-307.
- [15] Haq, I.U. and Dahot, M.U., 2007a. Effect of permanent and temporary immersion systems on banana micropropagation. *Pak. J. Bot.* 39: 1763-1772.
- [16] Haq, I.U. and Dahot, M.U., 2007b. Morphophysiological aspects of micropropagating banana under different hormonal conditions. *Asian. J. Plant Sci.* 6: 496-501.
- [17] Haq, I.U. and Dahot, M.U., 2007c. Effect of immersion systems on chlorophyll contents in micropropagating banana. *Afr. J. Biotechnol.*6: 1095-1101.
- [18] Haq, I.U., Dahot, M.U., Khan, S. and Kousar, N., 2009. Screening of banana bunchy top diseased plants: A way to control its spreading. *Plant Omics J.* 2(4): 175-180.
- [19] Haq, I.U., Faheeda, S., Dahot, M.U., Shahrukh, Um-e-aiman, 2008. *In vitro* multiplication of banana (*Musa* spp.) under different NaCl stresses. *Pak. J. Biotechnol.* 4: 25-30.
- [20] Haq, I.U., Shahrukh, Um-E-Aiman, Khan, S., Parveen, N., Fatima, M. and Dahot, M.U., 2014. Certain growth related attributes of bunchy top virus infected banana under ex-vitro conditions. *Afr. J. Biotechnol.* 13(18): 1876-1882.
- [21] Haq, I.U., Soomro, F., Parveen, N., Dahot, M.U. and Mirbahar, A.A., 2011. Certain growth related attributes

of micropropagated banana under different salinity levels. *Pak. J. Bot.* **43(3)**: 1655-1658.

- [22] Hartmann, T., Colledge, M. and Lumsden, P., 2005. Responses of different varieties of *Lolium perenne* to salinity. Annual Conference of the Society for Experimental Biology, Lancashire.
- [23] Kaufman, M.R. and Eckard, A.N., 1971. Evaluation of water stress control by polyethylene glycols by analysis of guttation. *Plant Physiol.* **47**: 453-456.
- [24] Lawlor, D.W., 1970. Absorption of Polyethylene glycols by plants and their effects on plant growth. *New Phytol.* **69**: 501-513.
- [25] Lichtenthaler, H.K., 1987. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods Enzymol.* **148**: 350-382.
- [26] Lutts, S., Almansouri, M. and Kinet, J.M., 2004. Salinity and water stress have contrasting effects on the relationship between growth and cell viability during and after stress exposure in durum wheat callus. *Plant Sci.* **167**: 9-18.
- [27] Miller, G.L., 1959. Use of dinitrosalicylic acid reagent for the determination of reducing sugar. *Anal. Chem.* **31**: 426-429.
- [28] Mobambo, K.N., Gauhl, F., Vuylsteke, D., Ortiz, R., Pasberg-Gauhl, C. and Swennen, R., 1993. Yield loss on plantain from Black Sigatoka leaf spot and field performance of resistant hybrids. *Field crops Res.* **35**: 35-42.
- [29] Morris, A.W. and Riley, J.P., 1963. The determination of nitrate in sea water. *Anal. Chem. Acta* **29**: 272-279.
- [30] Murashige, T. and Skoog, F., 1962. A revised medium for rapid growth and bioassay with tobacco tissue cultures. *Physiol. Plant.* **15**: 473-497.
- [31] Ozyigit, I.I., Kahraman, M.V. and Ercan, O., 2007. Relation between explant age, total phenols and regeneration response in tissue cultured cotton (*Gossypium hirsutum* L.). *Afr. J. Biotechnol.* **6**: 3-8.
- [32] Ruf, R.H., Eckard, E.R. and Gifford, R.O., 1967. Compounds of osmotic adjustment of plants to rapid changes in root medium osmotic pressure. *Soil Sci.* **104**: 159-162.
- [33] Sadeghian, S.Y. and Yavari, N., 2004. Effect of water deficit stress on germination and early seedling growth in sugar beet. *J. Agron. Crop Sci.* **190**: 138-144.
- [34] Slama Ghnaya, T., Hessini, K., Messedi, D., Savoure, A. and Abdelly, C., 2007. Comparative study of the effects of mannitol and PEG osmotic stress on growth and solute accumulation in *Sesuvium portulacastrum*. *Environ. Exp. Bot.* **61**: 10-17.
- [35] Stover, R.H., 1972. Banana, Plantain and Abaca diseases. Commonwealth Mycological Institute, England, pp. 316.
- [36] Turner, N.C., 1981, Techniques and experimental approaches for the measurement of plant water status. *Plant and Soil* **58**: 39-366.
- [37] Vanden, B.L. and Zeng, Y.J., 2006. Response of South African indigenous grass species to drought stress induced by polyethylene glycol (PEG) 6000. *Afr. J. Bot.* **72**: 284-286.

## Author Profile



**Dr. Ikram-ul-Haq**: **2000: M.Sc.** degree (Plant Physiology) from University of Agriculture, Faisalabad, Pakistan. **2002: M.Phil.** degree (Plant Genetic Transformation) from University of Agriculture, Faisalabad, Pakistan; **2012: Ph.D.** degree (Plant Biotechnology) from University of Sindh, Jamshoro, Pakistan; **2012-15: Post-Doctorate:** University of Groningen, Netherland; **Research and Teaching Exp.:** 13 yrs. **Research Specializations:** Cell Stress Physiology in Bacterial and Plants. **Teaching Specializations:** Teaching courses such as Genetic Engineering, Breeding; Cell Hybridization, Cell Ecophysiology. Currently, working as Post-Doctorate Fellow at University of Groningen, Netherlands.