


2.4 Two Point Discrimination:

- Explanation of the procedure to the patient
- Patient position: Sitting straight with hands by the side.
- Trial: Patient will be given a trial to get accustomed to the procedure

Measurement: The Two point discrimination of diabetics will be measured for dominant hand using aesthesiometer. Sensation was tested on a sensory dermatome of the palm, two points were tested on each dermatome C6, C7, C8. Two points of C6 on thumb, two points of C7 on middle finger, two points of C8 on little finger. The two points of the caliper were applied at the same time (stationary two point discrimination test) using weight of the caliper alone. Subjects were asked whether they felt the point as two or one. The minimal distance at which two points were discriminated was measured. The distance between two points of the caliper with which testing was started was 4cms. the distance was decreased further if the subject responded correctly at least three times out of five. The distance of two points was progressively decreased till the subject started perceiving them as one. The scores for two point discrimination test were recorded and analyzed for the comparisons.


3. Stastical Analysis

All the data are presented as Mean±SD. Independent 2 tailed t test is used to find the differences in two point discrimination and grip strength among diabetic and non diabetic. A total of 60 patients (44 males and 16 females) were included for the study. The mean values in diabetic group for age, weight, height and BMI 48.03 ±6.0 years, 67.10±12.5kg, 157.7±10.65 cm and 26.97±4.38 respectively. The mean values in Non diabetic group for age, weight, Height, and BMI are 45.87±4.12 years, 67.90±1.1352 Kg, 162.40±12.13 cm and 25.8±5.2 respectively. The age, weight, height, and BMI showed no difference between the two groups and have the $P \leq 0.05$ as shown in Graph 1.1.




Graph 1.1

The two point discrimination was analyzed using Independent 2 tailed t-test. The mean values for two point discrimination at C6, C7 and C8 of group 1 are 7.43±3 mm, 11.23±3mm and 8.4± 5mm and group 2 are 5.6±2mm, 8.6±3mm and 8.4±3mm. The result showed statistically significant difference between the group ($t_{(1, 11)} = 2.79, 2.74, -0.06$ as mention in the graph 1.2



Graph 1.2

There was shown statistically significant difference between the two groups (group 1 and group 2). The mean values for grip strength of group 1 are (28.74± 9.63) and group 2 (37.5± 15.02) and significant at $p \leq 0.05$ as mention in the Graph 1.3



Graph 1.3

4. Discussion

The Objective of this study was to see the difference in measure between two point discrimination and grip strength among diabetic and non diabetic. Result revealed that statistically significant difference was found between two groups. The present study included 30 type II diabetic patients (experimental group) and 30 non diabetic (control group) including 44 males and 16 females. Two point discrimination and grip strength of the dominant hand of each group was measured. Difference of values between each group was calculated. The result of the study showed that there was significant difference ($p=0.01$) found in the grip strength of diabetic and non diabetic. There is decrease in the grip strength in type 2 diabetic than non diabetic. However, the result of our study confirmed the findings of Ezema et al (2012), Cetinus et al (2004) who have independently reported a decrease in hand grip strength in type2 diabetes.^(2, 3)

Peterson K. F. et al stated that the possible reason for muscle weakness may be insulin resistance as it reduces the rate of glycogen synthesis in muscle⁽⁴⁾. Muscle glycogen is used in type IIb fibres, using anaerobic pathway and in type II a fibres using both aerobic and anaerobic metabolism⁽⁵⁾. The highly potent type IIa fibre permit sustained efforts, due to oxidation of glycogen by the mitochondrion.⁽⁶⁾ Therefore, muscle glycogen content may be particularly decreased in type II a fibres in diabetes⁽⁷⁾. Fuel disposal is also of poorer quality due to insulin resistance and is associated with lower muscle performance⁽⁸⁾.

However, it contradicts the study of Anderson et al (1997) and Anderson et al (2004) who opposed this view and insisted that grip strength is not compromised in long standing type2 diabetes.^(9,10) Upper extremity grip strength evaluation is important as it is required for the basic functions such as manipulating and holding objects. Grip strength is the pre requisite for upper limb functions. Grip strength plays important role in daily activities in working age population. There is very important role of stable grip in every day to day activities of life such as opening a jar, driving a car, driving scooter, riding bike, writing, riding bicycle, carrying a bag of groceries and holding a book, holding a comb etc

The result of this study also showed that there is significant difference in two point discrimination in the dermatomes of hand which are responsible for sensation in hand among the diabetic and non diabetic. Sensations are very important for grip it is the role of sensation which gives the information to the nervous system that how much force is require for gripping an object. Sensations are important for grip as they bring about accuracy in the grip by giving sensory feedback to the nervous system. In this study the two point discrimination was tested because it is the only sensation which can be quantified. The evaluation of sensibility in the hand of the diabetic patient is of great importance in order to provide proper identification of the group that to what extent the sensory changes are coming. Sensory changes in the hand would help to detect the involvement of upper limb neuropathy. A shift in the age of onset of type2 diabetics has been observed in the population with sedentary life style. A huge number of the working population has shifted from manual labour associated with the agriculture sector to physically less demanding office jobs.

Various studies have been done in older population but there is lack of literature focusing on skeletal muscle strength of diabetes in middle age group. It is important to evaluate muscle strength of this age group so that appropriate remedial measures can be taken as from the time of diagnosis because the physical disability or functional dependence is difficult or nearly impossible to rehabilitate later.

The present study took into account the middle aged type II diabetics because currently the greatest number of individuals affected with type II diabetics is between age group of 40-59 years. In this study, two point discrimination sensations were checked in the dermatomes C6, C7 and C8. C6 dermatome is for thumb, thumb which is most important because of its relation with other digits, its loss can affect hand function greatly. C7 is for the middle finger, in flexion middle finger is the strongest and it is important for precision and power grip. Two point discrimination in C7 shows significant difference ($p=0.01$) among the diabetic and non diabetic. But there was no significant difference ($p=0.95$) in two point discrimination which is provided by dermatomes of C8 that is the little finger. In terms of functional impairment there is only 10% role of little finger which is very less even in normal population⁽¹¹⁾.

5. Clinical Relevance

If grip strength and two point discrimination are assessed at the time of diagnosis and routine assessments are done during clinical follow up, deterioration in grip strength and development of disability in the later stage can be prevented. Preventive modalities like resistance training can be incorporated to decrease the rate of deterioration of muscle strength. Strength training from time of diagnosis will help in maintaining the insulin level. Thus it will be helpful in delaying the late musculoskeletal complications occurring in type II diabetes patients due to deterioration in muscle strength. Latest research conducted by Adrienne Visocchi et al (2010) show that handgrip exercises make blood vessels more flexible, improve blood vessel function, and lower high blood pressure. The hand exercises performed in this study significantly helped to lower systolic blood pressure but Diastolic blood pressure did not change.¹²

6. Limitation of the Study

- 1) Duration of diabetes has been taken into account from time of diagnosis however individual may have subclinical type II diabetes for many years without knowing until hyperglycaemic and neuropathic symptoms manifest thus posing great difficulty in tracking the effect response trend between the duration of diabetes and muscle strength.
- 2) Study was performed on a relatively small sample size. (n=60)
- 3) Data collected was from a specific geographical region hence the result could not be generalized.

7. Future Research

- 1) Future research with larger sample size can be undertaken.
- 2) Data for two point discrimination can be generated for various cutaneous areas of the body.
- 3) Correlation with level of hyperglycaemia affecting grip strength and sensations in type 2 diabetes.

8. Conclusion

This result of this study showed that there is difference in two point discrimination and decrease in grip strength in type II diabetes patients as compared to non diabetics. The present study will be helpful for type II diabetics to prevent further deterioration in hand sensations and muscle strength.

References

- [1] Mohan, S. Sandeep, R. Deepa, B. Shah and C. Varghese, Epidemiology of type 2 diabetes: Indian scenario, Indian Journal of Medical Research 125, March 2007, 217-230
- [2] Gerard Said. Diabetic Neuropathy—a review. Nat Clin Pract Neurol 2007;3:331-40.
- [3] J.G.E,mer., and Spinner, D., 1980: Sensibility Testing, Management of Peripheral Nerve Problems. Philadelphia, PA: WB Saunders Co. 3-15

- [4] V. Mathiowetz, K.Weber, G. Volland, N. Kashman, Reliability and validity of grip and pinch strength evaluations, *J. Hand Surg. 9A* (1984) 222–226.
- [5] Ezema, Iwelu et al, *AJPARS* Vol. 4, Nos. 1 & 2, June 2012, pp. 67 – 71
- [6] Ercan Cetinus, Mehmet Akif Buyukbese et al, *journal of Diabetes Research and Clinical Practice* 70 (2005) 278–286.
- [7] .Peterson Kf ,shulman Shulman GI,Pathogenesis of skeletal muscle insulin resistance in type2 diabetes mellitus, *American journal of cardiology* 2002;90:11G-18G
- [8] ShermanW.M. Metabolism of sugars and physical performance.*Am J Clin Nutr* 1995;62:228S-241S
- [9] He J,Kelley DE. Muscle glycogen content in type2 diabetes mellitus.*Am j physiology endocrinol Metab* 2004;287:E1002-7.
- [10] Anderson,H.,P.C.Gadeberg,B. Brock and J.Jakobsen 1997.Muscular atrophy in diabetic neuropathy: A stereological magnetic resonance imaging study.*Diabetologia* 40:1062-1069
- [11] Andersen H, Nielsen S,Mogensen CE,Jakobsen J:Muscle strength in type2 diabetes.*Diabetes* 2004;53:1543-1548
- [12] Magee, David J. *Orthopedic physical assessment* edition pg 419. Elsevier Health Sciences, 2014.
- [13]Cassandra Stiller-Moldovan, "The effects of isometric handgrip exercise on post-exercise hypotension, ambulatory arterial blood pressure and heart rate variability in individuals medicated for hypertension" (2010)Paper 279.

