

# Parasites on Appendicularia (*Oikopleura (Vexillaria) dioica*) from Sea of Marmara, Turkey

M. Levent Artüz

Sevinç-Erdal İnönü Foundation, Department of Marine Sciences, Anadoluhisarı Toplarönü No: 8, 34810, Istanbul, Turkey  
e-mail: levent@artuz.com

**Abstract:** In the various sites in Sea of Marmara, Appendicularian *Oikopleura (Vexillaria) dioica* were invaded by *Oodinium* sp. This study provides the first evidence of *Oodinium* sp. in the Sea of Marmara, Turkey. High probability of a fatal attack to the zooplankton communities will affect directly to the biodiversity via food web and finally to the fishery of the area. The effect may grow or/and spread exponentially to the Black Sea and Mediterranean Sea, because of the oceanographical and biological specialities of the Sea of Marmara and its connections through Turkish straits. This study emphasizes the consequences of parasitic infection in the case of the special position of Sea of Marmara.

**Keywords:** Appendicularia, *Oodinium*, Parasites, Sea of Marmara, Zooplankton.

## 1. Introduction

Parasitism may reduce zooplankton's general condition with lethal effect [1], and it has developed effects on the biodiversity and to the sustainability of stocks, which appears important to assess this case.

Representatives of the genus *Oodinium* are common parasites of Appendicularia. They have been classified as Infraphylum Dinoflagellata Bütschli 1885; Class Dinophyceae Fritsch, 1927; Order Blastodiniales Chatton, 1906; Family Oodiniaceae Chatton 1920. The genus *Oodinium* Chatton, 1912 consists of four species – *Oodinium acanthometrae* Chacon, 1964; *Oodinium fritillariae* Chatton, 1912; *Oodinium inlandicum* T.Horiguchi & S.Ohtsuka, 2001; *Oodinium pouchetii* (Lemmermann) Chatton, 1912 [2].

A large number of dinoflagellates are known to parasitize marine vertebrates and invertebrates e.g., [3], [4], [5]. One of these, the family Oodiniaceae, has members which are characterized by being ectoparasitic and possessing suboval to fusiform cells with well-developed absorption apparatuses in the trophont stage [6].

*Oodinium* spp. has been noted in many species of Appendicularia, including *Oikopleura (Vexillaria) dioica* Fol, 1872. This study provides the first evidence of *Oodinium* sp. in the Sea of Marmara, Turkey.


## 2. Materials and Methods

Plankton samples were collected between years 2007 and 2012 by annual surveys as part of the MAREM (Marmara Environmental Monitoring) project. The details of hauls are given in Table 1 and the station locations are given in Fig. 1.

Plankton net with a 650 mm net diameter and of 0.180 mm mesh size were used as sampling gear. Two replicates were taken at each station. For the vertical haul, the net lowered to the maximum depth and hauled all the water column, for oblique hauls used closed nets with same sizes from the depth of thermocline layer (average 25 m depth) and horizontal hauls were realized of 0.5m depth, with 5 min duration.

Samples were preserved immediately after collection, in a 4% solution of borax-buffered formaldehyde in seawater. After fixation, samples were precipitated by waiting; liquid level minimized material were rinsed and random sample replicas of 10 pieces of each 1 cm<sup>3</sup> were picked up from each sample group.

Infected and non-infected individuals for each 1cm<sup>3</sup> replica were counted and the average percentages were calculated. All samples are bar-coded and listed in MAREM database.



**Figure 1:** Map of the Sea of Marmara and the location of the sampling stations

### 3. Environment

The entire system of the Turkish Straits and the Sea of Marmara extends from the Aegean Sea to the Black Sea through the Dardanelles (60 km), the Sea of Marmara (210 km), and the Bosphorus (31 km) [7].

The total length of the system thus is approx. 300 km because of the great differences in density between the waters of the Mediterranean and those of the Black Sea, there is a two-layered current system along the Sea of Marmara and the Turkish Straits, flowing in opposite directions [7].

The water masses of the Black Sea are entirely different from those of the Mediterranean proper. In the Black Sea, the combination of precipitation and runoff exceeds evaporation, and accordingly a surface layer of relatively low salinity and correspondingly low density is present [8].

Vertical mixing tends to reduce the density of the deeper surface layer (approx. 200 m), which therefore has a lower density than that of the water in the Aegean Sea at the same depth [9], [10], [11].

Owing to the distribution of density, the surface waters of the Black Sea flow through the Bosphorus and the Dardanelles into the Aegean basin, and Mediterranean water flows in the other direction along the bottom.

The outflow from the Black Sea basin is a function of its water budget and carries runoff from the large rivers and other surface waters that flows into the Black Sea towards the Mediterranean [7].

The outflow also is a function of the excess of precipitation that has come down on the 423 000 km<sup>2</sup> surface of the Black Sea.

Intensive mixing takes place along the Bosphorus as well as in the other parts of the straits. The salinity of the inflowing Mediterranean water, which is over 38.5‰ Sal at the entrance of the Dardanelles, decreases slowly, with the distance travelled in the Sea of Marmara down to 29‰ Sal, where the bottom current enters the Black Sea at the northern end of the Bosphorus.

At the same time, the salinity of the outflowing water increases from 16‰ Sal to 30‰ Sal when leaving the Dardanelles. This increase is a function of the distance travelled by the water.

The out- and inflowing water masses are separated by a well-defined layer of transition, which oscillates up and down according to the contours of the bottom.

This transition layer also represents the discontinuity layer for temperature and salinity and hence is called thermohaline, and is practically null below 200 m [7]. In other words, a Black Sea-originated water mass with a low salinity

and a Mediterranean-originated water mass with high salinity are located one over the other, in basins of 1000 m and deeper that are surrounded and limited by the shallow shelf areas.

The effect is a variable temperature of the surface water mass, fluctuating, dependent on meteorological conditions, between 6 and 29°C annually. In contrast, 14.2°C is the constant mean temperature of the water under the thermocline interface [7].

### 4. Results

Total 250 infected individuals of *Oikopleura (Vexillaria) dioica* were examined for presence of *Oodinium* spp. (Fig. 2. h) The mean number of *Oodinium* sp. on *O. (Vexillaria) dioica* individuals were  $4.07 \pm 2.32$ , with a maximum 11. (mean n° ± st.dev.).

The points of attachment of *Oodinium* sp. on *O. (Vexillaria) dioica* were: tail fin (mainly on the proximal side); trunk (mainly on the ovary section and on the stomach section respectively) (Fig. 2. a, b).

Percentage of the infected individuals of *O. (Vexillaria) dioica* was  $13.7\% \pm 6.16$  (mean% ± st.dev), the sampling period and station based percentage of infected individuals to uninfected individuals are given in table 1.



Figure 2: (a) *Oodinium* sp. on *O. dioica* (b) *Oodinium* sp. on *O. dioica* detail.

**Table 1:** Date and station based distribution of the percentages of infected individuals to the non infected individuals. Mean values of 10 replicates of 1cm<sup>3</sup> from each station. (%±st. dev.) (BC) Beginning and (EC) Ending Coordinates of haul, Station Depth and (HM) Hauling Method (V; vertical, O; Oblique, H; horizontal)

Date	Station nr	Depth (m)	BC	EC	HM	Total N <sup>o</sup> of Individuals	Infested Individuals (%)
09/08/2007	36	41	40° 31.967' N / 026° 59.983' E	40° 31.900' N / 026° 59.850' E	H	141	11.8±1.61
05/08/2009	17	427	40° 40.617' N / 028° 33.933' E	40° 40.450' N / 028° 34.500' E	O	98	13.9±1.51
06/08/2009	24	45	40° 31.883' N / 027° 59.917' E		V	112	23.1±2.10
11/08/2009	40	1100	40° 48.233' N / 027° 27.550' E	40° 48.167' N / 027° 27.433' E	O	75	7.9±1.56
11/08/2010	17	455	40° 40.417' N / 028° 33.100' E	40° 40.167' N / 028° 32.283' E	O	81	13.1±2.20
06/08/2011	30	38	40° 21.900' N / 027° 42.983' E	40° 21.833' N / 027° 43.017' E	H	136	4.8±1.41
07/08/2011	40	800	40° 48.167' N / 027° 26.500' E	40° 48.200' N / 027° 25.500' E	O	90	5.5±0.97
07/08/2011	40	1000	40° 48.183' N / 027° 27.600' E	40° 47.733' N / 027° 26.667' E	O	88	10.1±1.00
07/08/2011	34	187	40° 54.233' N / 027° 33.633' E		V	108	12.5±2.69
08/08/2011	35	55	40° 28.750' N / 026° 59.533' E	40° 28.683' N / 026° 59.483' E	H	144	16.2±1.25
22/07/2012	19	59	40° 56.600' N / 028° 33.583' E		V	132	17.9±2.79
26/07/2012	3	1200	40° 42.400' N / 029° 04.350' E	40° 42.450' N / 029° 04.967' E	O	107	16.4±3.80
28/07/2012	26	44	40° 25.083' N / 028° 02.967' E		V	131	24.5±2.39

## 5. Discussion

During plankton surveys in Sea of Marmara, specimens of the appendicularian *Oikopleura dioica* infected with *Oodinium* spp. were collected. This study describes the parasitic attack of *Oodinium* spp. to appendicularia and the potential impacts, in case of the unique conditions of the Sea of Marmara.

This finding seems important, Appendicularians are important for the fishery of the area directly and via food web [12]. Capitanio et al. (1997) [13] also indicated that the appendicularia are important food for anchovy and anchovy is the most consummated and landed economically important fish species of the district. Any possible exponential negative impact that originated from the beginning of the food chain, will effect directly to the fishery economy of the area.

The enormous effects would arise by the spread of any kind of the Sea of Marmara sourced negativity, especially given that it merges Black Sea originated upper layer water mass with Mediterranean Sea through Aegean Sea and also Mediterranean Sea originated limited scale of water mass with Black Sea, as a conveyor.

Meanwhile the Sea of Marmara has been subjected to various human impacts that have led to changes in the ecology of this inland sea.

The main problem in the Sea of Marmara since 1980 is the growing pollution. All the settled areas around the Sea of Marmara are discharging the wastes using 'deep sea discharges' directly under the pycnocline–thermocline layer in the Sea of Marmara, without any treatment [7].

In my opinion, the long term changing of the chemical environmental variables such as pH changes (Suppl. Data), increasing pesticide concentration [14] and/or increasing amounts of chlorine in the Sea of Marmara, that originated in large quantities as a result of the cleaning fouling organisms

on the inside of the direct discharge pipes of sewage and power plants, discharged directly to the water column, are effecting to the hosts immunity and making them more susceptible to the parasitic attack; but this hypothesis requires further study.

The occurrence of a possible attack of the parasite *Oodinium* spp. to *O. (Vexillaria) dioica* is significant because the subject has high social and economic importance. Besides the direct affect of growing pollution, the uniqueness of the Sea of Marmara from oceanographical and geological point of view plays an important role for the social and economic structure of the area.

High probability of a fatal attack to the zooplankton communities will affect directly to the biodiversity via food web and finally to the fishery of the area.

The effect may grow or/and spread exponentially to the adjacent seas, because of the oceanographical and biological specialities of the Sea of Marmara and its connections trough Turkish straits.

It can be imagined the Sea of Marmara as a biological corridor between two totally different basins with different characteristics such as Black Sea and Mediterranean Sea.

## 6. Acknowledgements

I acknowledge the Sevinç-Erdal İnönü Foundation that oversaw the long-term Marmara Environmental Monitoring project and O. Bülent Artüz for their assistance with programming, data acquisition, and data merging, and especially for the electronic field support for the project. I also acknowledge the crew of the vessel 'Oktay 4' for their help in collecting the material.

## References

- [1] Borgsteede, FHM (1996) The effect of parasites on wildlife. *Veterinary quarterly* 18(sup3): 138-140
- [2] Guiry, Michael D. (2015) *Ellobiopsis* Caullery, 1910. In: Guiry, M.D. & Guiry, G.M. (2015). *AlgaeBase*. World-wide electronic publication, National University of Ireland, Galway (taxonomic information republished from *AlgaeBase* with permission of M.D. Guiry). Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=562628> on 2015-04-07
- [3] Chatton E (1920) Les Péridiniens parasites. Morphologie, reproduction, éthologie. *Arch. Zool. exp. gen.* 59: 1-475.
- [4] Cachon J, Cachon M (1987) Parasitic dinoflagellates, p. 571-610. In *The Biology of Dinoflagellates* (ed. Taylor, FJR). Botanical Monographs Vol. 21. Blackwell Scientific Publications, Oxford.
- [5] Ohtsuka S, Nagasawa K, Gejima K (2000) Review of parasites of marine zooplankton. *Bull. Plankton Soc. Jpn* 47: 1-16. (in Japanese with English abstract)
- [6] Horiguchi T, Ohtsuka S (2001) *Oodinium inlandicum* sp. nov. (Blastodinales, Dinophyta), a new ectoparasitic dinoflagellate infecting. *Plankton Biol. Ecol.* 48(2), 85-95.
- [7] Artüz ML, Okay IA, Mater B, Artüz OB, Gürseler G, Okay N, (2007) Scientific Aspects of the Sea of Marmara. Union of Turkish Bar Associations publication, Istanbul (in Turkish)
- [8] Artüz İ, Baykut F, Aydın A (1982) Scientific Aspects of the Black Sea. Istanbul University Publication, Engineering Faculty, Fatih publisher, Istanbul (in Turkish)
- [9] Artüz ML, Artüz OB, Gülen D, Torcu KH, Üzen E, Aydemir A, Sönmez B, Akdemir D, Bakacak A (2009) Changing Oceanographic Conditions of the Sea Of Marmara (MAREM) 2008 survey reports. T.C. Piri Reis University publication, Istanbul (in Turkish)
- [10] Artüz ML, Artüz OB, Aydın A, Gülen D, Koc HT, Akdemir D, Aydemir A, Sönmez B, Üstün F, Serter U, Küçük Ö, Bakacak A, Uzun D, Bulut FM (2010a) Changing Oceanographic Conditions of the Sea Of Marmara (MAREM) 2009 working reports. Marmara University publication, Istanbul (in Turkish)
- [11] Artüz ML, Artüz OB, Aydın A, Gülen D, Koc HT, Yalçın B, Akdemir D, Aydemir A, Sönmez B, Üstün F, Ulunehir G, Serter U, Küçük Ö, Kasap H, Baki C, Erzaim N, Bulut FM, Özdemir F (2010b) Changing Oceanographic Conditions of the Sea Of Marmara (MAREM) 2010 working reports. Marmara University publication, Istanbul (in Turkish)
- [12] Shiganova TA, Bulgakova YV (2000) Effects of gelatinous plankton on Black Sea and Sea of Azov fish and their food resources. *ICES Journal of Marine Science: Journal du Conseil* 57(3): 641-648
- [13] Capitanio FL, Pájaro M, Esnal GB (1997) Appendicularians (Chordata, Tunicata) in the diet of anchovy (*Engraulis anchoita*) in the Argentine Sea. *Scientia Marina* 61(1): 9-15
- [14] Aksu A, Taşkın ÖS (2012) Organochlorine residue and toxic metal (Pb, Cd and Cr) levels in the surface sediments of the Marmara Sea and the coast of Istanbul, Turkey. *Marine pollution bulletin* 64(5): 1060-1062.