# Antibacterial Spectrum and Activity Guided Evaluation of Bark of *Sonneratia Alba*

# Mitter C. S.

Department of Zoology, Sathaye College, Vile-Parle (E), Mumbai, 400 057, Maharashtra, India

Abstract: The present study has been undertaken to investigate bioactive compounds from bark of Sonneratia alba. Extracts were made in acetone, ethyl acetate and water by using powder of bark of S. alba. Bioassay of primary screening and MIC were done by disc diffusion method using various gram negative and gram positive bacteria. Maximum inhibition was shown against Klebsiella pneumoneae by acetone and aqueous extract and against Corynebacterium diphtheriae by aqueous extracts. MIC of all potent extracts displayed promising inhibitory activity against susceptible microorganisms. Activity Guided Fractionation of highly potent extracts indicated active principles mainly dissolved in polar solvents.

Keywords: Bioactive compounds, S. alba, Bioassay, MIC and activity guided fractionation

## 1. Introduction

Since ancient times plants have played an important role in the discovery of new therapeutic agents. Much of the scientific efforts made in the past few decades with medicinal plants have focused on documenting the uses of traditional medicine, effectiveness of particular remedies characterizing medicinal plants to their compound and testing plant compounds *in vitro*. [1], [2].

A special feature of higher plants is their capacity to produce a large number of secondary metabolites which are produced for self defense. [3]. Human beings still rely on plants to combat bacterial and fungal infections. Microorganisms have developed a resistance to many antibiotics and as a result, immense clinical problems in the treatment of infectious diseases have been created. [4].

## 2. Literature Survey

Mangroves are 'coastal woodlands' or marine tidal forests comprising of trees, shrubs, palms, epipyhytes, ground ferns and grasses. They occupy large tracts along sheltered coasts, estuaries and in deltas. They are specially adapted to withstand salinity, wave action and can grow in poor soil. Mangroves occur approximately in 112 countries. [5]. The Sundarbans are the largest mangrove forest in the world, both in size as well as biodiversity. Mangroves are important source of tannins used for dying, leather production and oil drilling. They provide food, timber, variety of traditional products and folklore medicine. [6] and [7]. Mangroves are biochemically unique producing a wide array of novel natural product. [8]. Several biological activities are attributed to different mangroves. [9], [6] and [7]

Various phytochemicals are well known for their antimicrobial properties and can be of great significance in therapeutic treatments. Increasing resistance to the antimicrobial agents is very common in patients with various infections. [10].

*S. alba* consists secondary metabolites like steroids, polyphenols, glycosides and condensed as well as hydrostable tannins. The pneumatophores of *S. alba* and *S.* 

*caseolaris* are used in making corks, fishing floats and heels of shoes. **[11]**. The leaves, trunk and bark have been reported for its antoxidant properties. **[12]**. Folklore medicinal uses of *S. alba* include treatment of various skin ailments. **[13]**.

Activity guided fractionation of various mangroves were studied. **[14]**, **[15]**. Characterization of active principles of *S. apetela* was studied by various researchers. **[16]**, **[17]** and **[18]**. *S. apetela*, *R. mucrunata* and *B. cylindrica* were found to be highly potent against various pathogens. Antimicrobial activities of various parts of *S. alba* made in organic solvents were also investigated. **[19]** and **[20]**.

## **3. Problem Definition**

There is an urgent need to explore and discover new antimicrobials with diverse chemical and novel mechanical action for new and re-emerging infectious diseases. In recent years secondary plant metabolites, previously with unknown pharmacological activity have been extensively investigated as a source of medicinal agents. In the context of this a study was undertaken to evaluate the activity of bark of *S. alba* against several gram positive and gram negative bacteria *in vitro*.

## 4. Materials and Methods

#### **Plant Material**

Mangrove species *S.alba* was collected from Mumbra estuaries, District Thane, Maharashtra, India. The bark was washed with tap water and dried in shade, finely powdered and stored in airtight bottle.

#### **Preparation of Extracts**

1 gram of air dried powder of bark was immersed in 20 ml of organic solvents acetone, ethyl acetate, and distilled water separately to make crude extracts. It was incubated at room temperature for 48 hours at 150 rpm on an orbital shaker. The suspensions were filtered, made 20 ml and stored at 4°C for antimicrobial activity.

#### Microorganisms

Six gram negative bacteria *Escherichia coli* (*E.coli*), *Klebsiella pneumoneae* (*K.p*), *Salmonella typhii* (*S.t*), *Salmonella paratyphii A* (*Sp.A*), *Salmonella paratyphii B* (*Sp.B*) and *Pseudomonas aeruginosa* (*Ps.aeru*), four gram positive bacteria *Staphylococcus aureus* (*S.a*), *Bacillus subtilis* (*B.subt*), *Streptococcus pyogens* (*S.pyo*) and *Corynebacterium diphtheriae* (*C.d*) were used for bioassay. All the strains were collected from Haffkine Institute, Parel, Mumbai and sub cultured at regular intervals. These microbes were maintained on nutrient agar slant and stored at 4°C.

#### Media used

Nutrient agar and nutrient broth (Hi Media).

#### **Preparation of Inoculums**

Active cultures for experiments were prepared by transferring a loopful of culture to nutrient agar slant (Hi Media), incubated at 37°C for 24 hours for bacterial proliferation.

#### **Bioassay**

Paper disc method **[21]** was employed for testing antibacterial activity. Sterile sugar tubes with 20-25 ml sterile nutrient agar and18-24 hours old 0.4 ml of culture made in nutrient broth of each pathogen, mixed and poured on each sterile plate. The plates were kept aside for agar solidification. Paper disc (5 mm in diameter) soaked into plant extract were placed onto the plates and the plates were incubated at 37°C for 24 hours **[22]** to allow the bacterial growth. The antibacterial activity of the bark extract was determined by comparing the diameter of the zone of inhibition with that of respective controls (acetone, ethyl acetate, and distilled water). For each bacterial strain, controls were maintained where pure solvents were used instead of the extract. The experiment was conducted in triplicate and the mean values are represented.

#### **Minimum Inhibitory Concentration (MIC)**

In primary screening, the crude extracts which displayed zone of inhibition more than 15 mm in diameter were selected for MIC studies. Each extract was evaporated at room temperature to dry residues and then using fresh solvent, different concentrations viz.10mg/ml, 5mg/ml, 1mg/ml and 0.5mg/ml were prepared. MIC was conducted with all concentrations by above mentioned disc diffusion method.

#### **Bioactivity Guided Fractionation:**

Acetone Extract: The extract was evaporated to dryness. Then chloroform, petroleum ether, ethyl acetate, acetone, ethanol, methanol and water were added successively with centrifugation at 3000 rpm for 20 minutes. All the fractions of solvent filtrates separated and kept in airtight borosil bottles at 4°C and tested by disc diffusion method.

Aqueous extract: The water extract was evaporated to dryness. Then chloroform, petroleum ether, ethyl acetate, acetone, ethanol methanol and water were added successively with centrifugation at 3000 rpm for 20 minutes. All the fractions of solvent filtrates separated and kept in

airtight borosil bottles at  $4^{\circ}$ C and tested by disc diffusion method.

## 5. Result

Antimicrobial assay: In vitro antimicrobial activity is summarized in **Table-I.** Overall all bark extracts exhibited zone of inhibition ranging between 12mm - 22mm against sensitive pathogenic strains. Comparatively less activity was noted by ethyl acetate. Maximum inhibitory activity was shown with zone of inhibition exhibited by acetone extract against K. pneumoneae (22mm) followed by 21 mm zone of inhibition against C. diphtheriae by aqueous extract. Furthermore, it was observed that B.subtillis, S. paratyphi-A, S. paratyphi-B and Ps. aeruginosa were highly resistant against all test extracts. MIC of acetone extract revealed antimicrobial activity at 5 mg/ml against sensitive strains and by aqueous extract and at 1 mg/ml against susceptible bacteria.

Bioactivity Guided Fractionation indicated that active principles are dissolved in polar solvents.

# 6. Discussion

This investigation by primary screening reveals that active principals are dissolved in various solvent with different potency. Higher degree of potent bioactive phytochemicals is in acetone and in aqueous extracts. It was noted that in hot acetone extract of leaves S. apetala showed active principles are present. [16]. Other researchers [19]; [20]; [23] noted that extracts made in organic solvents such as methanol, ethanol, ethyl acetate, chloroform and water of various parts of S. alba were active against S. aureus, B. cereus, E. coli and Ps. aeruginosa. Similar findings were also noticed for various mangrove species. [14]; [15]; [16]. These earlier observations support the present study which indicates mangroves are loaded with antimicrobials which are dissolved in organic solvents. Bioactivity Guided Fractionation indicated that active constituents are dissolved in polar solvents. Similar findings were noted in activity guided fractionation of other mangrove plants occurring along Mumbai coast. [14]; [15]; [24]; [18]. These findings are in agreement with the present investigations.

# 7. Conclusion

This investigation supports that bioactive principles present in *S. alba* are active against sensitive gram positive and gram negative bacteria. It also reveals that mangrove species contains methanol, ethanol, acetone and water soluble antimicrobials. Further attempts may be made to isolate active principals which can be use in drug development.

# 8. Future Scope

It is suggested that the entire plant may be further explored for their active ingredients and further investigation is needed for its chemical composition in view of its pharmaceutical importance. Therefore it is suggested that methanol, ethanol, acetone and water can be use to extract bioactive principles from the mangrove species *S. alba*.

## International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2013): 6.14 | Impact Factor (2013): 4.438

 Table 1: Primary Screening of crude bark extracts of S.
 alba (Inhibition Zones in mm)

|         | B.subt | S.pyo | S.a | K.p | C.d | E.coli | S.t | Sp.A | Sp.B | Ps.aeru |
|---------|--------|-------|-----|-----|-----|--------|-----|------|------|---------|
| Acetone | -      | 18    | 15  | 22  | -   | -      | 16  | -    | -    | -       |
| Ethyl   | -      | 13    | 15  | 15  | 14  | -      | 15  | -    | -    | -       |
| acetate |        |       |     |     |     |        |     |      |      |         |
| Water   | -      | 13    | -   | 22  | 21  | 12     | -   | -    | -    | -       |

Keywords: - No zone of inhibition.

 Table 2: MIC of potent extracts

| Solvent       | Pathogen | Concentration mg/ml. |    |    |     |  |  |  |
|---------------|----------|----------------------|----|----|-----|--|--|--|
|               |          | 10                   | 5  | 1  | 0.5 |  |  |  |
| Acetone       | S.pyo    | 20                   | 15 | 1  | -   |  |  |  |
|               | S.a      | 18                   | 14 | 1  | -   |  |  |  |
|               | K.p      | 17                   | 12 | -  | -   |  |  |  |
|               | S.t      | 17                   | 12 | -  | -   |  |  |  |
| Ethyl acetate | S.a      | 15                   | -  | -  | -   |  |  |  |
|               | K.p      | 14                   | -  | -  | -   |  |  |  |
|               | S.t      | 14                   | -  | -  | -   |  |  |  |
| Water         | K.p      | 26                   | 19 | 13 | -   |  |  |  |
|               | C.d      | 26                   | 18 | 13 | -   |  |  |  |

Keywords: No zone of inhibition.

| Tables 3: Activity | Guided fractionation |
|--------------------|----------------------|
|                    |                      |

| Extracts | Pathogens     | Fractions |   |   |    |    |    |    |   |
|----------|---------------|-----------|---|---|----|----|----|----|---|
|          |               | а         | b | с | d  | e  | f  | g  | h |
| Acetone  | Kl. Pneumonia | —         |   | — | 12 | 13 | 13 | _  |   |
|          | S.pyogens     |           | _ |   | 13 | 12 | 13 |    |   |
|          | S.aureus      | —         |   | — | 13 | 14 | 14 | _  |   |
|          | S. typhi      | _         | _ | _ | 11 | 12 | 13 |    |   |
| Water    | C.diphtheriae | —         |   | — |    | 13 | 13 | 15 |   |
|          | Kl. Pneumonia |           | _ |   |    | 12 | 11 | 13 |   |

Keywords: -- No zone inhibition, petroleum ether

a- chloroform

- b- Petroleum ether
- c- ethyl acetate
- d- acetone
- e- ethanol
- f- methanol
- g- water

# References

- [1] Fowler M.W. (2006). "Plants and Medicines and Man." Journal-Science, Food and Agriculture, 86:17971804.
- [2] Gertsh J (2009). "How Scientific is the Science in Ethnopharmacology? Historical perspectives and epistemological problems." J. Ethpharm, 122@17783).
- [3] Evan J.S. pattison and Morris P: "Antimicrobial agent in plant culture" edited by P. Morris, A. Scarges, Cambridge university, London-1986, pp:12.
- [4] Davies J, 1994. 'Inactivation of Antibiotic And The Dissemination Of Resistant Gene.' Science Journal, 264; Pg. 375-382
- [5] Bandaranayake W.M, 1998. Traditional and Medicinal Use of Mangroves. Mangrove and Salt Marshes, 2: 133-148.
- [6] Kathiresan K. and Qasim S.Z. (2005). 'Biodiversity of mangrove ecosystems.' Hindustan Publishing Corporation (India) New Delhi, pp.5.
- [7] Bandaranayke, W.M. (2002). 'Bioactivities, bioactive compounds and chemical constituents of mangrove

plants.' Wetlands Ecology and Management, 10: 421-452.

- [8] Hogg R.W and Gillan F.T, (1984). Fatty acids, steroids and hydrocarbon in the leaves eleven species of mangroves. *Phytochemistry*.23: 93-97.
- [9] Kokpol U. and Chittawong V., (1987). Chemical constituents of *Acanthus illicifolius* Linn. And biological activity, UNESCO regional on the chemistry of mangrove plants, Bangkok, Thailand. 40-42.
- [10] Obadoni B.O and Ochuko P.O (2001) Phytochemical Studies and Comparative efficacy of the crude extracts of some Homostatic plants in Edo and Delta States of Nigeria Global J. Pure.Appl.Sci.Vol 8.203-208.
- [11] Walsh G.E., (1997). Exploitation of Mangal. In: Ecosystems of the World. Chapman V.J. (Ed.). Pub: Elsevier Scientific, New York. 335-347.
- [12] Wada, K., A. Kinjo, M. Uehara and K. Takara, 2002. Antioxidant activities of mangrove trees. Science Links jpn. 189-192.
- [13] Wang R.J and Z.Y. Chen, 2002. 'Systematics and Biogeography Study On The Family Sonnoretiaceae.' Guihaia, 22:214-219.
- [14] Shah N.J, (2005). Pharmacognosy Studies on Methanol Extracts of Mangrove Species Aegicerus corniculatum. Ph.D Thesis (Life Sciences), University of Mumbai.
- [15] Mane S, and Jadhav B.L., (2004). Pharmacognosical Investigations of Bruguiera cylindrical. In: Proceedings of National Symposium, Biosciences, Biotechnology and Bio-enterprises. 236-240.
- [16] Dhane J.M., (2005). Studies on the Acetone Soluble Bioactive Principles of Sonneratia apetala leaf. Ph.D Thesis (Life Sciences), University of Mumbai.
- [17] Maurya B. R., (2001). "Studies on Screening of Bioactive Compounds of Mangrove Plants Along Revas Coast." M.Sc. Thesis (Life Sciences) University of Mumbai, Mumbai, India. P 49.
- [18] Jhurani B.S, 2009. 'Studies On The Bioactivity Guided Isolation And Purification Of Natural Compounds *R. apiculata*.' Ph.D Thesis (Life Science) University of Mumbai, 400098.
- [19] Haq I., Hossain S., et. al., (2014). Antioxidant and Antibacterial Activities of Different Extracts of Fractions of a Mangrove Plant S. alba. International Journal of Agriculture and Biology. 707-714.
- [20] Saad S., Taher M., et. al., (2012). In Vitro Antimicrobial Activity of Mangrove Plant Sonneratia Alba. Asian Pacific Journal of Trop. Biomed. 2(6): 427-429.
- [21] Gould J.C and Bowie J.H, 1952. 'The determination of bacterial sensitivity to antibiotics.' Edinb. Med. J. 59: 178.
- [22] Cappucino J.G and Sherman N, 1999. Microbiology a laboratory manual, Addison Wesley Langman Inc., California, 4<sup>th</sup> edition, 263.
- [23] Sahoo G., Mulla S., et. al., (2012). Antibacterial Activity of Mangrove Leaf Extracts Against Human Pathogens. Scientific Publification of the Indian Pharmaceutical Association. Vol-74, issue-4. Pp.348-351.
- [24] Kale A.P., (2006). Screening of Pesticidal Principals from Mangrove Species Occuring along Mumbai Coast. Ph.D Thesis (Life Sciences), University of Mumbai.

# **Author Profile**



**Prof. (Dr.) Chhaya S. Mitter** is Head of Zoology Department, Sathaye College, Vile Parle (E), University of Mumbai, Mumbai-400057, Maharashtra, India. Associate Professor for 26 years. Successfully

completed university research projects. Published various research papers. Worked on various bodies in the college and university. Active in social fields. Worked as associate editor for Bio-chemical Science.