



**Characterization of *Trichoderma*.**

After the purification process, each colony of *Trichoderma* isolates was taken using a 0.5 cm bore cores then placed in the middle of new sterile solid PDA and incubated at room temperature (25°C - 27°C) for further used in macroscopic and microscopic observation. Identification of *Trichoderma* done by macroscopic and microscopic observation. The macroscopic identification was conducted by observing the morphology, colony growth pattern, density, color, mycelia zoning, and the diameter of *Trichoderma* colonies grown in the sterile solid PDA medium. Microscopically, the identification was performed by using a microscope observed at the age of 3 days.

For the microscopic observation, the procedure stages started with cleaning the the object glass with 70% alcohol previously before few drops of Lactophenol-Cotton Blue solution was given at the center of the glass object. A part of *Trichoderma* culture was taken aseptically using a loopful then placed on the glass object that has been spilled with the Lactophenol-Cotton Blue and subsequently covered with cover glass and observed under a microscope. Microscopic observations were made using a light microscope (Trinokuler Zeiss Primo Star iLED, 400x) and fungal colonies were identified by observing the conidia, conidiophores and phyalid described by Barnett and Hunter (1978).

**Assessment of seedling growth enhancement ability.**

**Table 1:** Cultural characteristics of *Trichoderma* indigenous strains on PDA medium isolated from rhizosphere of Maize growing in dry land of Maros, Takalar and Jeneponto regencies

<i>Trichoderma</i> Origine	Colony character				Zonation
	Texture	Density	Surface color	Reverse color	
Maros	Thick and velvety	Media	Dark green	Colorless	Concentric zones with green on first and third zones and white on second zone
Takalar	Thick and velvety	Media	Dark green	Hyaline with whiteish at the centre	Concentric zones with green on first and third zones and white on second zone
Jeneponto	Thick and velvety	Media	Dark green and white	Hyaline with dark green at the centre	Concentric zone with mix of green and white on first zone, white on second zone and green on third zone

Twenty four hours after inoculation, all isolates formed colonies with smooth colorless misellia and had colony diameter of 40 mm. At 48 hours, the misellia began to thicken as cotton and white misellia and colony size increased up to 60 mm. Colony showed a radial growth pattern with a concentric rings of green and white color. All isolates formed conidia, conidiophore, and phyalid. Both isolates from Maros and Takalar showed subglobose conidia shaped, slightly conidiophores branches formed, phyalid shaped of flask and ampulliform. Jeneponto isolate had subglobose form of conidia, conidiophores with many branch, phyalid shape of flask, ampulliform and lageniform (Table 2 and Figure 1c). At 96 hour after inoculation, the misellia growing thick and coarse grainy, and the white color slowly turn into green. The rapid grow of colonies was observed after 5 days where the diameter of the colony reached environ 90 mm. In this phase the missellia were abundant, thick and velvety and covered the plate (Figure 1a and 1b).

To evaluate the effect of *Trichoderma* isolates on seed germination and seedling growth, three isolates (Maros, Takalar and Jeneponto) were inoculated on maize seed. Inoculation was done by soaking with approximately  $1 \times 10^6$  spores per ml. One hundred maize seeds soaked in with 10 ml of *Trichoderma* spore suspension then planted in planting box and placed in green house. Percentage of seed germination was counted at 48, 72 and 96 h after incubation. 10 days after inoculation, plant response parameters such as shoot length, root length, dry root weight, dry shoot weight were measured in 10 plants after uprooting the plants and washing them under running tap water to remove residual soil from the roots.

**Statistical Analysis**


The experimental design used was a completely randomized design (CRD). Each isolate was repeated four times (400 seed/treatment), therefore there were 16 boxes of planting in total including controls. Means were analyzed by analysis of variance (ANOVA) and LSD test at 5% significant level.

**3. Results**

*Trichoderma* was found in all samples collected from Maize rhizosphere of Maros, Takalar, and Jeneponto regencies. The fungal isolates have colonies with thick and velvety texture and dark green color. The colonies of all origin grew covering the media. Upper and lower surface show a dark green color with a ring pattern evident in green (Table 1).

**Table 2:** Microscopical characteristics of *Trichoderma* indigenous strains grew on PDA medium isolated from rhizosphere of maize growing in dry land of Maros, Takalar and Jeneponto regencies


Observation	Isolates		
	Maros	Takalar	Jeneponto
<b>Conidia</b>			
Shape	Subglobose	Subglobose	Subglobose
Color	Green	Green	Green
Surface	Smooth	Smooth	Smooth
<b>Conidiophores</b>			
Surface	Smooth	Smooth	Smooth
Color	Hyaline	Hyaline	Hyaline
Branching	Few	Few	Plenty
<b>Phialide</b>			
Shape	Flask, ampulliform	Flask, ampulliform	Flask, ampulliform, lageniform
Color	Hyaline	Hyaline	Hyaline
Surface	Smooth	Smooth	Smooth




**Figure 1:** The colony of seven days *Trichoderma* isolates growing on PDA media (a); 'Reverse of colony' of seven


days old *Trichoderma* isolates growing on PDA media (b); Microscopic morphology of four days old *Trichoderma* isolates (400 x) (c).

No impact was observed within 48, 72 and 96 hours of incubation on maize seed germination after inoculation by *Trichoderma* of Maros, Takalar and Jeneponto isolates (Figure 2). While, these three isolates of *Trichoderma* treatment affect significantly root length, shoot length, dry root weight, and dry shoot weight of 10 days old seedling. The increase of four parameters above by Maros isolate compared to control was respectively 24.7%, 20.0%, 60.0%, and 57.1%, by Takalar isolates was respectively 16.7%, 17.4%, 60.0% and 42.9%, and by Jeneponto isolate was respectively 40.8%, 17.7%, 80.0%, and 42.9%. Root length treated by Jeneponto isolate was significantly different with that treated by Takalar isolate (Figure 3).



**Figure 2:** Percentage of maize seed germination at 48 hours, 72 hours, and 96 hours after treatment by *Trichoderma* of Maros, Takalar, and Jeneponto isolates through seed soaking. Means followed with the same letter were not significantly different ( $P \leq 0.05$ ) according to LSD test.





**Figure 3.** Root length (A), shoot length (B), dry root weight (C) and dry shoot weight (D) of ten days maize seedling after treatment by *Trichoderma* of Maros, Takalar, Jeneponto isolates through seed soaking. Means followed with the same letter were not significantly different ( $P \leq 0.05$ ) according to LSD test.

#### 4. Discussion

As *Trichoderma* has living habitat in agricultural lands and forests make the *Trichoderma* can be isolated from almost all agricultural land (Choi et al., 2003; Kredics et al., 2014). We also found *Trichoderma* isolates in all soil samples of maize rhizosphere from Maros, Takalar, and Jeneponto regencies. Cultural and morphological characteristic of Maros and Takalar isolates showed the similitude, while Jeneponto isolates indicated the difference. It mean apparently that isolates of Maros and Takalar have same species and isolate of Jeneponto classified in other species.

Maize seed germination was not affected by the application *Trichoderma* isolates. This finding is consistent with the results of Azarmi et al. (2011), Celar and Valic (2005) and Hajieghrari (2010).. This may be attributed to the utilization of existing food reserves in seeds such as carbohydrates, proteins, fats, amino acids and growth hormone to germinate. Seed germination is a process of morphological and biochemical changes that are influenced by the absorption of water, the chemical composition of seeds and phytohormones (Bewley and Black, 1985; Copeland and McDonald, 1995). Therefore, seed germination was not influenced by external factor including *Trichoderma* treatment.

All growth parameters of maize plant includes plant height, root length, shoot dry weight and root dry weight treated by three isolates of *Trichoderma* were significantly different compared to the control plants (Table 3). These increased growth response of plants depended mainly on the ability of *Trichoderma* to survive and develop in the rhizosphere (Harman et al., 2004). The rhizosphere is among the common ecological niches for *Trichoderma* spp. and provides opportunities for both biotrophy and saprotrophy on plant root exudates ( Druzhinina et al., 2011). The increasing root biomass by *Trichoderma*, may be due to auxine-mediated response pathways and also be mediated by a decrease in the level of the plant hormone ethylene (Wang et al., 2002; Contreras-Cornejo et al., 2009). In case of *T.asperelloides* T203, this species have  $\alpha$ -1-aminocyclopropane-1-carboxylate (ACC) deaminase gene (*acc1*) that encodes an enzyme which cleaves ACC that is expressed during the interaction with roots of

*Brassica napus*. ACC is a key intermediate in ethylene biosynthesis and *acc1* enzyme offers a mechanism for facilitating the formation of longer roots (Viterbo et al., 2010). In addition, the *Trichoderma* genomes contain many genes that encode nitrilase (Kubicek et al., 2011). These nitrilases may have a role either in hydrolysing  $\beta$ -cyano-L-alanine, a metabolite which is formed from cyanide released during the final step of ethylene biosynthesis, or in converting the plant metabolite indole-3-acetonitrile to indole-3-acetic acid (IAA), a hormone that promotes the growth of plant roots (Piotrowski and Volmer, 2006). The formation of longer roots by three isolates of *Trichoderma* permit apparently maize to have more capability in nutrient uptake, resulting in more shoot growth. Comparing to Maros and Takalar isolates, Jeneponto isolate enhance more formation of longer roots and therefore more potential for it use in improving maize vigor.

#### 5. Acknowledgement

The authors are heartily very thankful to plant disease laboratory Mr. Ardan, Kamaruddin, and Muh. Said Baco during this study.

#### References

- [1] Akladios, S. A and S. M. Abbas. 2012. Application of *Trichoderma harzianum* T22 as a biofertilizer supporting maize growth. African Journal of Biotechnology 11: 8672-8683
- [2] Azarmi, R., B. Hajieghrari, and A. Giglou. 2011. Effect of *Trichoderma* isolates on tomato seedling growth response and nutrient uptake. African Journal of Biotechnology 10 : 5850-5855
- [3] Barnett, H.L., and B.B. Hunter. 1978. Illustrated Genera of Imperfect Fungi, fourth ed. MacMillan Pub. Company, New York, USA, 217p.
- [4] Bewley, J.D. and M. Black. 1985. Seeds. Physiology of Development and Germination. Plenum Press, New York.
- [5] Celar, F. and N. Valic. 2005. Effects of *Trichoderma* spp. and *Gliocladium roseum* culture filtrates on seed germination of vegetables and maize. Journal of Plant Diseases and Protection 112: 343–350.



- [6] Choi, I. Y., S. B. Hong and M. C. Yadav. 2003. Molecular and Morphological Characterization of Green Mold, *Trichoderma* spp. isolated from Oyster Mushrooms. *Mycobiology* 31: 74-80.
- [7] Contreras-Cornejo HA, Macias-Rodriguez L, Cortes-Penagos C, Lopez-Bucio J. *Trichoderma virens*, a plant beneficial fungus, enhances biomass production and promotes lateral root growth through an auxin-dependent mechanism in *Arabidopsis*. *Plant Physiol* 2009;149:1579–1592.
- [8] Copeland, L.O. and M.B. McDonald. 1995. *Seeds Science and Technology*. Chapman and Hall, USA.
- [9] Druzhinina IS, Seidl-Seiboth V, Herrera-Estrella A, Horwitz BA, Kenerley CM, Monte E, Mukherjee PK, Zeilinger S, Grigoriev IV, Kubicek CP. *Trichoderma*: the genomics of opportunistic success. *Mycobiology* 2011;9:749-759.
- [10] Hajjeharari, B. 2010. Effects of some Iranian *Trichoderma* isolates on maize seed germination and seedling vigor. *African Journal of Biotechnology* 9 : 4342-4347.
- [11] Harman G.E. 2006. Overview of mechanisms and uses of *Trichoderma* spp. *Phytopathology* 96: 190-194.
- [12] Harman G.E., C.R. Howell, A. Vitrebo, I. Chet, and M. Lorito. 2004. *Trichoderma* species opportunistic, arivulent plant symbionts. *Nat. Rev. Microbiol.* 2: 43-56.
- [13] Hermosa R, Viterbo A, Chet I, Monte E. Plant-beneficial effects of *Trichoderma* and of its genes. *Microbiology* 2012; 158:17–25.
- [14] Kredics, L., L Hatvani, S. Naeimi, P. Kormoczi, L. Manczinger, C. Vagvolgyi, I. Druzhinina. 2014. Biodiversity of the Genus *Hypocrea/Trichoderma* in Different Habitats. *Biotechnology and Biology of Trichoderma*: 3-18
- [15] Mariola R. C., O. Rodriguez-Galan, T. Benitez, S. Sousa, M. Rey, A. Llobell, and J. Delgado-Jarana. 2007. Microscopic and Transcriptome Analyses of Early Colonization of Tomato Roots by *Trichoderma harzianum*. *International Microbiology* 10:19–27.
- [16] Nawrocka, J. and U. Małolepsza. 2013. Diversity in plant systemic resistance induced by *Trichoderma*. *Biological Control* 67:149-156.
- [17] Piotrowski M, Volmer JJ. Cyanide metabolism in higher plants: cyanoalanine hydratase is a NIT4 homolog. *Plant Mol. Biol* 2009;61:111–122.
- [18] Rahman, A., M. F. Begum, M. Rahman, M. A. Bari, G. N. M. Ilias, and M. F. Alam. 2011. Isolation and Identification of *Trichoderma* Species from Different Habitats and Their Use for Bioconversion of Solid Waste. *Turk J Biol* 35 :183-194.
- [19] Rosmana A, Samuels GJ, Ismaiel A, Ibrahim ES, Chaverri P, Herawati Y, Asman A. *Trichoderma asperellum*: A Dominant Endophyte Species in Cacao Grown in Sulawesi with Potential for Controlling Vascular Streak Dieback Disease. *Tropical Plant Pathology* 2015 March 03 [Epub]. <http://dx.doi.org/10.1007/s40858-015-0004-1>.
- [20] Samolski I, Rincon AM, Pinzon LM, Viterbo A, Monte E. The qid74 gene from *Trichoderma harzianum* has a role in root architecture and plant biofertilization. *Microbiology* 2012;158:129–138.
- [21] Schuster A, Schmoll M. Biology and biotechnology of *Trichoderma*. *Applied Microbiology and Biotechnology* 2010;87:787–799.
- [22] Shoresh, M., F. Mastouri and G. E. Harman. 2010. Induced systemic resistance and plant responses to fungal biocontrol agents. *Annu Rev Phytopathol* 48: 21–43.
- [23] Stewart, A. and R. Hill. 2014. Applications of *Trichoderma* in Plant Growth Promotion. *Biotechnology and Biology of Trichoderma* 4571: 415-425.
- [24] Viterbo A, Landau U, Kim S, Chernin L, Chet I. Characterization of ACC deaminase from the biocontrol and plant growth-promoting agent *Trichoderma asperellum* T203. *FEMS Microbiol. Lett* 2010;305:42–48.
- [25] Wang K, Li H, Ecker J. Ethylene biosynthesis and signaling networks. *Plant Cell* 2002;14:131-151.