

For the 38 items related to essential newborn care, each correct answer was awarded with a score of '1' and a score of '0' was awarded for the wrong answer. The data obtained was analyzed in terms of descriptive and inferential statistics. Data obtained were tabulated and analyzed in terms of objectives of the study using descriptive and inferential statistics.


3. Results

I) Finding on knowledge regarding essential newborn care among primipara mothers.

Table 1.a: Distribution of knowledge scores among primipara mothers about essential newborn care in study group, n=30

Knowledge scores	Pretest		Post test	
	Frequency	Percentage	Frequency	Percentage
Good (> 20) ($\bar{X} + SD$)	04	13.3%	30	100%
Average (15 – 20) ($\bar{X} - SD$) to ($\bar{X} + SD$)	21	70%	-	-
Poor (< 15) ($\bar{X} - SD$)	05	16.7%	-	-

Table 1.a Revealed that in pretest primipara mothers 21 (70%) had average knowledge and 5 (16.7%) had poor knowledge whereas 4 (13.3%) had good knowledge. In post test all primipara mothers 30 (100%) had good knowledge. This shows that there is gain in knowledge scores after administering video assisted teaching programme regarding essential newborn care.




Graph 1: Cylindrical graph showing percentage distribution of primipara mothers according to pretest and post test knowledge scores in study group

Table 1.b: Distribution of knowledge scores among primipara mothers regarding essential care in control group, n = 30

Knowledge scores	Pretest		Post test	
	Frequency	Percentage	Frequency	Percentage
Good (> 20) ($\bar{X} + SD$)	05	16.7%	10	33.3%
Average (15 – 20) ($\bar{X} - SD$) to ($\bar{X} + SD$)	20	66.6%	18	60%
Poor (< 15) ($\bar{X} - SD$)	05	16.7%	02	6.7%

Table 1.b Revealed that in pretest test primipara mothers were 20 (66.6%) had average knowledge and 5(16.7%) had good knowledge whereas 5 (16.7%) had poor knowledge. In post test 18 (60%) had average knowledge and 10 (33.3%) had good knowledge whereas only 2 (6.7%) had poor knowledge.



Graph 2: Cylindrical graph showing percentage distribution of primipara mothers according to pretest and post test knowledge scores in control group.

II: Pretest and post test mean percentage of knowledge scores in different areas of knowledge.

Table 2.a: Pretest and post test mean percentage knowledge scores of primipara mothers on different components of essential newborn care in study group, n=30

Area of knowledge	Total scores	Total pretest scores	Total post test scores	Pretest mean%	Post test mean%	Actual gain score %
Cleanliness/ Personal hygiene	330	149	265	45.15%	80.30%	35.15%
Thermal Protection	300	99	211	33%	70.33%	37.33%
Breastfeeding	330	153	240	46.36%	72.72%	26.36%
Immunization	180	89	180	49.44%	76.66%	27.22%

Table 2.a Revealed that there was 35.15% of actual gain in knowledge in cleanliness/ personal hygiene, thermal protection was 37.33%, breastfeeding was 26.36% and immunization was 27.22%.

Table 2.b: Pretest and post test mean percentage knowledge scores of primipara mothers on different components of essential newborn care in control group, n=30

Area of knowledge	Total scores	Total pretest scores	Total post test scores	Pretest mean%	Post test mean%	Actual gain score %
Cleanliness/ Personal hygiene	330	159	180	48.18%	54.54%	6.36%
Thermal Protection	300	85	119	28.33%	39.66%	11.33%
Breastfeeding	330	148	185	44.84%	56.06%	11.22%
Immunization	180	95	109	52.77%	60.55%	7.78%

Table 2.b Revealed that in cleanliness/ personal hygiene the actual gain of knowledge was 6.36%, thermal protection was 11.33%, breastfeeding was 11.22% and immunization was 7.78%.

Table 3 b: Comparing knowledge scores in study and control group, n (30+30)=60

Group	Pretest	Posttest	Increased Scores	Paired t test	Df
Study	16.33±3.54	28.47±2.69	12.13±1.80	37.0179	29
Control	17.20±3.17	19.67±2.95	2.47± 1.53	8.8578	29

III. Evaluate the effectiveness of video assisted teaching programme in terms of gain in post test knowledge scores in study group as compared to control group.

Table 3 a: Mean pretest scores, post test scores and standard deviation of both study and control group, and test of significance by using unpaired 't' test, n (30+30) =60

	Pretest Mean ±SD	Post test Mean±SD
Study group	16.33±3.54	28.47±2.69
Control group	17.20±3.17	19.67±2.95
Unpaired 't' value	1.00	12.0742
P value	0.3215	0.00001*

*(P<0.05) t (n₁+n₂-2) = 58

Table 3 a : Comparison of pretest scores between study and control group by unpaired't' test yielded p value > 0.05 suggesting of no significant difference. While after comparing post test scores between both the groups p value was obtained 0.00001 which was less than 0.05. This suggested that there was significant difference observed between the groups. Therefore it can be inferred that video assisted teaching programme regarding essential newborn care was effective to improve the knowledge of primipara mothers.

Table 3 b: Revealed that mean post test score of study group was significantly more than the control group. This proved that video assisted teaching programme was effective.

4. Discussion

1. Finding related to knowledge regarding essential newborn care among primipara mothers in study and control group.

In the present study findings showed that in study group in pretest primipara mothers 21 (70%) had average knowledge, 5 (16.7%) had poor knowledge whereas 4 (13.3%) had good knowledge. In post test all primipara mothers 30 (100%) had good knowledge. In control group primipara mothers 20 (66.6%) had average knowledge, 5(16.7%) had good knowledge whereas 5 (16.7%) had poor knowledge in pretest. In post test 18 (60%) had average knowledge and 10 (33.3%) had good knowledge whereas only 2 (6.7%) had poor knowledge. This study was supported by a study done by Shrishail B⁶. Finding showed that in pretest postnatal

mothers 39(65%) had average knowledge, 10(16.66%) had poor knowledge and 11(18.33%) had good knowledge in the pretest. In post test all the mothers 60(100%) had good knowledge score.

2. Findings related to effectiveness of video assisted teaching programme in terms of gain in post test knowledge scores in study group as compared to control group.

In the present study findings in study group after applying paired 't' test results showed that 't' value was 37.017 which was significant at $p < 0.05$ level. This revealed that video assisted teaching programme was effective in gaining knowledge regarding essential newborn care. The findings in the control group after applying paired 't' test results showed that 't' value was 8.857 which was significant at $p < 0.05$ level. This revealed that there was increase in knowledge scores in the post test scores as compared to pre test scores.

The similar findings was supported by the study conducted by M.D Santhi and Kokilavani⁶ that the obtained 't' value in the experimental group was 37.26, which was significant at $p < 0.05$ level and in the control group 't' value was 3.77 which was significant at $p < 0.05$.

The findings of the mean post test scores between study and control group by unpaired 't' test yielded $p < 0.05$ suggesting significant difference ($t = 12.0742$). Similar findings were seen in the study conducted by M.D Santhi and Kokilavani⁶. The findings showed that unpaired 't' value between experimental and control group post test knowledge scores was 58.85 which was significant at $p < 0.05$.

5. Conclusion

The study findings showed that there was a significant difference in the knowledge scores of the study group when compared to control group. The unpaired 't' test between post test knowledge score in study group and post test knowledge score in control group showed that there was significant gain in knowledge in study group compared to control group. The results of this study inferred that Video Assisted Teaching Programme (VATP) was the best teaching and learning strategy as it enhanced the knowledge of primipara mothers regarding essential newborn care.

6. Acknowledgment

A heartfelt appreciation goes to the my guide, Prof (Dr.) Sangeeta Kharde, HOD Dept of OBG (N), KLEU'S Institute Of Nursing Sciences, Belgaum and all the authority and staff nurses of KLE Dr. Prabhakar Kore Charitable Hospital, Belgaum.

References

- [1] Dhana K. Study on Immediate Newborn Care. StudyMode.com. 9 Jan 2013.
- [2] Fernandez G. Importance of a child in the family. Child development Institute: 4 Jan 2012.

- [3] Mother Newborn news. Community based management of newborn infections. July 2007 to Dec.2008; vol.3; No.1-2.
- [4] Park K. Text Book of Preventive and Social Medicine. 20th ed. Jabalpur: Banarsidas Bhanut publisher; 2008. p.423-25.
- [5] Shrishal B. Study to assess the effectiveness of planned teaching programme on knowledge regarding prevention of neonatal hypothermia among postnatal mothers. [PG Nursing thesis]. Belgaum: KLE University; 2013.
- [6] Santhi MD and Kokilavani. Effectiveness of video teaching programme on physical care of newborn among primigravida mothers. Reviews of Progress. 2013 June : 1(8); 1-7.

Author Profile

Ms. Oinam Mandakini Devi, M. Sc (N), Lecturer, Department of Obstetrics and Gynaecological Nursing, TripuraSundari College of Nursing, City off: 12, B.K. Road, Banamalipur, Agartala-799001, Tripura West, India.

Prof. (Dr.) Sangeeta Kharde, M. Sc (N) , Ph D and HOD, Department of Obstetrics and Gynaecological Nursing, K.L.E University's Institute of Nursing Sciences, Belgaum-590010, Karnataka, India.