N- Dimensional Plane Wave Solutions of Einstein's Field Equations in GR theory

I. S. Mohurley¹, R. K. Jumale², J. K. Jumale³

¹Department of Physics, Shri Dnyanesh Mahavidyalaya, Nawargaon, India

²Kaveri Nagar, Yavatmal Road Darwha, Distt. Yavatmal, (M.S.) India.

³ Department of Physics, R. S. Bidkar Arts, Comm. & Science College, Hinganghat, India

Abstract: In the present paper, we have studied $\left[\frac{1}{\sqrt{2}}(y+z)-t\right]$ - type and $\left[t\sqrt{2}/(y+z)\right]$ -type plane wave solutions of

Einstein's field equations in general theory of relativity in the case where the zero mass scalar field coupled with gravitational field and zero mass scalar field coupled with gravitational & electromagnetic field and established the existence of these two types of plane wave solutions in V_n . Furthermore we have considered the case of massive scalar field and shown that the non-existence of these two types of plane wave solutions in GR theory.

Keywords: Einstein's field equations in general relativity in case where the zero mass scalar field coupled with gravitational field and coupled with gravitational & electromagnetic field, *N*-dimensional space-time, etc.

1. Introduction

H.Takeno (1961) has obtained the non-flat plane wave solutions g_{ij} of the field equations $R_{ij} = 0$ and established the existence of (z-t)-type and (t/z)-type plane waves for purely gravitational case in four-dimensional empty region of space-time. On the lines of Takeno (1961), Thengane (2002) have also obtained the plane wave solutions g_{ij} of the field equations $R_{ij} = 0$ in purely gravitational case by reformulating Takeno's (1961) definition of plane wave as follows:

Definition A plane wave g_{ij} is a non-flat solution of field equations

 $R_{ij} = 0$, $[i, j = 1, 2, 3, \dots, (n-3), (n-2), (n-1), n]$ (1.1) in an empty region of space-time such that

 $g_{ij} = g_{ij}(Z), \ Z = Z(x^i),$

where $x^{i} = x^{1}, x^{2}, x^{3}, \dots, x^{n-3}, y, z, t$ (1.2) in some suitable co-ordinate system such that

$$g^{ij}Z_{,i}Z_{,j}=0, \ Z_{,i}=\frac{\partial Z}{\partial x^{i}},$$
 (1.3)

 $Z = Z(y, z, t), \ Z_{n-2} \neq 0, \ Z_{n-1} \neq 0, \ Z_{n} \neq 0.$ (1.4) In this definition, the signature convention adopted is $g_{rr} < 0, \ r = 1, 2, 3, \dots, (n-1)$

$$\begin{vmatrix} g_{rr} & g_{rs} \\ g_{sr} & g_{ss} \end{vmatrix} > 0, \begin{vmatrix} g_{rr} & g_{rs} & g_{rt} \\ g_{sr} & g_{ss} & g_{st} \\ g_{tr} & g_{ts} & g_{tt} \end{vmatrix} < 0 ,$$

$$\begin{vmatrix} g_{11} & g_{12} & \cdots & g_{1(n-1)} \\ g_{21} & g_{22} & \cdots & g_{2(n-1)} \\ \cdots & \cdots & \cdots & \cdots \\ g_{(n-1)1} & g_{(n-1)2} & \cdots & g_{(n-1)(n-1)} \end{vmatrix} < 0$$

when *n* is even

and
$$\begin{vmatrix} g_{11} & g_{12} & \cdots & g_{1(n-1)} \\ g_{21} & g_{22} & \cdots & g_{2(n-1)} \\ \cdots & \cdots & \cdots & \cdots \\ g_{(n-1)1} & g_{(n-1)2} & \cdots & g_{(n-1)(n-1)} \end{vmatrix} > 0$$

when *n* is odd, $g_{nn} > 0$ (1.5)

[not summed for r, s = 1, 2, 3, ..., (n-1)] and accordingly

 $g = \det(g_{ij}) < 0$ if *n* is even and $g = \det(g_{ij}) > 0$ if *n* is odd (1.6)

In the paper [3] by investigating the plane symmetric line elements

$$ds^{2} = \sum_{i=1}^{n-3} -A(dx^{i})^{2} - B[dy^{2} + dz^{2} - dt^{2}]$$
(1.7)
and
$$ds^{2} = \sum_{i=1}^{n-3} -A(dx^{i})^{2} - Z^{2}Bdy^{2} - Z^{2}Bdz^{2} + Bdt^{2}$$
(1.8)

we have obtained the relations of non-vanishing components of Ricci tensor as

$$P = 2R_{(n-2)(n-2)} = 2R_{(n-1)(n-1)} = R_{nn} = 2R_{(n-2)(n-1)} = -\sqrt{2}R_{(n-2)n} = -\sqrt{2}R_{(n-1)n},$$
(1.9)

Volume 4 Issue 5, May 2015 <u>www.ijsr.net</u> Licensed Under Creative Commons Attribution CC BY

1462

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2013): 6.14 | Impact Factor (2013): 4.438

and
$$P = \frac{(y+z)^2}{Z^2} R_{(n-2)(n-2)} = \frac{(y+z)^2}{Z^2} R_{(n-1)(n-1)} = \frac{(y+z)^2}{2} R_{nn} = \frac{(y+z)^2}{Z^2} R_{(n-2)(n-1)}$$

= $-\frac{(y+z)^2}{Z\sqrt{2}} R_{(n-2)n} = -\frac{(y+z)^2}{Z\sqrt{2}} R_{(n-1)n}$ (1.10)

for $\left[\frac{1}{\sqrt{2}}(y+z)-t\right]$ -type and $\left[t\sqrt{2}/(y+z)\right]$ -type plane waves respectively.

In the present paper, we investigate whether these two types of plane wave solutions exist in the case where zero mass scalar field coupled with gravitational field and the zero mass scalar field coupled with gravitational and electromagnetic field. Furthermore we consider the coupling of massive scalar field with gravitational field and the massive scalar field with gravitational and electromagnetic field in V_n to investigate the existence of these two types of plane wave solutions of Einstein's field equation

$$R_{ij} = (-8\pi)[T_{ij} - \frac{1}{2}g_{ij}T],$$

[*i*, *j* = 1,2,3,...,(*n*-3),(*n*-2),(*n*-1),*n*] (1.11)

where R_{ij} is the Ricci tensor of the space-time,

 T_{ii} is the energy momentum tensor,

 g_{ii} is the fundamental tensor of the space-time

and $T = T_i^i = g^{ij} T_{ij}$.

We study these two plane wave solutions separately.

Case I $\left[\frac{1}{\sqrt{2}}(y+z)-t\right]$ -type type plane wave solutions in V_n

2. Zero Mass Scalar Field Coupled with Gravitational Field in V_n

The energy momentum tensor of zero mass scalar field is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} V_k V^k],$$

[k = 1,2,3,...,(n-3),(n-2),(n-1),n] (2.1)

where *V* is scalar function of *Z* and $V_j = \frac{\partial V}{\partial x^j}$, $(x^j = x^1, x^2, x^3 \dots x^{n\cdot 3}, x^{n-2}, x^{n-1}, x^n \text{ i.e. } x^j = x^1, x^2, x^3 \dots x^{n\cdot 3}, y, z, t)$. Thus $\sqrt{2}V_{n-2} = \sqrt{2}V_{n-1} = -V_n = \overline{V}$ $\because V_1 = V_2 = V_3 = \dots = V_{n-3} = 0$ (2.2) where (-) bar denotes partial derivative w. r. to *Z* From line element (1.7) we have $g_{(n-2)(n-2)} = -B$, $g_{(n-1)(n-1)} = -B$,

$$g_{nn} = B, (2.3)$$

$$g^{(n-2)(n-2)} = -\frac{1}{B}, \qquad g^{(n-1)(n-1)} = -\frac{1}{B},$$

$$g^{nn} = \frac{1}{B} (2.4)$$

$$\Rightarrow V_k V^k = 0. (2.5)$$

Therefore equation (2.1) becomes

$$T_{ij} = \frac{1}{4\pi} [V_i V_j] \ (2.6)$$

And from equation (2.6), in zero mass scalar field

$$T = T_i^i = g^{ij} T_{ij} = 0 . (2.7)$$

Then using equations (2.6) and (2.7), Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j] \ (2.8)$$

which further gives

$$R_{(n-2)(n-2)} = -V,^{2}$$

$$R_{(n-1)(n-1)} = -\overline{V},^{2} R_{nn} = -2\overline{V},^{2} R_{(n-2)(n-1)} = -\overline{V},^{2}$$

$$R_{(n-2)n} = \sqrt{2} \overline{V},^{2} R_{(n-1)n} = \sqrt{2} \overline{V}^{2} (2.9)$$

Thus non-vanishing components of Ricci tensor are related as

$$2R_{(n-2)(n-2)} = 2R_{(n-1)(n-1)} = R_{nn} = 2R_{(n-2)(n-1)} = -\sqrt{2}R_{(n-2)n} = -\sqrt{2}R_{(n-1)n}.$$
 (2.10)

It is observed that the equation (2.10) is compatible with the equation (1.9) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $\left[\frac{1}{\sqrt{2}}(y+z)-t\right]$ -type plane wave solutions

exist in the case where zero mass scalar field is coupled with the gravitational field.

3. Zero Mass Scalar Field Coupled with Gravitational & Electromagnetic Field in V_n

In this case the energy momentum tensor is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} V_k V^k] + E_{ij}$$
(3.1)

where E_{ij} denotes electromagnetic energy momentum tensor.

Volume 4 Issue 5, May 2015

<u>www.ijsr.net</u>

Licensed Under Creative Commons Attribution CC BY

But as in the previous case for zero mass scalar field $V_k V^k = 0$ and $T = T_i^i = g^{ij}T_{ii} = 0$. (3.2)

Therefore equation (3.1) becomes

$$T_{ij} = \frac{1}{4\pi} [V_i V_j] + E_{ij} . (3.3)$$

Hence Einstein's field equation (1.11) becomes $R_{ij} = -2[V_iV_j + 4\pi E_{ij}]$. (3.4)

Then equation (3.4) yields

$$R_{(n-2)(n-2)} = -2[\frac{\overline{V}^{2}}{2} + 4\pi E_{(n-2)(n-2)}],$$

$$R_{(n-1)(n-1)} = -2[\frac{\overline{V}^{2}}{2} + 4\pi E_{(n-1)(n-1)}],$$

$$R_{nn} = -2[\overline{V}^{2} + 4\pi E_{nn}],$$

$$R_{(n-2)(n-1)} = -2[\frac{\overline{V}^{2}}{2} + 4\pi E_{(n-2)(n-1)}],$$

$$R_{(n-2)n} = -2[-\frac{\overline{V}^{2}}{\sqrt{2}} + 4\pi E_{(n-2)n}],$$

$$R_{(n-1)n} = -2[-\frac{\overline{V}^{2}}{\sqrt{2}} + 4\pi E_{(n-1)n}].$$
 (3.5)

All above non-vanishing components of Ricci tensor are related as

$$2R_{(n-2)(n-2)} = 2R_{(n-1)(n-1)} = R_{nn} = 2R_{(n-2)(n-1)} = -\sqrt{2}R_{(n-2)n} = -\sqrt{2}R_{(n-1)n}$$
(3.6)

It is observed that the equation (3.6) is compatible with equation (1.9) which is obtained in the case of purely gravitational field. Hence we have

Conclusion
$$\left[\frac{1}{\sqrt{2}}(y+z)-t\right]$$
-type type plane wave

solutions exists in the case where zero mass scalar field is coupled with gravitational & electromagnetic field.

4. Massive Scalar Field Coupled with Gravitational Field in V_n

The energy momentum tensor of massive scalar field is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} (V_k V^k - m^2 V^2)],$$

[k = 1,2,3,...,(n-3),(n-2),(n-1),n] (4.1)

where V is scalar function of Z and $V_j = \frac{OV}{\partial x^j}$,

 $(x^{j} = x^{1}, x^{2}, x^{3}...x^{n-3}, y, z, t)$ *m* is mass associated with the massive scalar field.

Thus $V_{n-2}\sqrt{2} = V_{n-1}\sqrt{2} = -V_n = \overline{V}$ $\therefore V_1 = V_2 = V_3 = \cdots = V_{n-3} = 0$ (4.2) where (-) bar denotes partial derivative w. r. to Z. From line element (1.7) we have

$$g_{(n-2)(n-2)} = -B, \qquad g_{(n-1)(n-1)} = -B \qquad g_{nn} = B \quad (4.3)$$
$$g^{(n-2)(n-2)} = -\frac{1}{B}, \qquad g^{(n-1)(n-1)} = -\frac{1}{B},$$
$$g^{nn} = \frac{1}{B} \quad (4.4)$$
$$\Rightarrow V_k V^k = 0. \quad (4.5)$$

Therefore equation (4.1) implies

$$T_{ij} = \frac{1}{4\pi} [V_i V_j + \frac{1}{2} g_{ij} m^2 V^2].$$
(4.6)

Equation (4.6) yields

$$T = T_i^{\ i} = g^{\ ij}T_{ij} = \frac{1}{4\pi} [\frac{n}{2}m^2V^2] . (4.7)$$

Using (4.6) and (4.7), Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j + (\frac{2-n}{4})g_{ij}m^2 V^2]$$
(4.8)

which further yields

$$\begin{aligned} R_{(n-2)(n-2)} &= -[\overline{V}^2 + (\frac{n-2}{2})Bm^2V^2],\\ R_{(n-1)(n-1)} &= -[\overline{V}^2 + (\frac{n-2}{2})Bm^2V^2],\\ R_{nn} &= -[2\overline{V}^2 - (\frac{n-2}{2})Bm^2V^2], R_{(n-2)(n-1)} = -\overline{V},^2\\ R_{(n-2)n} &= \overline{V}^2\sqrt{2}, R_{(n-1)n} = \overline{V}^2\sqrt{2}. (4.9) \end{aligned}$$

But from the line element (1.7) we have the relation of non-vanishing components of Ricci tensor as

$$2R_{(n-2)(n-2)} = 2R_{(n-1)(n-1)} = R_{nn} = 2R_{(n-2)(n-1)} = -\sqrt{2}R_{(n-2)n} = -\sqrt{2}R_{(n-1)n}$$
(4.10)

It is to be noted that here the equation (4.9) is incompatible with the equation (4.10) which is obtained in the case of purely gravitational field. Hence we have

Conclusion
$$\left[\frac{1}{\sqrt{2}}(y+z)-t\right]$$
-type plane wave solutions

doesn't exist in the case where massive scalar field is coupled with gravitational field.

Remark If $m^2 = 0$ then the equation (4.9) is compatible to (4.10) and we have a result as in the case where the zero mass scalar field is coupled with the gravitational field.

5. Massive Scalar Field Coupled with Gravitational & Electromagnetic Field in V_n

In this case the energy momentum tensor is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} (V_k V^k - m^2 V^2)] + E_{ij}$$
(5.1)

where E_{ij} denotes electromagnetic energy momentum tensor.

But as in the previous case for massive scalar field

Volume 4 Issue 5, May 2015

$$V_k V^k = 0$$
 and $T = T_i^i = g^{ij} T_{ij} = \frac{1}{4\pi} [\frac{n}{2} m^2 V^2]$. (5.2)

Therefore equation (5.1) becomes

$$T_{ij} = \frac{1}{4\pi} [V_i V_j + \frac{1}{2} g_{ij} m^2 V^2] + E_{ij} . (5.3)$$

Hence Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j + (\frac{2-n}{4})g_{ij}m^2 V^2 + 4\pi E_{ij}].$$
 (5.4)

Then from equation (5.4) we have

$$\begin{split} R_{(n-2)(n-2)} &= -[\overline{V}^2 + (\frac{n-2}{2})Bm^2V^2 + 8\pi E_{(n-2)(n-2)}],\\ R_{(n-1)(n-1)} &= -[\overline{V}^2 + (\frac{n-2}{2})Bm^2V^2 + 8\pi E_{(n-2)(n-2)}],\\ R_{nn} &= -[2\overline{V}^2 - (\frac{n-2}{2})Bm^2V^2 + 8\pi E_{nn}],\\ R_{(n-2)(n-1)} &= -[\overline{V}^2 + 8\pi E_{(n-2)(n-1)}],\\ R_{(n-2)n} &= -[-\sqrt{2}\overline{V}^2 + 8\pi E_{(n-2)n}],\\ R_{(n-1)n} &= -[-\sqrt{2}\overline{V}^2 + 8\pi E_{(n-1)n}]. (5.5) \end{split}$$

But line element (1.7) gives the relation of non-vanishing components of Ricci tensor as

$$2R_{(n-2)(n-2)} = 2R_{(n-1)(n-1)} = R_{nn} = 2R_{(n-2)(n-1)} = -R_{(n-2)n}\sqrt{2}$$
(5.6)

Here it has been observed that the equation (5.5) is incompatible with the equation (5.6) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $\left[\frac{1}{\sqrt{2}}(y+z)-t\right]$ -type plane wave solutions

of Einstein's field equation in general relativity doesn't exist in the case where massive scalar field is coupled with gravitational & electromagnetic field.

Remark If $m^2 = 0$ then the equation (5.5) is compatible to (5.6) and we have a result as in the case where the zero mass scalar field is coupled with the gravitational & electromagnetic field.

Case II we consider the case of $[t\sqrt{2}/(y+z)]$ -type plane wave solutions.

6. Zero Mass Scalar Field Coupled with Gravitational Field in V_n

The energy momentum tensor of zero mass scalar field is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} V_k V^k],$$

[k = 1,2,3,...,(n-3),(n-2),(n-1),n] (6.1)

where V is scalar function of Z and $V_j = \frac{\partial V}{\partial x^j}$, $(x^j = x^1, x^2, x^3 \dots x^{n-3}, y, z, t)$. Thus $V_1 = V_2 = V_3 = \dots, V_{n-3} = 0$ and $V_{(n-2)} = V_{(n-1)} = -V_n \frac{Z}{\sqrt{2}} = -\frac{\overline{VZ}}{(y+z)}$ (6.2) here (-) bar denotes partial derivative w. r. to Z From line element (1.8) we have $g_{(n-2)(n-2)} = -Z^2B$, $g_{(n-1)(n-1)} = -Z^2B$, $g_{nn} = B$,

(6.3)

$$g^{(n-2)(n-2)} = -\frac{1}{Z^2 B}, \qquad g^{(n-1)(n-1)} = -\frac{1}{Z^2 B}, \qquad g^{nn} = \frac{1}{B}$$

(6.4)

$$\Rightarrow V_k V^k = 0. (6.5)$$

Therefore equation (6.1)
$$\Rightarrow T_{ij} = \frac{1}{4\pi} [V_i V_j] . (6.6)$$

Equation (6.6) yields

$$T = T_i^i = g^{ij}T_{ij} = 0.$$
 (6.7)

Then from (6.6) and (6.7), Einstein's field equation (1.11) becomes $\sqrt{2} = -R_{(n-1)n}\sqrt{2}$

$$R_{ij} = -2[V_i V_j]$$
 (6.8)
which further gives

$$\begin{aligned} R_{(n-2)(n-2)} &= -2\left[\frac{\overline{V}^2 Z^2}{(y+z)^2}\right],\\ R_{(n-1)(n-1)} &= -2\left[\frac{\overline{V}^2 Z^2}{(y+z)^2}\right], R_{nn} = -2\left[\frac{2\overline{V}^2}{(y+z)^2}\right]\\ R_{(n-2)(n-1)} &= -2\left[\frac{\overline{V}^2 Z^2}{(y+z)^2}\right],\\ R_{(n-2)n} &= 2\sqrt{2}\left[\frac{\overline{V}^2 Z}{(y+z)^2}\right], R_{(n-1)n} = 2\sqrt{2}\left[\frac{\overline{V}^2 Z}{(y+z)^2}\right].\end{aligned}$$

(6.9)

Thus non-vanishing components of Ricci tensor are related as

$$\frac{(y+z)^2}{Z^2} R_{(n-2)(n-2)} = \frac{(y+z)^2}{Z^2} R_{(n-1)(n-1)} = \frac{(y+z)^2}{2} R_{nn} = \frac{(y+z)^2}{Z^2} R_{(n-2)(n-1)}$$
$$= -\frac{(y+z)^2}{Z\sqrt{2}} R_{(n-2)n} = -\frac{(y+z)^2}{Z\sqrt{2}} R_{(n-1)n}.$$
(6.10)

It is observed that the equation (6.10) is compatible with equation (1.10) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $[t\sqrt{2}/(y+z)]$ -type plane wave solutions exist in the case where zero mass scalar field is coupled with the gravitational field.

Volume 4 Issue 5, May 2015 www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

7. Zero Mass scalar field coupled with gravitational & electromagnetic field in V_n

In this case the energy momentum tensor is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} V_k V^k] + E_{ij}$$
(7.1)

where E_{ij} denotes electromagnetic energy momentum tensor.

But as in the previous case for zero mass scalar field $V_k V^k = 0$ and $T = T_i^i = g^{ij}T_{ii} = 0$. (7.2)

Therefore equation (7.1) becomes

$$T_{ij} = \frac{1}{4\pi} [V_i V_j] + E_{ij} . (7.3)$$

Hence Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j + 4\pi E_{ij}] \quad (7.4)$$

which further yields

$$R_{(n-2)(n-2)} = -2\left[\frac{\overline{V}^{2}Z^{2}}{(y+z)^{2}} + 4\pi E_{(n-2)(n-2)}\right],$$

$$R_{(n-1)(n-1)} = -2\left[\frac{\overline{V}^{2}Z^{2}}{(y+z)^{2}} + 4\pi E_{(n-1)(n-1)}\right],$$

$$R_{nn} = -2\left[\frac{2\overline{V}^{2}}{(y+z)^{2}} + 4\pi E_{nn}\right],$$

$$R_{(n-2)(n-1)} = -2\left[\frac{\overline{V}^{2}Z^{2}}{(y+z)^{2}} + 4\pi E_{(n-2)(n-1)}\right],$$

$$R_{(n-2)n} = -2\left[\frac{-\overline{V}^{2}Z\sqrt{2}}{(y+z)^{2}} + 4\pi E_{(n-2)n}\right],$$

$$R_{(n-1)n} = -2\left[\frac{-\overline{V}^{2}Z\sqrt{2}}{(y+z)^{2}} + 4\pi E_{(n-1)n}\right].$$
(7.5)

All above non-vanishing components of Ricci tensor are related as

$$\frac{(y+z)^2}{Z^2} R_{(n-2)(n-2)} = \frac{(y+z)^2}{Z^2} R_{(n-1)(n-1)} = \frac{(y+z)^2}{2} R_{nn}$$
$$= -\frac{(y+z)^2}{Z\sqrt{2}} R_{(n-2)n} = -\frac{(y+z)^2}{Z\sqrt{2}} R_{(n-1)n}.$$
(7.6)

It is observed that the equation (7.6) is compatible with equation (1.10) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $[t\sqrt{2}/(y+z)]$ -type plane wave solutions exist in the case where zero mass scalar field is coupled with gravitational & electromagnetic field.

8. Massive scalar field coupled with gravitational field in V_n

The energy momentum tensor of massive scalar field is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} (V_k V^k - m^2 V^2)],$$

[k = 1,2,3,...,(n-3),(n-2),(n-1),n] (8.1)
where V is scalar function of Z and $V_j = \frac{\partial V}{\partial x^j},$
(x^j = x¹, x², x³...xⁿ⁻³, y, z, t),

m is mass associated with the massive scalar field.

Thus
$$V_{(n-2)} = V_{(n-1)} = -V_n \frac{Z}{\sqrt{2}} = -\frac{VZ}{(y+z)}$$
,
 $\therefore V_1 = V_2 = V_3 = \cdots, V_{n-3} = 0$ (8.2)

where (-) bar denotes partial derivative w. r. to Z. From line element (1.8) we have

$$g_{(n-2)(n-2)} = -Z^{2}B , \quad g_{(n-1)(n-1)} = -Z^{2}B , \quad g_{nn} = B ,$$

$$g^{(n-2)(n-2)} = -\frac{1}{Z^{2}B}, \quad g^{(n-1)(n-1)} = -\frac{1}{Z^{2}B}, \quad g^{nn} = \frac{1}{B}$$
(8.3)
(8.3)
(8.4)

 $\Rightarrow V_k V^k = 0. (8.5)$

Therefore equation (8.1) becomes

$$T_{ij} = \frac{1}{4\pi} [V_i V_j + \frac{1}{2} g_{ij} m^2 V^2].$$
(8.6)

From equation (8.6) we have

$$T = T_i^{\ i} = g^{\ ij} T_{ij} = \frac{1}{4\pi} [\frac{n}{2} m^2 V^2]. \ (8.7)$$

Then from (8.6) and (8.7), Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j + (\frac{2-n}{4})g_{ij}m^2 V^2]$$
(8.8)

which further yields

$$\begin{split} R_{(n-2)(n-2)} &= -2[\frac{\overline{V}^2 Z^2}{(y+z)^2} + (\frac{n-2}{4})Z^2 Bm^2 V^2], \\ &= R_{(n-\overline{Z})^{n-1}}^{(y+z)^2} \frac{R_{(n}2[\frac{\overline{V}^2 Z^2}{(y+z)^2} + \frac{Z^2 Bm^2 V^2}{2}], \\ R_{nn} &= -2[\frac{2\overline{V}^2}{(y+z)^2} - (\frac{n-2}{4})Bm^2 V^2], \\ R_{(n-2)(n-1)} &= -2[\frac{\overline{V}^2 Z^2}{(y+z)^2}] \\ R_{(n-2)n} &= -2[\frac{-\overline{V}^2 Z \sqrt{2}}{(y+z)^2}], R_{(n-1)n} = -2[\frac{-\overline{V}^2 Z \sqrt{2}}{(y+z)^2}] \end{split}$$

(8.9)

But from the line element (1.8) we have the relation of non-vanishing components of Ricci tensor as

<u>www.ijsr.net</u> Licensed Under Creative Commons Attribution CC BY

Volume 4 Issue 5, May 2015

$$\frac{(y+z)^2}{Z^2} R_{(n-2)(n-2)} = \frac{(y+z)^2}{Z^2} R_{(n-1)(n-1)} = \frac{(y+z)^2}{2} R_{nn} = \frac{(y+z)^2}{Z^2} R_{(n-2)(n-1)}$$
$$= \frac{-(y+z)^2}{Z\sqrt{2}} R_{(n-2)n} = \frac{-(y+z)^2}{Z\sqrt{2}} R_{(n-1)n}$$
(8.10)

Here we observed that the equation (8.9) is incompatible with the equation (8.10) which is obtained in the case of purely gravitational field. Hence we have

2

Conclusion $[t\sqrt{2}/(y+z)]$ -type plane wave solutions of Einstein's field equation in general theory of relativity doesn't exist in the case where massive scalar field is coupled with the gravitational field.

Remark If $m^2 = 0$ then the equation (8.9) is compatible to (8.10) and we have a result as in the case where the zero mass scalar field is coupled with the gravitational field.

9. Massive scalar field coupled with gravitational & electromagnetic field in V_n

In this case the energy momentum tensor is given by

$$T_{ij} = \frac{1}{4\pi} [V_i V_j - \frac{1}{2} g_{ij} (V_k V^k - m^2 V^2)] + E_{ij}$$
(9.1)

where E_{ij} denotes electromagnetic energy momentum tensor.

But as in the previous case for massive scalar field

$$V_k V^k = 0$$
 and $T = T_i^i = g^{ij} T_{ij} = \frac{1}{4\pi} [\frac{n}{2} m^2 V^2].$ (9.2)

Therefore equation (9.1) becomes

$$\frac{(y+z)^2}{Z^2} R_{(n-2)(n-2)} = \frac{(y+z)^2}{Z^2} R_{(n-1)(n-1)} = \frac{(y+z)^2}{2} R_{nn} = \frac{(y+z)^2}{Z^2} R_{(n-2)(n-1)}$$
$$-(y+z)^2 = -(y+z)^2$$

$$=\frac{-(y+z)^2}{Z\sqrt{2}}R_{(n-2)n}=\frac{-(y+z)^2}{Z\sqrt{2}}R_{(n-1)n}.$$
 (9.6)

Here we observed that the equation (9.5) is incompatible with the equation (9.6) which is obtained in the case of purely gravitational field. Hence we have

Conclusion $[t\sqrt{2}/(y+z)]$ -type plane wave solutions of Einstein's field equation doesn't exist in the case where massive scalar field is coupled with gravitational and electromagnetic field.

Remark If $m^2 = 0$ then the equation (9.5) is compatible to (9.6) and we have a result as in the case where the zero mass scalar field is coupled with the gravitational and electromagnetic field.

References

$$T_{ij} = \frac{1}{4\pi} [V_i V_j + \frac{1}{2} g_{ij} m^2 V^2] + E_{ij} .$$
(9.3)

Hence Einstein's field equation (1.11) becomes

$$R_{ij} = -2[V_i V_j + (\frac{2-n}{4})g_{ij}m^2 V^2 + 4\pi E_{ij}].$$
(9.4)

And from equation (9.4) we have

$$\begin{split} R_{(n-2)(n-2)} &= -2\left[\frac{VZ^2}{(y+z)^2} + (\frac{n-2}{4})Z^2Bm^2V^2 + 8\pi E_{(n-2)(n-2)}\right],\\ R_{(n-1)(n-1)} &= -2\left[\frac{\overline{V}^2Z^2}{(y+z)^2} + (\frac{n-2}{4})Z^2Bm^2V^2 + 8\pi E_{(n-1)(n-1)}\right]\\ R_{nn} &= -2\left[\frac{2\overline{V}^2}{(y+z)^2} - (\frac{n-2}{4})Bm^2V^2 + 8\pi E_{nn}\right],\\ R_{(n-2)(n-1)} &= -2\left[\frac{\overline{V}^2Z^2}{(y+z)^2} + 8\pi E_{(n-2)(n-1)}\right],\\ R_{(n-2)n} &= -2\left[\frac{-\overline{V}^2Z\sqrt{2}}{(y+z)^2} + 8\pi E_{(n-2)n}\right],\\ R_{(n-1)n} &= -2\left[-\frac{\overline{V}^2Z\sqrt{2}}{(y+z)^2} + 8\pi E_{(n-1)n}\right]. (9.5) \end{split}$$

But line element (1.8) gives the relation of non-vanishing components of Ricci tensor as

[1] B. G. Ambatkar (2015) : $\left[\frac{1}{\sqrt{2}}(y+z)-t\right]$ -type and

 $[t\sqrt{2}/(y+z)]$ -type four and J K Jumale dimensional plane wave solutions of Einstein's field equations in general relativity. *International Journal of Engineering* and Science Vol.5, Issue 2, Feb. 2015, pp. 30-38.

[2] B. G. Ambatkar (2015) : $\left[\frac{1}{\sqrt{2}}(y+z)-t\right]$ -type and

 $[t\sqrt{2}/(y+z)]$ -type plane wave and J K Jumale solutions of Einstein's field equations in V_6 . Online International Interdisciplinary Research. Vol.V.Issue III, May- June.2015

[3] Thengane K D (2003) : Solutions of field equations

$$R_{ij} = 0$$
 for $\left[\frac{1}{\sqrt{2}}(y+z) - t\right]$ -type and

Volume 4 Issue 5, May 2015

<u>www.ijsr.net</u>

Licensed Under Creative Commons Attribution CC BY

 $\left[t\sqrt{2}/(y+z)\right]$ -type plane waves in N-dimensional space- time (I).

[4] H Takeno(1961) : The Mathematical theory of plane gravitational wave in general relativity". Scientific report of the Research Institute to theoretical physics. Hiroshima, University, Takchara, Hiroshima-ken, Japan-1961