

Leaf- fall initiation	4	-----	5	49	-----	-----	-----	50
Flower - initiation	11	6	10	7	7	9	10	20
Flower -full initiation	13	7	11	7	9	10	13	21
Flower-end	20	17	15	18	28	24	17	38
Fruit - initiation	12	7	11	7	9	10	12	22
Fruit- mature	25	26	48	17	26	26	24	7
Fruit- dehiscence	-----	----	-----	19	27	27	-----	-----
Fruit-drop	26	27	-----	-----	28	-----	25	-----

Table 7: Range and average of germination percentages for saplings and mature tree species during 2014/2015 season in Aswan

Tree species	Germination % for saplings		Germination % for mature trees	
	Range	Average	Range	Average
<i>Azadirachta indica</i>	51.23- 63.17	54.27	38.47- 56.23	40.25
<i>Chrysophyllum oliviforme</i>	55.47- 70.35	60.55	51.33- 66.17	56.23
<i>Dalbergia sissoo</i>	85.55- 93.36	90.23	81.32- 88.33	85.23
<i>Inga edulis</i>	51.34- 71.33	65.17	36.55- 65.15	55.18
<i>Khaya senegalensis</i>	95.45- 100	98.14	94.23- 99.11	96.17
<i>Spathodea campanulata</i>	33.17- 51.27	45.33	24.18- 45.47	40.56
<i>Syzygium cuminii</i>	96.47- 100	99.15	85.33- 97.23	94.17
<i>Tamarindus indica</i>	65.33- 81.17	71.33	45.45- 74.55	65.45

References

- [1] **Arbonnier, M., 2004.** Trees, Shrubs and Lianas of West Africa Dry Zones. CIRAD, Margraf Pub. GMBH: Germany, 292-293.
- [2] **Augsburger, C.K., Bartlett, E.A., 2003.** Differences in leaf phenology between juvenile and adult trees in a temperate deciduous forest. *Tree Physiology*. 23: 517-525.
- [3] **Badeck, F.W., Bondeau, A., Bottcher, K., Doktor, D., Lucht, W., Schaber, J., Sitch, S., 2004.** Responses of spring phenology to climate change. *New Phytol* 162:295-309.
- [4] **Bangarwa, K.S., Singh, V.P., 1994.** Floral Biology and crossing techniques in *Dalbergia sissoo* Roxb In *Dalbergia*. Proc. of an International workshop. Ed. Westley SB and Roshrtko JM. Publ. Nitrogen fixing tree association, USA.
- [5] **Borchert, R., 1994.** Soil and stem water storage determine phenology and distribution of tropical dry forest trees. *Ecology*. 75:1437-1449.
- [6] **Borchert, R., Rivera, G., 2001.** Photoperiodic control of seasonal development and dormancy in tropical stem-succulent trees. *Tree Physiology* 2: 213-221.
- [7] **Burt, B.L., 1970.** The evolution and taxonomic significance of a subterranean ovary in certain monocotyledons. *Israel Journal of Botany* 19: 77-90.
- [8] **Cayan, D.R., Kammerdiener, S.A., Dettinger, M.D., Caprio, J.M., Peterson, D.H., 2001.** Changes in the onset of spring in the western United States. *Bull Am Met Soc* 82:399-415
- [9] **Dewald, L.E., Steiner, K.C., 1986.** Phenology, height increment and cold tolerance of *Alnus glutinosa* population in a common environment. *Silvae Genetica* 35: 205-211.
- [10] **El-Siddig, K., Ebert, G., Ludders, P., 1999.** Tamarind (*Tamarindus indica*): a review on a multipurpose tree with promising future in the Sudan. *J. Applied Botany* 73: 202-205.
- [11] **Facao, M.A., Clement, C.R., 2000.** Phenology and productivity of *Inga edulis* in central Amazonia. *J. Acta Amazonica*, 30 (2): 173-180.
- [12] **F.A.O., 1982.** Fruit-bearing forest trees: technical notes. FAO- Forestry- Paper, No. 34, pp.177
- [13] **Gill, D.S., Amthor, J.S., Bormann, F.H., 1998.** Leaf phenology, photosynthesis, and the persistence of saplings and shrubs in a mature northern hardwood forest. *Tree Physiol*. 18:281-289.
- [14] **Harrington, R.A., Brown, B.J., Reich, P.B., 1989.** Ecophysiology of exotic and native shrubs in southern Wisconsin (USA). I. Relation-ship of leaf characteristics, resource availability, and phenology to seasonal patterns of carbon gain. *Oecologia* 80:356-367.
- [15] **Karmer, K., 1997.** Phenology and growth of European trees in relation to climate change. pp. 39-50.
- [16] **Khamis, M.H., 2013.** Phenological Variations of Three Urban Forest Trees Grown in North and Upper Egypt. *Alex. J. Agric. Res.* 58 (2): 71-81.
- [17] **Kikim, A., Yadava, P.S., 2001.** Phenology of tree species in subtropical forests of Manipur in north eastern India. *Tropical Ecology* 42: 269-276.
- [18] **Lei, T.T., Lechowicz, M.J., 1990.** Shade adaptation and shade tolerance in saplings of three Acer species from eastern North America. *Oecologia* 84:224-228.
- [19] **Little, E.L., Wadsworth, F.L., 1964.** Common trees of Puerto Rico and the Virgin Islands. *Agric. Handb.* 249, U.S. Dept. Agric., Washington, DC. p.548
- [20] **Murali, K.S., Sukumar, R., 1994.** Reproductive phenology of a tropical dry forest in Mudumala, Southern India. *Journal of Ecology* 82: 759-767.
- [21] **Oliveira, P.E., 1998.** Fenologi'a e biologia reproductiva das espe'cies de cerrado.
- [22] **In S.M. Sana and S.P. de Almeida [ed], Cerrado, ambiente e flora, 169-192.** Embrapa, Penaltina, DF, Brasil.
- [23] **Opler, P.A., Frankie, G.W., Baker, H.G., 1980.** Comparative phenological studies of treelet and shrubs species in tropical wet and dry forests in the lowlands of Costa Rica. *Journal of Ecology* 68:167-188.
- [24] **Petanidou, T., W.N. Ellis, Margaris, N.S., Vokou, D., 1995.** Con842 straints of flowering phenology in a

- phryganic (East Mediterranean shrub) community. *American Journal of Botany* 82: 607-620.
- [25] **Prasad, R., 1983.** Neem-the miracle tree for meeting Indian's growing energy needs. *Bioenergy News*, 1:43-47.
- [26] **Prasad, S.N., M., Hedge, 1986.** Phenology and seasonality in the tropical deciduous forest of Bandipur, South India. *Proceedings of Indian Academy of Sciences (Plant Sciences)* 96: 121-133.
- [27] **Rangaiah, K., S.P., Rao, A.J.S., Raju, 2004.** Bird-pollination and fruiting phenology in *Spathodea campanulata* Beauv. (Bignoniaceae). *J. Beitrage zur Biologie der Pflanzen*, 73 (3): 395-408.
- [28] **Rivera, G., Elliott, S., Caldas, L.S., Nicolossi, G., Coradin, V.T.R., Borchert, R., 2002.** Increasing day length induces spring flushing of tropical dry forest trees in the absence of rain. *Trees* 16: 445-456.
- [29] **Seiwa, K., 1998.** Advantages of early germination for growth and survival of seedlings of *Acer monouder* different overstorey phenologies in deciduous broad-leaved forests. *J. Ecol.* 86:219-228.
- [30] **Seiwa, K., 1999a.** Changes in leaf phenology are dependent on tree height in *Acer mono*, a deciduous broad-leaved tree. *Ann. Bot.* 83: 355-361.
- [31] **Seiwa, K., 1999b.** Ontogenetic changes in leaf phenology of *Ulmus davidiana* var. *japonica*, a deciduous broad-leaved tree. *Tree Physiol.* 19:793-797.
- [32] **Singh, J.S., Singh, V.K., 1992.** Phenology of seasonally dry tropical forest. *Current Science* 63: 684 - 688.
- [33] **Singh, K.P., Kushwaha, C.P., 2005a.** Emerging paradigms of tree phenology in dry tropics. *Current Science* 89: 964-975.
- [34] **Singh, K.P., Kushwaha, C.P., 2006.** Diversity of flowering and fruiting phenology of trees in a tropical deciduous forest in India. *Annals of Botany* 97: 265-276.
- [35] **Swenson, U.I.F., Richardson, J.E., Bartish, I.V., 2008.** "Multi-gene phylogeny of the pantropical subfamily Chrysophylloideae (Sapotaceae): Evidence of generic polyphyly and extensive morphological homoplasy". *Cladistics* 24 (6):1006-1031.
- [36] **Tutin, C.E.G., Fernandez, M., 1993.** Relationships between minimum temperature and fruit production in some tropical forest trees in Gabon. *Journal of Tropical Ecology* 9: 241-248.
- [37] **Wright, S.J., 1991.** Seasonal drought and the phenology of understory shrubs in a tropical moist forest. *Ecology* 72:1643-1652
- [38] **Yadav, R.K., Yadav, A.S., 2008.** Phenology of selected woody species in a tropical dry deciduous forest in Rajasthan, India. *Tropical Ecology* 49(1):25-34.
- [39] **Zhang, G., Song, Q., Yang, D., 2006.** Phenology of *Ficus racemosa* in Xishuangbanna, Southwest China. *Biotropica* 38: 334-341.