


- and its evaluation in broiler diets,” *Animal Feed Science and Technology*, 141, 129–140, 2008.
- [37] Ramasamy Ramasubburayan, Palanisamy Iyapparaj, Kanaharaja Joselin Subhashini, Manohar Navin Chandran, Arunachalam Palavesam, Grasian Immanuel, “Characterization and Nutritional Quality of Formic Acid Silage Developed from Marine Fishery Waste and their Potential Utilization as Feed Stuff for Common Carp *Cyprinus carpio* Fingerlings,” *Turkish Journal of Fisheries and Aquatic Science*, 13, 281-289, 2013.
- [38] K.B. Rao, M.S. Mithyantha, L.S. Devi, N.G. Perur, “Nutrient content of some new ragi varieties,” *Mysore J. Agric. Sci.*, 7, 4, 562-565, 1973.
- [39] P.U. Rao, “Evaluation of protein quality of brown and white ragi (*Eleusine coracana*) before and after malting,” *Food Chemistry* 51, 433- 436, 1994.
- [40] R. Sridhar, G. Lakshminarayana, “Contents of total lipids and lipid classes and composition of fatty acids in small millets: Foxtail (*Setaria italica*), Proso (*Panicum Miliceum*) and Finger Millet (*Eleusine coracana*),” *Cereal Chemistry*, 71, 4, 355-359, 1994.
- [41] B.S. Gunashree, R. Selva Kumar, R. Roobini G. Venkateswaran, “Nutrients and antinutrients of ragi and wheat as influenced by traditional processes,” *Int.J.Curr.Microbiol.App.Sci.*, 3, 7, 720-736, 2014.

Author Profile

Prof. G. H. Philip, Head and BOS, Department of Zoology, Sri Krishnadevaraya University, Anantapuramu, AP, India.

Mr. R. Sreedhar, Research Scholar, Department of Zoology, Sri Krishnadevaraya University, Anantapuramu, AP, India






Figure 2: Concentration of nutrients in fish waste and in plant material after addition of different concentration of waste of *Catla catla*