















- [15] ZoricalekaGrujiS. , SvetlanalukiI.,SkubanS.and TrifunoviS.2004. The Synthesis and Characterization of Complexes of Zinc (II) , Cadmium (II) , Platinum (II) and Palladium (II) with Potassium 3-Dithiocarboxy-3-Aza-5-Aminopentanoate.J.Serb. Chem. Soc..69(2):173-143.
- [16] FiggisB. and Hitchman M. 2000. Ligand Field Theory and ItsApplications. Wiley, New York.
- [17] CarolinSh. and Mahasin F. 2012.Synthesis,Spectroscopic Studyof Pt(IV),Au (III) ,Rh(III),Co(II) and V(IV) Complexes with Sodium {5-(P-NitroPhenyl)-4 -Phenyl-1,2,4-Triazole-3-Dithiocarbamato Hydrazide}and CytotoxicityAssay on Rhabdo-omyosarcoma CellLineof Heavy Metals. J. Baghdad for Sci..9(4):668-679 .
- [18] Young, D. C. (2001). "Computational Chemistry", John Wiley & Sons.
- [19] <sup>1</sup>L.J.V.Piddock; J.Applied Bacteriology; 68; 307; (1990).
- [20] Elzahany, E.A.,Hegab, K.H; Khalil, S.K.H. and Youssef, N.S. Synthesis, Characterization and Biological activity of some transition metal complexes with Schiff Bases Derived from 2- Formylidole, Salicyladehyde and N-amino Rhodanine. Aust. J Basic Appl. Scien. 2008. Vol.2(2) pp: 210-220.
- [21] Chandra, S.; Jain D. ; Sharma. A.K. and Sharma, p. coordination modes of Schiff base pentadentate derivative of u. amino anti- pyrine with cobalt (II) metal ions : synthesis, spectroscopic and Antimicrobial studies, Molecules 2009. Vol. 14(1). Pp174-190 .
- [22] IsraaQaisYousif and Mahasin Faisal Alias, synthesis, structural study, Antimicrobial activity and theoretical treatment of 3d-metal complexes involving Schiff base of piperonal derivative AJPS, 2013, vol. 13. No1, pp 1-13.