

Association, held by P.A Leggat and D.R Smith (12) among 285 dentists. 57.5% of the subjects reported cervix pain; 53.7% - lower back pain; 50.3% - shoulder pain; and 24.6% - neck pain.

A study, conducted in Hong Kong among 500 dentists (13) by T.K Lun Li et al. in 2002 established that the most common health problem for the dentists is MSD (43%), followed by allergic dermatitis of the hands (24%) and instrument caused injuries (17%). The prevalent pain was mostly in the lower back and the cervix.

R.S Acharya et al. studied 150 dentists from Nepal (2). The distribution of the musculoskeletal disorders for the last 12 months was as followed: neck pain – 52.4%; lower back pain -52.4%; shoulder pain- 49.5%; upper back – 41.7%; wrist/hand – 36.9%.

A phone interview held by E. Milerad and L. Ekenvall (15) showed that 44% of the dentists and 29% of the pharmacists reported neck pain; while 51% of the dentists and 23% of the pharmacists experienced shoulder pain. Musculoskeletal symptoms of the forearm were present predominantly with the dentists - 12% against just 1% for the pharmacists.

In Thailand S. Chowanadisai, B. Kukiattrakoon et al. (7) surveyed 220 Thailand dentists in 1997. 78% of the respondents reported musculoskeletal disorders, followed by percutaneous injuries - 50%.

A. Al-Kholani et al. (4) studied 136 dentists in a public dental care clinic in Yemen and found out that back pain affected 80% of the subjects, followed by neck pain (58.8%) and shoulder pain (47%). There results are close to those most of the studies. Another study, conducted by P.Dajpratham (8) involving 164 dentists displayed extremely high MSD incidence – 96.3%. Most widely spread was shoulder pain (72.2%), next came neck pain (70.3%) and lower back pain (50.6%). Hayes et al. (10) found out that musculoskeletal disorder-related pain among dental professionals (dentists, Students and hygienists) varied between 64% and 93%. Most common proved to be neck pain (64,29%), followed by lower back pain (57,94%) and shoulder pain - 48,41% in dental students, while these figures were respectively 36,3% - 60,1% for back pain and 60%-69,5% for neck pain.

The aim of this study is to establish the distribution of musculoskeletal disorders among the dentists in Bulgaria.

The study objectives cover:

1. Determining the incidence of pain, as a major MSD symptom among the dentists.
2. Outlining the pain character, localization and intensity.

2. Material and Methods

The source information needed for the study objectives was gathered through surveys using written questionnaires distributed and filled out within the period October 2012 – April 2013. The questionnaire was designed by us specifically to serve the objectives of the study. The survey was conducted with the support of BDU and regional bodies in Sofia, Pleven, Shoumen, Varna, and Vratza. A total of 1300 questionnaires were prepared and disseminated, of which we got back 700 (return rate – 53.84%). They comprised of 27 close questions (fixed answers) on paper, each having a unique entry code.

The age groups covered by the study are almost equally presented, except for the 66+ age group: age group 25-35 – 163 participants (23,35%), 36-45 - 171 individuals (24,50%), the 46-55 age group is the most numerous - 196 individuals (28,08%), 56-65 age group - 139 (19.91%), and the least presented age group of 66+ - only 29 (4,15%) i.e. all age groups were included. The relative share of dentists having over 20-years length of service is distinctly high (54.33%). It suggests more representative measurement/reporting of the impact of dental-specific factors, causing MSD. The major share of dentists, participating in the study, is as follows: general practitioners (58.03%), followed by the group, practicing mostly conservative dental care (12.49%) and prosthetic dental medicine (10.29%). Comparatively lower is the share of those practicing specific narrow specialties such as: oral surgery (7.40%), pediatric dental medicine (6.47%), periodontology (3.12%) and the least share is that of orthodontology (2.20%).

3. Results

Pain incidence and character for the dentists in Bulgaria

Significant share of dentists – 569 (81.87%) reported musculoskeletal type of pain they associated with their professional activity. The share of dentists with no such complaints is tiny – 126 (18.13%). We entered a question on assessing the type of pain among the dentists with MSD in order to be able to characterize its nature. The study results demonstrate that the major pain type is chronic (85.84%) (Fig.1).




Figure 1: Chart presentation of MSD-related pain incidence and nature

Pain localization

The study subjects listed the zone/zones of musculoskeletal pain due to their professional activity. The results reveal that it is present mostly in the neck and shoulder area –320 (32.92%), and almost equally distributed in the remaining

zones – for 230 (23.66%) of the respondents the pain is localized in the upper back area; for 216 (22.22%) – in the lower back area, while upper extremities pain is listed by 206 (21.19%) (Fig.2).




Figure 2: Pain Zones (N=566)

Pain Intensity

The respondents used their individual discretion in defining the pain intensity as mild, moderate, strong and severe/unbearable. Most of the respondents - 388 (64.88%)

have defined it as moderate, 132 (22.07%) – as mild. It is disturbing that not that small percent of the respondents define it as strong -73 (12.21%) and severe/unbearable 5 (0.84%) (Fig.3).




Figure 3: Musculoskeletal Pain Level (N=569)

Pain Duration

The survey participants determined pain duration in time as followed: experiencing pain for years - 381 respondents (67.55%), for months – 148 (26.24%), and for weeks – 35 (6.21%) (Fig.4). These results illustrate the severity of the problem.




Figure 4: Period of pain manifestation in time (N=564)

4. Discussion

Based on published data we avail of, the distribution of musculoskeletal disorders among the dentists in Bulgaria falls within similar ranges. Our data display that 81.87% of our practitioners have MSD-related pain. This gives us reason to conclude that the occupational factors, conducive to the development of MSD are similar for all dentists, irrespective of their dental practice geographical region.

Our study has found out that the Bulgarian dentists suffer mostly of chronic pain (85.84%), predominantly defining its intensity as moderate (64.88%). Lower levels are reported in a Nepal study – 55.9% and a study conducted in Greece, where the pain in 30% of the dentists is of a chronic nature (3).

Besides pain character and intensity, we also focused on its localization. The main areas of pain manifestation among the Bulgarian dentists prove to be neck and shoulder

(32.92%), while comparatively lower, yet relatively equal are the shares of the other three area types listed, namely upper and lower back (23.66% and 22.22%) and upper extremities (21.19%). For this specific indicator our data partially overlap with those officially published.

The neck and shoulder pain type distribution is key both for dentists in Bulgaria and abroad, though our country level is comparatively lower – 32.92%. The levels indicated in officially published papers, that we have access to, are significantly higher. For example neck pain goes to 63% of the dentists in Nebraska, 65% in Denmark, 72% in Sweden, 81,9% in Nigeria, 58% - in Australia, 38% - in Israel, 58,8% - Nepal, 56,3% - Poland, 28% в – in Iran, 58,8% in Yemen, 54.4% in Saudi Arabia, and 26% in Greece.

For the remaining three pain zones the officially published data are also higher: for Saudi Arabia the study results report 73.5% back pain; in Canada this percentage is 57%, out of which 35.5% for lower back pain; in Yemen the lower back pain distribution is in the range of 80%; in Nepal lower back pain – 52.4%, while upper back pain – 41.7% and upper extremities – 36.9%; in Australia back pain is present in 53.7%; in India this level is 75,5%; Iran - 33%; Finland - 38%; Poland – 60.1% and 44% - in the upper extremities; for Greece – 46% lower back pain; Denmark – 50%.

5. Conclusions

- 1) The level of MSD-related pain among the dentists in Bulgaria is very high (81.87%), similar to the officially published data.
- 2) The majority of the respondents (67.55%) state that this MSD-provoked pain exists for years.
- 3) The largest is the share of neck and shoulder pain - 32.92%, while for the other three zones – upper and lower back and the upper extremities area, pain distribution is almost equal ranging from 21.19% to 23.66%.
- 4) The prevailing pain intensity specified by our dentists is of the “moderate” type – 64.88%. Much lower is the number of respondents, classifying it as “mild” (22.07%) and “strong” (12.21%).
- 5) The major pain type, reported by the dentists, has a “chronic nature” - 85,84%.

All the data in this introducing article on the severity of the problems, related to MSD among dentists and the conclusions made give us grounds for future research on both the relationship of MSD to the socio-demographic and occupational factors and pain management methods.

References

- [1] Abiodun-Solanke IM, Agbaje JO, Ajayi DM, Arotiba JT. Prevalence of neck and back pain among dentists and dental auxiliaries in South-western Nigeria, Afr J Med Med Sci. 2010 Jun;39(2):137-42.
- [2] Acharya RS, Acharya S, Pradhan A, Oraibi S. Musculoskeletal disorders among dentists in Nepal. Journal of Nepal Dental Association 2010, vol 11, No 2, Jul-Dec:107-113
- [3] Alexopoulos EC, Stathi I, Charizani F. Prevalence of musculoskeletal disorders in dentists. BMC Musculoskelet Disord 2004;5:16
- [4] Al-Kholani A: Musculoskeletal Symptoms among Dentists in Yemen. The International Journal of Occupational Safety and Ergonomics (JOSE), 2011
- [5] Auguston TE, Morken T. Musculoskeletal problems among dental health personnel: a survey of the public dental health services in Hordaland (in Norwegian). Tidsskr Nor Laegeforen. 1996;116:2776-80.
- [6] Biller FE. Occupational hazards in dental practice. Oral Hyg 1946;36:1994.
- [7] Chowanadisai S, Kukiattrakoon B, Yapong B, Kedjarune U, Leggat PA. Occupational health problems of dentists in Southern Thailand. Int Dent J 2000;50:36-40.
- [8] Dajpratham P, Ployetch T, Kiattavorncharoen S, Boonsirirath K. Prevalence and associated factors of musculoskeletal pain among the dental personnel in a dental school. J Med Assoc Thai 2010; 93: 714-721.
- [9] Finsen L, Christiansen H, Bakke M: Musculoskeletal disorders among dentists and variation in dental work. Appl Ergon 1998; 29:119-125
- [10] Hayes MJ, Smith DR, Cockrell D. A systematic review of musculoskeletal disorders among dental professionals. Int J Dent Hyg 2009b; 7: 159-165
- [11] Kierklo A, Kobus A, Jaworska M, Botuliński B. Work-related musculoskeletal disorders among dentists - a questionnaire survey. Ann Agric Environ Med. 2011 Jun;18(1):79-84
- [12] Leggat PA, Smith DR: Musculoskeletal disorders self-reported by dentists in Queensland, Australia. Aust Dent J 2006;51(4):324-7
- [13] Li TKL, Lo ECM, Wong HHA: Self-reported occupation-related health problems in Hong Kong dentists. Hong Kong Dental Journal; 2006; v. 3: 39-44
- [14] Marshall ED, Duncombe LM, Robinson RQ, Kilreath SL. Muscu- loskeletal symptoms in New South Wales dentists. Aust Dent J 1997; 42:240-6.
- [15] Milerad E, Ekenwall L: Symptoms of the neck and upper extremities in dentists. Scand J Work Environ Health 1990; 16:129-134
- [16] Puriene A, Aleksejuniene J, Petrauskiene J, Balciuniene I, Janulyte V. Self-reported occupational health issues among Lithuanian dentists. Ind Health 46(4);369-74 (2008), PMID 18716385
- [17] Shrestha BP, Singh GK, Niraula SR. Work related complaints among dentists. JNMA 2008 Apr-Jun;47(170):77-81
- [18] Shugars DA, Miller D, Williams D, Fishburne C, Strickland D. Musculoskeletal pain among general dentists. Gen Dent 1987;35: 272-6.
- [19] Szymanska J. Disorders of the musculoskeletal system among dentists from the aspect of ergonomics and prophylaxis. Ann Agric Environ Med 2002; 9(2):169-173

Author Profile

Dr Peter Bojinov is in Department of Public Dental Health, FDM-Sofia, Sofia 1431, G.Sofiiski 1