

Grey Musk Shrew (*Suncus murinus*) a Favorite Prey of Barn Owl (*Tyto alba*) in Calicut, Kerala

Ajitha K. V.¹, Bobby Jose², Vineesh P. J.³

PG and Research Department of Zoology, St. Joseph's College, Devagiri, University of Calicut, Kerala, India

Abstract: The Barn Owl (*Tyto alba*) is a widely distributed resident nocturnal raptor. The pellets of Barn owl were collected and studied from the zoology wing of St. Joseph's college, Calicut, Kerala during April-July 2013. Regurgitated pellets of barn owl were analyzed to understand the dietary composition. The aim of the study is to find out seasonal difference of food preferred by the barn owl, estimating the average size of each pellet to assess the difference in total intake of food during rainy and summer seasons, to compare the amount of bones in each pellet to know the availability of shrew during the rainy and summer season. It was found that the diet comprised of exclusively the rodent *Suncus murinus* and its consumption among two seasons was not similar.

Keywords: Barn owl, *Tyto alba*, Grey musk shrew, *Suncus murinus*, pellets, skull, rodent

1. Introduction

Barn owl is a cosmopolitan species found all over the world except Antarctica [9]. There is so many stories, myths, believe and superstitions about barn owl all over the world. This bird is notoriously associated with death, evil spirit and misfortune. This may be due to its horror appearance, call and nocturnal activity.

2. Literature Review

Extensive studies have been conducted on the diet, feeding and nesting habits of barn owl in different parts of the world by various research workers. (C.D.Marti 1972&1974, Alivizators & Gounter 1999, Bose & Guidali 2001, Mushtaq-ul-Hassan *et al.* 2007 a,c, Leonardi & Arte 2006, Margini & Facure 2008, Nadeem *et al* 2012) The available literature on the Indian Barn owl is comparatively less and is confined to feeding behavior and diet composition (Neelananarayanan 2007 a, b, Santhanakrishan *et al.* 2010, Unfortunately, we have little research studies on barn owls of Kerala. (Jayson & Babu 2009).

3. Study Area

The zoology wing of St. Joseph's College Devagiri, Calicut (N 11° 15' 51.0761" & E 75° 50' 6.8575") selected as the study area during the period of April-July 2013. An adult barn owl has inhabited in the lab for last two years. This area is always free from all other disturbances especially in night time. That may be one of the reasons for selecting this area as its territory by the bird.

4. Materials and Methods

Total 60 pellets were collected in summer and monsoon. The length and width of pellet were first measured using a verniers caliper (with an accuracy of 0.05mm). The samples were then weighed using electronic balance (with accuracy to 0.01g). To identify the food items preferred by barn owl, pellet analysis must be done. Pellets were analyzed following the methods adopted in earlier studies [6]. At the time of

analysis each pellet was put in warm water for softening. The pellet materials were carefully opened up using tweezers. During pellet examination skulls, bones, feathers, insect remains were separated out for identification. Collected bones and hairs are examined. To identify the skull present in the pellets, rodents, particularly rats were trapped from the surrounding area, killed and boiled in water. Separated bones and skulls compared with pellet compositions. Mean abundance and standard deviation of different food components in pellet were calculated on an excel sheet and tabulated. Normality of collected data was tested using Jarque-Bera test [11]. Affinity between the size of pellet between two seasons and difference in availability of shrew during the rainy and summer season were assessed with t test.

5. Result

All pellets are compact, hard, oval shape and dark green in color. Skulls, bones, and hairs were embedded into the pellet. There is a significant difference between the size of pellet during two seasons ($t= 4.38$, $P= 0.0001$) and availability of shrew during the rainy and summer season ($t= 5.87$, $P> 0.0001$). The most striking feature of this study is the presence of at least one mammalian (rodent) skull in all 60 pellets. Among these, 3 pellets were contained more than one skull that may be depending up on the size of the pellet and the food taken by the barn owl. All the skulls in the 60 pellets were similar in size and structure. These entire skulls were well analyzed and studied. For better identification these skulls were compared with skull of other local rodents trapped from that area such as bandicoot rat (*Bandicoot malabarica*), house rat (*Rattus rattus*) and grey musk shrew (*Suncus murinus*). Skulls in the pellets were exactly matches the skull of grey musk shrew. Pellet skulls were entirely different from the local rodent's skull in their size and structure. Skulls obtained from all 60 pellets were unique and was found to be the skull of Grey musk shrew (*Suncus murinus*). This particular species is identified with the help of experts. In spite of these skulls and bones, negligible amount of insect remains (order: Coleoptera) were found in some pellets.

Volume 4 Issue 7, July 2015

www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

Table 1: Mean and Standard deviation of pellets in two seasons

Features of Pellets	Summer	Monsoon
	Mean ± SD	Mean ± SD
Length (mm)	46.8 ± 7.4	42.3 ± 5.0
Width (mm)	32.7 ± 1.7	31.8 ± 5.1
Dry weight (mg)	2696.7 ± 485.7	2068.0 ± 618.7
Weight of bones (mg)	1676.0 ± 408.1	1090.4 ± 363.3
Weight of scat (mg)	1020.7 ± 457.5	977.6 ± 420.1



Plate 1: Skull image of *Suncus murinus*

6. Discussion

The size of the pellets in the study is similar to the previous reports. The t-test results indicate the reduction in shrew availability during monsoon season, in this locality and the absence of alternate food during monsoon season. It means that the consumption of grey musk shrew in the diet of the Barn Owls among two seasons was not similar, and was different. The unavailability of grey musk shrew during monsoon in the study area, inversely affects the weight of pellets and bones in each pellet.

Several investigations on the diet of the Barn Owl in different geographical regions confirmed that small mammals were the main components of the diet [1], [2], [5], [8] & [10]. From this study it has been proved that it prefers only *Suncus murinus* even though the other prey availability was more in this area. It means that *Suncus murinus* may be the favorite food item of barn owl. Previous studies had revealed that diet of the barn owl includes heavy consumption of bats [15] & [23] but very negligible amount had noticed in Madurai, Tamil Nadu [22]. We hadn't found any remains of bats in all 60 pellets.

Intake of small birds as the prey had been identified in many reports by various workers, exclusively species of sparrows, finches, starling and thrushes [22]. We couldn't find bird remains in the samples. Reptiles and amphibians were absent in every pellets. Arthropod preyed up on are exclusively insects belongs to Coleoptera, Isoptera, Orthoptera and Hymenoptera [8], [20] & [22]. In our study we could have found out certain remains of coleoptera in two pellets.

7. Conclusions

The study was mainly concentrated on the presence of unique skulls of *Suncus murinus* in all 60 pellets. It revealed

that Barn owl shows a special preference for our commonest shrew, Grey musk shrew (*Suncus murinus*) which is generally distributed throughout peninsula. From our study it is clear that there is a variation in the diet of these birds depending on the area and availability of food items. In Calicut it feeds exclusively on *Suncus murinus*, even though other species of rodents were common in the campus.

8. Acknowledgement

We thank Dr. P.O. Nameer (Professor and Head, Department of Wildlife Science College of Forestry, KAU, Thrissur, Kerala) who helped us to identify the skull of *Suncus murinus*. One of the authors (Ajitha K.V) expresses her gratitude to the Kerala State Govt. for granting her the E-grants Scholarship (Govt.of Kerala) to pursue doctoral research.

References

- [1] A.Travaini, J.A.Donazar, O., Ceballos , A.Rodriguez, F. Hiraldo, M.Delibes, "Food habits of common Barn-owls along an elevational gradient in Andean Argentine Patagonia". *Journal of Raptor Research*, 31(1), pp 59–64, 1997.
- [2] C.D Marti, "Feeding ecology of four sympatric owls" *Condor*, 76, pp 45-61, 1974.
- [3] C.D. Marti, „The years of barn owl prey data from a Colorado nest site“, *Wilson Bulletin*, 85, pp 85– 86, 1972.
- [4] E.A. Jayson, S. Babu "Studies on owls of southern Western Ghats", KFRI research report, 2009.
- [5] F.Pezzo, F.Morimando, "Food habits of the barn owl, *Tyto alba*, in a Mediterranean rural area: Comparison with the diet of two sympatric carnivores", *Italian Journal of Zoology*, 62, pp 369– 373. 1995.
- [6] F.W. Schueler, "A new method of preparing owls pellets boiling in NaOH", *Bird Banding*, 43, pp 142. 1972.
- [7] G. Leonardi, G.L.D. Arte, „Food habits of the Barn Owl (*Tyto alba*) in a steppe area of Tunisia" *Journal of Arid Environment*, 65, pp 677–681, 2006.
- [8] H.Alivizatos , V. Gountner, "Winter diet of the Barn Owl (*Tyto alba*) and Long-eared Owl (*Asio otus*) on northeastern Greece: a comparison" *Journal of Raptor Research*, 33, pp160–163, 1999.
- [9] I.R.Taylor, "Barn Owls. *Predator-prey Relationships and Conservation*" Cambridge University Press, Cambridge, England, 1994.
- [10] J.A .Gubanyi, R.M. Case, G Wingfield, "Diet and nesting success of Barn Owls breeding in western Nebraska". *American Midland Naturalist*, 127, pp 224–232. 1992.
- [11] J.H. McDonald, "Handbook of Biological Statistics "Sparky House Publishing, Baltimore , 2008.
- [12] L. Magrini, K.G.Facure, "Barn Owl (*Tyto alba*) predation on small mammals and its role in the control of hantavirus natural reservoirs in a periurban area in southeastern Brazil" *Brazilian Journal of Biology*, 68(4), pp 733–740, 2008.

- [13] M.Bose, F.Guidali, "Seasonal and geographic differences in the diet of the Barn Owl in an agroecosystem in Northern Italy". *Journal of Raptor Research*, 35(3), pp 240–246,2001.
- [14] M. Mahmood-ul-Hassan, M.A. Beg , M.Mushtaq-ul Hassan," Locality related changes in the diet of Barn Owl (*Tyto alba stertens*) in agroecosystem in Central Punjab, Pakistan". *Wilson Journal of Ornithology*, 119(3), pp 479–483,2007a.
- [15] M.I.Bellocq, " A review of the trophic ecology of the Barn Owl in Argentina." *Journal of Raptor Research*, 34(2), pp 108–119. 2000.
- [16] M.Mahmood-ul-Hassan, M.A.Beg ,H. Ali, "Seasonal variation in the diet of the barn owl *Tyto alba stertens* in central Punjab, Pakistan". *Acta Zoological Sinica*, 53(3), pp 431–436 ,2007c.
- [17] M.S Nadeem, S.M.K. Imran, T. Mahmood, , A.R. Kayani, S.I. Shah," A Comparative study of the diets of Barn Owl (*Tyto alba*) and Spotted Owlet (*Athene brama*) inhabiting Ahmadpur East, Southern Punjab, Pakistan" *Animal Biology*, 62(1), pp 13–28. 2012.
- [18] P.Neelanarayanan, "Diet of Barn Owl *Tyto alba stertens* Hartert, (1929) in a portion of Cauvery delta, Tamil Nadu, India." *Zoos' Print Journal*, 22(8), pp 2777–2781. 2007a.
- [19] P.Neelanarayanan," Technique for estimation of barn owl (*Tyto alba stertens* Hartert, 1929) prey biomass with special reference to mandible length body weight ratio of small mammals" *Zoos' Print Journal*, 22(1), pp 2519–2521, 2007b.
- [20] R. Santhanakrishnan, "Ecology of Barn Owl, *Tyto alba* (Scopoli) with special reference to its population, feeding and breeding in Mayiladuthurai, Tamil Nadu, South India." Ph.D. Thesis, Bharathidasan University, Trichy, India, 1995.
- [21] R.Santhanakrishnan, "Studies on population, food habits and nesting of Barn Owl, *Tyto alba*(Scopoli) in a portion of Cauvery river basin." M.Phil. Thesis, Bharathidasan University, Trichy, India 1987.
- [22] R.Santhanakrishnan, A.M.S.Ali, U.Anbarasan, "Diet Variations of the Barn Owl *Tyto alba* (Scopoli, 1769) in Madurai District, Tamil Nadu, Southern India". *Podoces*, 5(2), pp 95–103, 2010b.
- [23] R.Sommer , H.Zoller, D.Kock, W.Bohme , A.Griesau," Feeding of the barn owl, *Tyto alba* with first record of the European free-tailed bat, *Tadarida teniotis* on the island of Ibiza (Spain, Balearics)." *Folia Zoologica*, 54(4), pp 364–370. 2005.
- [24] S.T.Alvarez-Castaneda, N. Cardenas, L. Mendez "Analysis of mammal remains from owl pellets (*Tyto alba*), in a suburban area in Baja California". *Journal of Arid Environment*, 59, pp 59–69, 2004.



Dr. Boby Jose, Associate Professor, PG and Research Department of Zoology, St. Joseph's College, Devagiri, University of Calicut, Kerala, India. Research area is ORNITHOLOGY, PhD on the Ecological Isolation of Babblers (*Turdoides* spp.)



Dr. Vineesh.P.J. Assistant Professor, PG and Research Department of Zoology St. Joseph's College Devagiri University of, Calicut, Kerala, India. Research area is Entomology, PhD on the Forest Litter Entomology.

Author Profile



Ajitha K.V. A PhD candidate in the PG and Research Department of Zoology, St. Joseph's College, Devagiri, University of Calicut, Kerala, India. Research area is ORNITHOLOGY, study on the Kole Wetland Birds and Egrets of Muriyad, Thrissur District, Kerala, South India