

- [10] Stuart. B. 2004. Infrared Spectroscopy , John Wiley and sons Ltd, England.
- [11] Nakomato N. 2009. Infrared and Raman Spectra of Inorganic and Coordination Compounds , 6th.ed., John Wiley and Sons, Inc., New Jersey.
- [12] Robert M. Silverstein, Francis X. Webster and David J. Kiemle . 2005. Spectrometric Identification of Organic Compound , 7thed. Jon-wiley. New York .
- [13] Shakir M. and Azim Y. 2005. Metal Directed Synthesis of Hexaazamacrocyclic Complexes of Co (II), Ni (II), Cu (II) and Zn (II) and Their Physico-Chemical Studies, Polish J.Chem. 79,1115-1121.
- [14] Majeed A. ,Yousif E. and Farina Y.2010.Synthesis and Characterization of Transition Metal Complexes of 2-Thioacetic Acid Benzothiazole Ligand ,J. Al-Nahrain University (Science) . 13(1):36-42.
- [15] Zahid H. Chohan, Hazoor A. Shad and Nasim F. H.2009.Synthesis, Characterization and Biological Properties of Sulfonamide Derived Compounds and Their Transition Metal Complexes, Appl. Organometal. Chem. 23:Issue 8 , 319-328.
- [16] Imam H , Kumar B. and Shafayat M.D.2011.Mixed Ligand Complexes of Transition Metal Chelates of 1-nitroso-2-naphthol and 8-hydroxyquinoline with Picolinic Acid and Quinaldinic acid Oriental ,J. Chemistry. 27 (1): 287-291.
- [17] Suttan D. (1968).Electronic Spectra of Transition Metal Complexes. Mc Grow-Hill, Publishing, London.
- [18] B.N. Figgis and M. A. Hitchman. (2000). Ligand Field Theory And its Application.Wiley-VCH, New York, Singapore, Toronto.
- [19] Hashim.C.S and Alais.M.F . (2012). Synthesis, spectroscopic study of Pt (IV), Au (III), Rh(III),Co(II) and V (IV) complexes with sodium[5-(p-nitro phenyl) - /4-phenyl-1,2,4-triazole-3-dithiocarbamate] and cytotoxicity assay on rhabdomyosarcoma cellline of heavy metals J.Baghdad Sci. ,9(4).
- [20] Jabur .H. K., Alias .M F. and Al-Azez Kareem.T. A.. (2012) .preparation, characterization and biological activity of some complexes of potassium 2-carbomethoxy amino-5-trithiocarbonate 1,3,4-thiadiazole. J. Baghdad .Sci., 9(3).
- [21] Mukkanti .K. and Singh .R. P. . (1988).Synthesis and characterization of Ru(III), Rh(III), Pt(IV) and Ir(III) thiosemicarbazone complexes Proc. Indian Acad. Sci (Chem. Sci) 100(1):21-26.
- [22] Datt .N.K. and Chakder. N.C. (1971). Element of the first transitional period J.Inorg.Nucl.Chem,33:393.
- [23] Colinand .R.L. and Holt. S.R .(1963).Inorg.Chem,2,8.
- [24] Chow .S.T. and Mcauliffe .C.A. (1995).Metal complexes of amino acids and derivatives—XIII Some nickel(II) complexes of L-arginine zwitterions or L-argininate anions. A novel ionic → bidentate perchlorate rearrangement brought about by pressure[1–5].J.Inorg.Nucl.Chem,37(4). P : 1059-1064.
- [25] Arar .H.B. (2006).Trans.Met.Chem.,1,239,(5).
- [26] Nicholls .D. (1973).The Chemistry of Iron,Cobalt and Nickel.1st Ed., Pergman Press,Oxford .
- [27] Deshmuke .K.G. and Bhoje. R.A. (1978).Spectral, thermal and magnetic properties of Ni(II) complexes of p-phenylazo-salicylaldehydes.J. Inorg. Nucl.chem,40(1),135-136.
- [28] Nath .P. and Dhumwad. S.D. (2012). Synthesis, characterization and antimicrobial studies of Co(II), Ni(II), Cu(II) and Zn(II) complexes derived from a Schiff base of 2-[(4-Methyl-2-oxo-2H-chromen-7-yl)oxy]acetohydrazide with 3-formyl-2-hydroxy quinoline and 3-formyl-2-mercapto quinoline.J.Chem. Pharm.l Reas.,4(1),P: 851-865.
- [29] Alias .M. F., Kassum.H. and Shakir .C.(2013). Synthesis, spectral, thermal and antibacterial studies of Cd(II), Mn(II) and Fe(III) complexes containing trithiocarbonate 1,3,4-thiadiazole moiety.J.King Saud Univ. Sci.V 25,(2), P 157–166.