



















- from aqueous solution using modified corn stalk. *Journal of Hazardous of material*. 176: 650-656.
- [5] Athar, M., and Vohara, S.B., (1995). Heavy metals and Environment. Wiley Eastern Ltd., New Age International Publishers Ltd. New Delhi. pp. 204-206
- [6] Abbas, H.S., Thamer, J.M., and Jenan, A., (2012). Equilibrium and kinetic studies of adsorption of heavy metals onto activated carbon. *Canadian Journal on Chemical Engineering and Technology*. 3(4): 86-92.
- [7] Jarup L., (2014). Hazard of heavy metal contamination. *British Medical Bulletin*. 68(1): 167-182.
- [8] Nandal, M., Hooda R., Dhania, G., (2014). Tea waste as sorbent for the removal of heavy metals from waste water. *International Journal of current Engineering and Technology*. 4(1): 243- 245.
- [9] Doris, K.L., Zhang, Y., and Shukka. A., (2000). The removal of heavy metal from aqueous solution by sawdust adsorption. *Journal of Hazardous Material*. 8(2): 33-42.
- [10] Mathialagan, T.; Viraraghavan, T. (2002) Adsorption of Cd from aqueous solutions by perlite, *J. Hazard. Mater.* 94 : 291–303.
- [11] Inbaraj, S. B.; Wang, J. S.; Lu, J. F.; Siao, F. Y.; & Chen, B. H. (2009). Adsorption of toxic mercury (II) by an extracellular biopolymer poly(c-glutamic acid). *Bioresource Technology*, 100, 200–207.
- [12] Zhang, F. S., Nriagu, J. O., & Itoh, H. (2005). Mercury removal from water using activated carbons derived from organic sewage sludge. *Water Research*, 39, 389–395.
- [13] Nathaniel, O., Edith, B.J., and Sulaiman, O.I., (2012). Removal of Pb<sup>(II)</sup> and Ni<sup>(II)</sup> ions from aqueous solutions by adsorption onto activated locust bean (*Parkia biglobosa*) husk. *Archives of Applied Science Research*. 4(6): 2308-2309.
- [14] Bailey, S.E., Olin, T.J., Bricka, R.M., and Adrian, D., (1999). A review of potentially low-cost sorbent for heavy metals. *Journal of Water Research*. 33(11): 2469-2479.
- [15] Kumar, U., (2006). Agricultural products and by-products as a low cost adsorbent for heavy metal removal from water and wastewater: A review. *Scientific Research and Essay*, 1(2): 33-37.
- [16] Nasim A.K., Shaliza I. and Piarapakaran S., (2004). Review paper; elimination of heavy metals from wastewater using agricultural wastes as adsorbents. *Malaysian Journal of Science*, 23: 43-51.
- [17] Surchi, K.M.S., (2011). Agricultural wastes as low cost adsorbents for Pb Removal: Kinetics, equilibrium and thermodynamics. *International Journal of Chemistry*, 3(3): 103-112
- [18] Hala, A.H., (2013). Removal of heavy metals from waste water using agricultural and industrial waste as adsorbents. *Housing and Building National Research Center Journal*. 9: 267282.
- [19] Langmuir I., (1918) The adsorption of gases on plane surfaces of glass, mica, and platinum, *J. Am. Chem. Soc.* 40 :1361–1368.
- [20] Freundlich H., (1907) Veber die adsorption in loesungen (Adsorption in solution), *Z. Physik. Chem.* 57 : 385–470.
- [21] Igwe, J. C.; Abia, A. A. (2007) Equilibrium sorption isotherm studies of Cd(II) Pb(II) and Zn(II) ions detoxification from waste water using unmodified and EDTA-modified maize husk. *Electron. J. Biotechnol.* 10: 536–548.
- [22] Ho, Y. S. and McKay, G., (1998), A comparison of chemisorption kinetic models applied to pollutant removal on various sorbents, *ProSafety and Environmental. Protection*, 76B(4):332-340
- [23] Horsfall Jr., M.; Spiff, A. I.; Abia, A. A. (2004) Studies on the influence of mercaptoacetic acid (MAA) modification of cassava (*Manihot esculenta* Cranz) waste biomass on the adsorption of Cu<sup>(II)</sup> and Cd<sup>(II)</sup> from aqueous solution. *Bull Korean. Chem. Soc.* 25: 969–976.
- [24] Mahamadi C, Nharingo T (2010) Utilization of water hyacinth weed (*Eichhornia crassipes*) for the removal of Pb(II) Cd(II) and Zn(II) from aquatic environments: An adsorption isotherm study. *Environ. Technol.* 31:1221–1228.
- [25] Vijayaraghavan K.; Padmesh, T. V. N.; Palanivelu, K.; Velan, M. (2006) Biosorption of nickel(II) ions onto *Sargassum wightii*: application of two-parameter and three-parameter isotherm models. *J. Hazard Mater.* 133: 304–308.
- [26] Mondal, D. K.; Nandi, B. K.; Purkait, M. K. (2013) Removal of mercury (II) from aqueous solution using bamboo leaf powder: Equilibrium, thermodynamic and kinetic studies, *Journal of Environmental Chemical Engineering* 1: 891–898
- [27] Han, R. P.; Zhang, L. J.; Song, C.; Zhang, M.M.; Zhu, H. M.; Zhang, L. J. (2010). Characterization of modified wheat straw, kinetic and equilibrium study about copper ion and methylene blue adsorption in batch mode, *Carbohydr. Polym.* 79: 1140–1149
- [28] Chakravarty, P., Sarma, N.S., and Sarma, H.P., (2010). Biosorption of cadmium (II) from aqueous solution using heart wood powder of *Areca catechu*. *Chemical Engineering Journal*. 162: 949-955.
- [29] Uluozlu, O.D.; Sari, A.; Tuzen, M.; Soylak, M. (2008) Biosorption of Pb(II) and Cr(III) from aqueous solution by lichen (*Parmelinatiaceae*) biomass, *Biores. Technol.* 99: 2972–2980.
- [30] Srivastava, V.C.; Swamy, M.M.; Mall, I.D.; Prasad, B.; Mishra, I.M. (2006) Adsorptive removal of phenol by bagasse fly ash and activated carbon: equilibrium, kinetics and thermodynamics, *Colloids Surf. A: Physicochem. Eng. Aspects* 272: 89–104
- [31] Garg U., Kaur, M.P., Jawa G.K., Sud D., and Garg, V.K., (2008). Removal of cadmium (II) from aqueous solutions by adsorption on agricultural biomass. *Journal of hazardous material*. 154: 1149- 1157.