

References

- [1] Adhikary SP, 2002, Utilization of region specific cyanobacteria as biofertilizer for rice-a case study from Orissa; Conference paper.*BiotechnolMicr Sustainable Utilization* 47-56.
- [2] Bhagwati J.,2013, High Commission of India London.,Dialogue With India Series II,India House, May 21
- [3] Bhattacharyya S., Nayak B., Choudhary, NK., 2011. Response of diazotropic cyanobacteria *Nostoccarneum* under pesticide and UV-s tress.*Chemosphere* 84: 131-135.
- [4] Bhattacharyya S., Nayak B., Deep P.R, Sahu J K, 2014. Methyl parathion induced changes in primary photochemistry of a natural nitrogen engineer *Nostocpunctiforme*. *Indian Journal of Applied Research*. 4(7): 53-56
- [5] Bloom D. E. 2011.,Population Dynamics in India and Implications for Economic Growth, Harvard School of Public Health,<http://www.hsph.harvard.edu/pgda/working.html>
- [6] Choudhary, KK,2011. Occurance of nitrogen fixing cyanobacteria during different stages of paddy cultivation, Bangladesh *J.Plant taxon*. 18(1): 73-76
- [7] Deep P.R., Bhattacharyya S.,Nayak B.,2013. Cyanobacteria in wetlands of the industrialized Sambalpur district of India.*Aquatic Biosystem*, Biomed 9,14.
- [8] Desikachary TV:*Cyanophyta*, Indian Council of Agricultural Research, New Delhi.1959, 686 pp.
- [9] Komárek J and Anagnostidis K: Cyanoprokaryota-2.Teil/ 2nd Part: Oscillatoriales. In: Büdel B, Krienitz L, Gärtner G and SchagerlM(ed.), *SüßwasserfloravonMittleuropa* 19/2. Elsevier/Spektrum, Heidelberg 2005.
- [10] Komárek J, Anagnostidis K: Cyanoprokaryota-1.Teil/ 1st Part: Chroococcales. In: Ettl H, Gärtner G and Heynig H, Mollenhauer D (ed.), *SüßwasserfloravonMittleuropa* 19/1. Elsevier/Spektrum, Heidelberg 1998.
- [11]Nayak S, Prasanna R:Soil pH and its role in cyanobacterial abundance and diversity in rice field soils.*ApplEcol Environ Res* 2007, **5**: 103 – 113
- [12]Nilsson, M., Bhattacharya, J., Rai, AN., Bergman, B., 2002, Colonization of roots of rice (*Oryza sativa*) by symbiotic *Nostoc* strains, *New Phytologist*159,517-525.
- [13]Odisha Agriculture Statistics, 2008-2009. Directorate of Agriculture and Food Production, India, Bhubaneshwar.
- [14]Olson, J.M., & Blankenship, R.E. 2004, Thinking about the evolution of photosynthesis, *Photosynthesis Research* 80: 373–386.
- [15]Rashtriya Chemicals and Fertilizers Ltd . 2012-2013, Sustainability Report,
- [16]Sahu J K, Bhattacharyya S, Deep P R, Nayak B.,2015.Relative Tolerance and Nitrogenase activity of Several Heterocystous Cyanobacteria to Herbicide, Hiltachlor, 50 EC.*International Journal of Toxicological and Pharmacological Research* 2015; 7(1); 17-22
- [17]Singh R.N., 1961. Role of Blue Green Algae in nitrogen economy in India agriculture. Indian agriculture research institute- New Delhi 175
- [18]Tripathy I. C., Ayyappan S. and Adhikary S. P., 1990.Isolation and growth of local heterocystous cyanobacteria for using as biofertiliser.National Symposium on cyanobacterial Nitrogen Fixation, New Delhi, pp 275-278.
- [19]Venkataraman G.S., 1981. Blue green Algae; a possible remedy to nitrogen scarcity.*Current Science*. 50: 253-256.