

Synthesis, Characterization and Antimicrobial Studies of Mn(II) Co(II) AND Zn(II) SCHIFF Base Complexes Derived from L-Arginine and 2-Hydroxy-1-Naphthaldehyde

Iniama, Grace E¹, Iorkpiligh, Terungwa Isaac²

¹Department of Pure and Applied Chemistry, University of Calabar, Nigeria

²Department of Chemistry, College of Education Oju, Benue State, Nigeria

Abstract: *The ligand, L-arginine-2-hydroxy-1-naphthaldehyde (LAHN) was synthesized and used for the preparation of Mn(II), Co(II) and Zn(II) complexes. These were characterized using infrared, electronic absorption data, molar conductivity and elemental analysis. The infrared spectra data of this ligand and its metal complexes show that the ligand is a bidentate molecule which coordinated to the metal ions through the azomethine nitrogen and the carboxylate oxygen atom of the L-arginine-2-hydroxy-1-naphthaldehyde, the Schiff base. From the electronic absorption spectra data available, tetrahedral geometry have been assigned to the prepared complexes. The ligand and all complexes have good antimicrobial effect on Escherichia coli, Staphylococcus aureus, Salmonella typhi and Candida albicans and Aspergillus niger strains at different concentrations. The complexes show enhanced activities and their activities against the microorganisms increased with increasing concentrations.*

Keywords: Schiff base complex, bidentate ligand, antimicrobial activity

1. Introduction

Schiff bases also called imines are characterized by the presence of the azomethine functional group (-C=N-) and are usually formed by condensation of an aldehyde or ketone with a primary amine [1]. This condensation reaction is a facile reaction due to the good electrophilic characteristic properties of the carbonyl towards the amine group [2]. Over the years, Schiff bases have been used extensively as versatile ligands with transition metals which are able to form very stable complexes because they donate electrons into the vacant d-orbitals of the metal atoms /ions and in some cases these ligands accept electrons from these low oxidation state metals thus effecting strong bonding through synergetism [3]. It has been observed that such complexation is the explanation for their therapeutic effects against many diseases including cancer [4]. This versatility of Schiff base ligands is shown in the biological, analytical and industrial applications of their metal complexes. This also makes further investigations in this area highly desirable [5], and as such, current researches dealing with metal complexes of hetero atoms in Schiff bases have expanded their applications enormously to include a vast diverse fields of human endeavors.

The interest on Schiff bases and their metal complexes is heightened by the fact that they provide synthetic docking models for development of new chemotherapeutic drugs that fit well into binding sites of macromolecular targets [6]. This is usually achieved by providing unhindered entry sites for Deoxyribonucleic Acid (DNA) structure and conformation [7]. Many Schiff bases and their complexes are known to show promising antifungal, antibacterial, antiviral including anti HIV and anticancer activities [8]. In addition, some

Schiff bases containing N & O donor atoms are also effective anti-corrosion agents and serve in many other analytical purposes in organic and inorganic chemistry fields, hence their wide applications in industries. [9][10]. This research work reports on the synthesis of Mn(II), Co(II) and Zn(II) complexes of L-arginine-2-hydroxy-1-naphthaldehyde, their characterization and their antimicrobial effectiveness.

2. Experimental

Synthesis of L-arginine-2-hydroxy-1-naphthaldehyde (LAHN)

Ethanol solutions of L-arginine (1.742g, 0.01mol) and 2-hydroxy-1-naphthaldehyde (1.722g, 0.01mol) were slowly mixed together while stirring. Three drops of glacial acetic acid were added to the mixture and magnetically stirred for two hours. The resulting yellow precipitate was filtered and washed several times with ethanol then dried over fused calcium chloride in a desiccator.

Synthesis of the metal(II) L-arginine-2-hydroxy-1-naphthaldehyde

The Schiff base ligand, L-arginine-2-hydroxy-1-naphthaldehyde (1.742g, 0.01M) was dissolved in 30ml ethanol. To this solution, warmed ethanolic solution of metal(II) chloride hydrate (0.01M) was added, followed by few drops of sodium hydroxide. This mixture was magnetically stirred for two hours and the precipitate formed was filtered, washed and dried in a desiccator over fused calcium chloride.

Table 1: Physical data of the compounds

| Compound | Molecular Formula | Colour | Melting point(°C) | Conductivity (Ω ⁻¹ cm ² mol ⁻¹) | % yield | Found (Cald) (%) | | | |
|---|---|--------|-------------------|---|---------|------------------|---------|--------|--------------------|
| | | | | | | M | C | H | N |
| LAHN | C ₁₇ H ₁₉ N ₄ O ₃ | Yellow | 203.7 | 1.17 | 87.73 | - | 63.25.8 | 17.1 | (62.7)(4.9) (16.8) |
| [Mn(LAHN) ₂]Cl ₂ | MnC ₃₄ H ₃₈ N ₈ O ₆ Cl ₂ | Brown | >300 | 1.65 | 54.04 | 7.1 | 52.3 | 4.9 | 14.4 |
| [Co(LAHN) ₂]Cl ₂ | CoC ₃₄ H ₃₈ N ₈ O ₆ Cl ₂ | Brown | 219.2 | 3.82 | 51.19 | 7.5 | 52.0 | 4.8 | 14.4 |
| [Zn(LAHN) ₂]Cl ₂ | ZnC ₃₄ H ₃₈ N ₈ O ₆ Cl ₂ | Yellow | 213.4 | 1.53 | 52.28 | 8.2 | 51.6 | 4.8 | 14.2 |
| | | | | | | (7.9)(51.5) | (4.8) | (14.2) | |

Table 2. Selected Infrared Bands (KBr disc, cm⁻¹)

| Compound | νOH | NH | C=O | C=N | COO ⁻ | M-N | M-O |
|---------------------------------------|-------|-------|-------|-------|------------------|------|------|
| LAHN | 3400 | 3797w | 1681s | 1629s | 1342s | 590w | 455w |
| Mn(LAHN) ₂ Cl ₂ | 3368s | 3760m | 1687s | 1629w | 1393s | 489s | 424s |
| Co(LAHN) ₂ Cl ₂ | 3386s | 3589w | 1687m | 1623w | 1356s | 581w | 469w |
| Zn(LAHN) ₂ Cl ₂ | 3240s | 3678w | 1681m | 1627w | 1346s | 606m | 455w |

Table 3: Electronic Spectral data

| Compound | Band(cm ⁻¹) | Assignment |
|---------------------------------------|-------------------------|---|
| LAHN | 47,846 | n-σ* |
| | 30,395 | π-π* |
| | 30,211 | n-π* |
| Mn(LAHN) ₂ Cl ₂ | 48,076 | n-σ* |
| | 30,487 | π-π* |
| | 24,027 | ⁶ A ₁ → ⁴ T ₁ (⁴ P, ² D) |
| | 21,038 | ⁶ A ₁ → ⁴ E(⁴ G) |
| Co(LAHN) ₂ Cl ₂ | 47,846 | n-σ* |
| | 30,487 | π-π* |
| | 25,000 | ⁴ A ₂ → ⁴ T ₁ |
| Zn(LAHN) ₂ Cl ₂ | 47,846 | n-σ* |
| | 38,759 | π-π* |
| | 30,120 | L → M |

Table 4: Antimicrobial studies of L-arginine-2-hydroxy-1-naphthaldehyde and its metal complexes

| Ligands/complexes | Zone of inhibition in (mm) | | | | | |
|---|----------------------------|-----------------|----------------|------------------|-------------------|-----------------|
| | Concentrations (µg/ml) | <i>S. typhi</i> | <i>E. coli</i> | <i>S. aureus</i> | <i>C. albican</i> | <i>A. niger</i> |
| LAHN | 1000 | 14 | 15 | 15 | 19 | 15 |
| | 500 | 9 | 9 | 10 | 9 | 10 |
| [Mn(LAHN) ₂]Cl ₂ | 1000 | 17 | 24 | 16 | 22 | 18 |
| | 500 | 10 | 10 | 10 | 12 | 12 |
| [Co(LAHN) ₂]Cl ₂ | 1000 | 14 | 12 | 12 | 10 | 14 |
| | 500 | 7 | 8 | 8 | 8 | 9 |
| [Zn(LAHN) ₂]Cl ₂ | 1000 | 23 | 22 | 17 | 25 | 21 |
| | 500 | 12 | 10 | 12 | 13 | 14 |
| Streptomycin | 1000 | 25 | 23 | 23 | - | - |
| | 500 | 15 | 15 | 13 | | |
| Ketoconazole | 1000 | - | - | - | 24 | 26 |
| | 500 | | | | 13 | 16 |

*Measurements are in mm diameter, ≥8.0 mm indicates no inhibition

3. Discussion

All the synthesized ligand and complexes are variously coloured solids; soluble in common coordinating solvents like DMF, DMSO, acetone and ethanol as shown in Table 1. They are stable in air and exist in powdery form indicating their polymeric nature. The molar conductance values are very low in the range of 1.17-3.82s cm²mol⁻¹ suggesting

their non electrolytic nature and only little degree of polarity. [11]. The melting points of the complexes are high indicating strong bonding system within the compounds. [12]

Infrared of LAHN and its complexes

The IR spectrum (Table 2) of the ligand gave a characteristic band at 3400cm⁻¹ assignable to ν(O-H) intramolecular

hydrogen bond. [13]. The band at 1629cm^{-1} is characteristic of azomethine $\nu(\text{C}=\text{N})$ stretching vibration while the bands at 1342cm^{-1} is assigned to carboxylate ion $\nu(\text{COO}^-)$. The $\nu(\text{C}=\text{N})$ vibration at 1629cm^{-1} in the ligands was shifted to lower wave number of 1625cm^{-1} , 1623cm^{-1} and 1624cm^{-1} at coordination to Mn(II), Co(II) and Zn(II) respectively. The $\nu(\text{O}-\text{H})$ intramolecular hydrogen bond at 3400cm^{-1} disappeared on complexation with another band emerging at 1342cm^{-1} in each complex, thus confirming participation of the carboxylate oxygen atom in coordination. The bands at 489, 581 and 610cm^{-1} in the spectra of the metal complexes are due to $\nu(\text{M}-\text{N})$ while the band at 424cm^{-1} , 460cm^{-1} and 455cm^{-1} are due to $\nu(\text{M}-\text{O})$. This shows that the ligand acted as a bidentate ligand coordinating to the metal ions through the carboxylate oxygen and azomethine or imino nitrogen.

Electronic spectra of LAHN and its complexes


The electronic absorption spectra are often very helpful in the evaluation of results furnished by other methods for structural identification. The electronic absorption measurements (Table 3) are hereby used for assigning the stereochemistry of metal ions [14]. The electronic absorption spectra of the Schiff base have bands at 47,846, 30,395 and $30,211\text{cm}^{-1}$. These are assigned $n - \sigma^*$, $\pi - \pi^*$ and $n - \pi^*$ transitions respectively which are attributed to intra-ligand transition which are due to the presence of lone pair of electrons on the hetero oxygen atoms and double bonds in the structure of the ligand [12][14]. The $\pi - \pi^*$ transitions are associated with azomethine group. In the spectra of the complexes, these bands were shifted to higher absorption wave numbers indicating the coordination of the ligand to the metal ions. The metal ions also show some d-d transitions. The spectrum of the $[\text{Mn}(\text{LAHN})_2]\text{Cl}_2$ gave a single d-d transition band at $24,038\text{cm}^{-1}$ assignable to forbidden ${}^6\text{A}_1 \rightarrow {}^4\text{A}_1$ transitions which are typical of tetrahedral manganese(II) complex [15]. The electronic spectrum of $[\text{Co}(\text{LAHN})_2]\text{Cl}_2$ has a single d-d transition at $25,000\text{cm}^{-1}$ which can be assigned to ${}^4\text{A}_2 \rightarrow {}^4\text{T}_1$ transition, a characteristic of tetrahedral cobalt(II) complex [15]. The electronic spectra of the $[\text{Zn}(\text{LAHN})_2]\text{Cl}_2$ complex gave no d-d transition as expected for a d^{10} configuration where all the electrons are paired. Thus giving the diamagnetic properties of the zinc complex [16].

The procedure used for screening is as reported by [17][18]. The synthesized compounds dissolved in DMF were examined *in vitro* by Paper Disk Diffusion method. In this method, all wares used were sterilized in a hot air oven. The gram-positive bacteria, *Staphylococcus aureus*, *Salmonella typhi* and gram-negative *Escherichia coli* and the fungi *Candida albican* and *Asperigillus niger* were subcultured and incubated for 6-8 hours in Nutrient and Sabouraud Dextrose agar respectively. The viable bacterial cells were evenly swabbed onto the Nutrient agar plates while the fungal spores were swabbed onto the Sabouraud Dextrose agar plates. The paper disks, 6.0 mm diameter, were soaked in the different test samples (concentrations, 1000 & 500 $\mu\text{g}/\text{ml}$), drained and using sterilized forceps placed in the agar plates. The plates were then incubated for 48 hours (bacteria) and 72 hours (fungi) at 37°C . After the incubation period, the zones of inhibition were measured in mm (Table 3).

Antimicrobial Activity

The ligands and the metal complexes show very good antimicrobial properties against *Escherichia coli*, *Staphylococcus aureus*, *Salmonella typhi* and *Candida albican* strains with the metal complexes having a higher inhibition activities (14–25 mm diameter inhibition zones) compared to the free ligands (14–19 mm diameters). In all the cases studied, the zinc complexes proved to be a better antimicrobial agent (up to 25 mm diameter inhibition zone against *C. albican*) followed by Mn(II) complexes (24 mm diameter against *E. coli*) as compared to the free ligands.

The following structure is proposed for the complexes



4. Conclusion

The Schiff base and novel complexes of Mn(II), Co(II) and Zn(II) L-arginine-2-hydroxynaphthaldehyde and the Schiff base ligand were synthesized and characterized using infrared spectra, electronic absorption spectra, molar conductivity and elemental analysis. These complexes and

the ligand were stable to air, non hygroscopic and variedly coloured. Their electronic and infrared spectra data revealed the bidentate nature of the ligand through which it coordinates to the metal ions using the azomethine nitrogen and carboxylate oxygen. Tetrahedral geometries are assigned to these complexes. The ligand and complexes showed very good antimicrobial properties against

Escherichia coli *Staphylococcus aureus* *Salmonella typhi* and *Candida albicans* strains with the metal complexes having a higher inhibition activities than the free ligand.

References

- [1] Graham, S. & Fryhle, C.B. (2004). *Organic Chemistry*. New York: John Wiley and Sons Inc. pp. 738-40.
- [2] Petil, S., Jadhav, S.D. & Petil, U.P. (2012). Natural Acid Catalyzed Synthesis of Schiff Base Under Solvent-Free Condition: As Green Approach. *Archives of Applied Science Research*, 4(2): 1074-1078.
- [3] Ayi, A. A. (2006). *Understanding Inorganic Chemistry Through Problems and Solutions. Second Edition*. Calabar: University of Calabar Press. pp. 355-529.
- [4] Singh, H.L., Singh, J. & Mukherjee, A. (2013). Synthesis, Spectral, and In Vitro Antibacterial Studies of Organosilicon(IV) Complexes with Schiff Bases Derived from Amino Acids. *Bioinorganic Chemistry and Applications*, 2013(1), Article ID 425832, 9 pages
- [5] Ashraf, M.A., Wajid A., Mahmood, K., Maah, M.J. & Yusoff, I. (2011). Spectral Investigation of the Activities of Amino Substituted Bases. *Oriental Journal of chemistry*, 27 (2): 363-372.
- [6] Kumar, S.S., Nath, P.S. & Kumar, P.A. (2013). In-silico identification and molecular Docking studies of quinolone Resistance Determining Region (QRDR) of E. Coli DNA Gyrase- A with ofloxacin Schiff Bases. *International Journal of Pharmtech. Research*, 5(4): 1794-1803.
- [7] Zhang, Y., Wang, X. & Ding, L. (2010). Interaction between tryptophan-vanillin Schiff base and herring sperm DNA. *Journal of the Serbian Chemical Society*, 75(9): 1191-1201.
- [8] Raj, A. A., Vinnarasi, J. & Rose, G.L. (2013). Molecular Docking of some Isatin Schiff Bases Using Hex. *World Journal of Pharmacy and Pharmaceutical Sciences*, 2(5): 3058-3064.
- [9] Mishra, A.P., Sharma, N. & Jain, R. (2012). Microwave synthesis, spectral, thermal and antimicrobial studies of some Ni(II) & Cu(II) Schiff Bases complexes. *Advances en quimica*, 7(1): 77-85.
- [10] Kotkar, S.N. & Juneja, H.D. (2013). Synthesis, Characterization, and Microbial studies of N.O Donor Schiff Base Polymeric Complexes. *Hindawi Publishing Corporation Journal of Chemistry* 1: 1-5.
- [11] Ogunniran, K.O., Ajanaku, K.O., James, O.O., Nwinyi, C.O. & Allensela, M.A. (2008). Cobalt(II) complexes of mixed antibiotics: Synthesis, Characterization, antimicrobial potential and their effect on alkaline phosphate activities of selected rat tissues. *International Journal of Physical Science*, 3(8): 177-182.
- [12] Didarul, A. C., Uddin C.M. & Hoque F. (2010). Dioxouranium(VI) Complexes of Some Bivalent Tridentate Schiff-base Ligands Containing ONS Donor. *Set Chiang Mai Journal of Science*. 37(3) : 443-450.
- [13] Munde A.S, Shelke V.A, Jadhav S.M, Kirdant A.S, Vaidya R.S, S. G. Shankarwar S.G, and
- [14] Raman, N., Ravichandran, S. & Thangaraja, C. (2004). Co(II), nickel(II) and Zn(II) Complexes of Schiff base derived from benzyl-2,4-dinitrophenylhydrazone with aniline. *Journal of Chemical Science*, 116(4): 215-219.
- [15] Osowole, A. A. & Ott, I. (2012). Synthesis, characterization, *invitro* Anticancer and antimicrobial properties of some metal(II) complexes of 4-[2,3-dihydro 1H-inden-4-ylimino)methyl] benzene-2,4-diol. *International Research Journal of Pure and Applied Chemistry*, 2(2): 156-169.
- [16] Mishra, A.P. & Gupta, P. (2011). Effect of Chelation On Therapeutic Potential of Drugs: Synthesis, Structure, Antimicrobial And Insecticidal Activity of 3d-Metal Complexes Involving Schiff-Bases. *Journal of Chemical and Pharmaceutical Research*, 3(2): 150-161.
- [17] Iniama, G.E., Nfor, E.N., Okon, E.D. and Iorkpilgh, T.I. (2014). Antimicrobial activities of synthesized zinc(II) mixed ligand complexes derived from 2-acetylpyridine-4-phenylsemicarbazone and nitrogen-sulphur monodentate ligands. *International Journal of Scientific & Technology Research*. 3 (11) 73-76.
- [18] Chondhekar T. K (2012) Synthesis, Characterization and Antimicrobial Activities of some Transition Metal Complexes of Biologically Active Asymmetrical Tetradentate Ligands. *Advances in Applied Science Research*, 3 (1): 175-182

Author Profile



Iniama, Grace E. holds a Ph.D degree in Inorganic Chemistry and a Master degree in Analytical Chemistry. She is currently a Senior Lecturer in the Department of Pure and Applied Chemistry, University of Calabar-Nigeria



Iorkpilgh, Terungwa I. holds a Master degree in Inorganic Chemistry from the University of Calabar, lectures at the College of Education, Oju and he is currently doing his Ph.D degree still in Inorganic Chemistry in the University.