

References

- [1] I. Sener, N. Sener, S. Eriskin. Synthesis and absorption spectra of some novel hetaryltetrakisazocalix[4]arene derivatives. *Dyes and Pigments*. **2013**. Vol.96. P.256-263.
- [2] J. O. Otutu, Synthesis and Application of azo dyaderived from 2-amino-1, 3,4-thiadizole-2-thiol on polyester fiber // *IJRRAS* 2013.v.15 .No..2.P.292-2963.
- [3] S.M. Abdallah Metal complexes of azo compounds derived from 4-acetamidophenol and substituted aniline// *Arabian Journal of Chemistry* (2012),v.3,No. 5, P.251–256
- [4] Dostanci J., Valentic N., Uscumlic G., Mijin D. Synthesis of 5-(substituted phenylazo)-6-hydroxy-4-methyl-3-cyano-2-pyridones from ethyl 3-oxo-2-(substituted phenylazo)butanoates. // *J. Serb. Chem. Soc.*, 2011. V. 76. No 4. P. 499–504.
- [5] Bustos C., Alvarez-Thon L., Carcamo G.J., Ibanezd A., Sanchez C. 2-[2-(3-Chlorophenyl)hydrazinylidene]-1,3-diphenylpropane-1,3-dione. // *Acta Cryst.*, 2011. V. E67. P. o1450–o1451.
- [6] Mahmoud A.A., El-Sayed G.M., Morsy N.S. Conductivity measurements of the deaggregation constants and the solvated species parameters of some azo (3-diketone) derivatives // *Al-Azhar J. Pharm. Sci.*, 1994. V. 14. P. 1-11.
- [7] Dunning T.H. Gaussian-basis sets for use in correlated molecular calculations .I. The atoms boron through neon and hydrogen. // *J. Chem. Phys.*, 1989. V. 90, № 2. P. 1007.
- [8] Kendall R.A., Dunning T.H., Harrison R.J. Electron affinities of the first-row atoms revisited. Systematic basis sets and wave functions // *J. Chem. Phys.*, 1992. V. 96. P. 6796.
- [9] Shchegol'kov E.V., Sadchikova E.V., Burgart Ya.V., Saloutina V.I.Synthesis and Structure of 4-Hydroxy-4-fluoroalkyl-1,4-dihydroimidazo[5,1-c][1,2,4]triazines. // *Russian Journal of Organic Chemistry*, 2009. V. 45. No. 4. P. 572–580.
- [10] Wilson A.K., Woon D.E., Peterson K.A., Dunning T.H. Gaussian basis sets for use in correlated molecular calculations. IX. The atoms gallium through krypton. // *J. Chem. Phys.*, 1999. V. 110. P. 7667.
- [11] Prascher B.P., Woon D.E., Peterson K.A., Dunning T.H., Wilson A.K. Gaussian basis sets for use in correlated molecular calculations. VII. Valence, core-valence, and scalar relativistic basis sets for Li, Be, Na, and Mg. // *Theor. Chem. Acc.*, 2011. V. 128. P. 69–82
- [12] Cabot JM, Fuguet E, Ràfols C, Roses M. Fast high-throughput method for the determination of acidity constants by capillary electrophoresis. II. Acidic internal standards. *J Chromatogr A*. 2010;1217(52):8340–5.
- [13] Jetse Reijenga, Arno van Hoof, Antonie van Loon, and Bram Teunissen, Development of Methods for the Determination of pK_a Values// *Anal Chem Insights*. 2013;V. 8.P. 53–71.
- [14] Tanner E.M. The infra-red absorption spectra of some α-phenylhydrazono-ketones and alcohols. // *Spectrochim. Acta*, 1959. V. 15. P. 20.
- [15] Shcheglova TL, BN Melnikov, Belokurova OA., Laboratory workshop on the use of dyes//Textbook. Ivanovo, 2002.P. 108.