

Antimicrobial Susceptibility Pattern and Microbiological Profile of Uropathogens among Patients in a Tertiary Care Hospital in Bihar

R R Prasad¹, Vijay Shree²

¹Department of Microbiology, Patna Medical College and Hospital, Patna

²Department of Community Medicine, Indira Gandhi Institute of Medical Sciences, Patna

Abstract: *Background:* Urinary tract infection is one of the most prevalent infections. Microorganisms causing UTI vary in their susceptibility to antimicrobials due to widespread use of inappropriate and empirical antibiotic therapy. Urinary tract infection (UTI) is also the most common nosocomial infection. *Objective:* This study was designed to determine the microbiological profile and antimicrobial susceptibility patterns of uropathogens among patients referred to Patna medical college and hospital Patna. *Material and Method:* A total of two hundred and fifty (250) samples of urine from patients attending different inpatient and outpatient departments were included in the study. Urine samples were inoculated on Nutrient agar, Blood agar and McConkey agar plates by streaking. Inoculated plates were then incubated aerobically at 37°C for 24 hours. After 24 hours of incubation, isolated colonies were picked up and Gram staining was done. Motility test and other biochemical tests were done for further identification of bacterial isolates using suitable Controls. Finally Antibiotic Susceptibility Test (AST) was performed to detect the degree of sensitivity or resistance of the pathogen isolated from the patient to an appropriate range of antimicrobial drugs on Mueller-Hinton agar (MHA) plates by Kirby-Bauer disc diffusion method. *Results and Conclusion:* Out of the total of two hundred and fifty (250) samples examined in the laboratory during the study period only one hundred two (102) patients were found to have bacteriological infection. Among the bacteriologically positive cases, UTI was more common in females. Among the bacterial isolates *Escherichia coli* was the commonest pathogen in both males and females, followed by *Klebsiella* species. Sparfloxacin and Gatifloxacin were the most effective antibiotics in vitro for the Gram Negative bacilli isolated, while Azithromycin was most effective against the Gram Positive cocci. The Gram Negative uropathogens showed a high degree of resistance to cephalosporins. While the Gram Positive cocci showed highest resistance to Norfloxacin and ofloxacin. It is due to the excessive use of antimicrobials for all sorts of infections that uropathogens responsible for UTI are increasingly showing resistance to antibiotics.

Keywords: Uropathogen, Urinary Tract Infection, AST, *E.coli*, Cephalosporin

1. Introduction

Urinary tract infections (UTIs) are one of the most common infectious diseases in clinical practice both in community and hospital settings. Due to its frequency and recurrence, UTI poses a real challenge to the medical professionals. UTI is classified into uncomplicated and complicated infections. The empirical choice of antimicrobial treatment is generally guided by susceptibility data provided by regional microbiological laboratories, however, since samples of uncomplicated UTIs are rarely sent for culture, these data are mainly found in complicated UTIs [1, 2].

Etiology of UTI is influenced by factors like age, sex, diabetes, urinary catheterization and others. Urinary tract infection are more common in women than men, It could be due to the proximity of genital tract and urethra /anus. Nearly 10% of women experience a UTI during their lifetime [3]. The spectrum of bacteria which causes complicated UTI is much broader than uncomplicated ones. Resistances to antimicrobial agents have undergone dramatic variations and consequently the empirical treatment of UTI needs constant updating of the antibiotic sensitivity of the main uropathogens of that area.

A patient is said to have a urinary tract infection, when there is the presence of over 1×10^5 organisms per ml in the midstream sample of urine [4]. The human urinary tract system comprises of the kidneys, ureters, bladder and

urethra. Infections in any of these anatomical sites are referred to as UTI. Infections extending to the bladder leads to cystitis while those involving the kidneys leads to pyelonephritis [5]. *Escherichia coli* is the most common cause of urinary tract infection [6,7] and accounts for approximately 90% of first urinary tract infection in young women [7]. The symptoms and signs include increased urinary frequency, dysuria, hematuria and pyuria. Flank pain is associated with upper tract infections. None of these symptoms or signs is specific for *Escherichia coli* infection [8].

Antimicrobial susceptibility profile in respect to causative microbes may significantly reduce morbidity and mortality, cost of treatment, and duration of hospitalization if duly provided to medical practitioners and clinicians in a rapid and timely fashion [9].

2. Material and Method

This study was carried out in Department of Microbiology Patna Medical College, Patna for 6 months, from march 2014 to September 2014. A total of two hundred fifty (250) midstream, clean catch urine samples were collected in wide-mouthed, sterile, screwcapped universal plastic containers and received in the laboratory routinely from patients attending OPD (Out patient department) and also from indoor. A brief clinical history of the patients and antibiotic intake was also taken.

The specimens collected were examined by microscopy and then put up for culture. The samples were inoculated on Nutrient agar, Blood agar and McConkey agar plates by streaking. Inoculated plates were then incubated aerobically at 37°C for 24 hours. After 24 hours of incubation, isolated colonies were picked up and Gram staining was done. Motility test and other biochemical tests were done for further identification of bacterial isolates. Control strains used were as follows:

- Escherichia coli*: ATCC 25922
- Staphylococcus aureus*: ATCC 25923
- Klebsiella pneumonia*: ATCC 700603
- Staphylococcus saprophyticus*: ATCC 15305
- Proteus mirabilis*: ATCC 7002
- Staphylococcus epidermidis*: ATCC 14990
- Pseudomonas aeruginosa*: ATCC 27853
- Enterococcus faecalis*: ATCC 29212

Finally, Antibiotic Susceptibility Test (AST) was performed to detect the degree of sensitivity or resistance of the pathogen isolated from the patient to an appropriate range of antimicrobial drugs. AST was done on Mueller-Hinton agar (MHA) plates by Kirby-Bauer disc diffusion technique [9] using commercially available antibiotic discs (HiMedia, Mumbai). Interpretation of results was done based on the diameter of the zone of inhibition as per guidelines laid down by CLSI (Clinical and Laboratory Standards Institutes). The antibiotics and the concentrations at which they were used were as follows:

- Sparfloxacin (5 µg) Norfloxacin (10 µg) Cefuroxime (30 µg)
- Cephalexin (30 µg) Ofloxacin (5 µg) Gentamycin (10 µg)
- Azithromycin (15 µg) Cotrimoxazole (25 µg) Gatifloxacin (5 µg)
- Ciprofloxacin (5 µg) Amoxicillin (30 µg) Ceftriaxone (30 µg) Amikacin (30 µg)

3. Results

Out of the total of two hundred fifty (250) samples received and examined in the laboratory during the study period only one hundred two (102) patients were found to have bacteriological infection (40.80%). The remaining samples of one hundred forty eight (148) patients were found to be sterile (59.20%).

Out of the one hundred two (102) positive samples, 36 (35.29%) were male patients and 66 (64.71%) were female patients [Table 1].

Table -2 shows that amongst the one hundred two (102) bacteriologically infected samples, *Escherichia coli* was the commonest pathogen (59.81%) responsible for urinary tract infection followed by *Klebsiella pneumoniae* (18.62%), *Staphylococcus aureus* & CONS (Coagulase negative staphylococcus) (15.68%), *Proteus mirabilis* (2.95%), *Pseudomonas aeruginosa* (1.96%) and mixed - Staph/ E.coli (0.98).

Out of sixty one (61) strains of *Escherichia coli* isolated from the urine culture, 45 strains were found to be sensitive to Gatifloxacin (73.77%) and Sparfloxacin (73.77%), Amikacin (59%), Cotrimoxazole (54.09%), Gentamycin (49.18%), Ceftriaxone (45.90%), Ciprofloxacin (41%) each

and Ofloxacin (39.34%). The remaining antibiotics were less effective.

Out of sixteen (16) strains of *Staphylococcus aureus* and CONS isolated from the urine culture, 15 were found to be sensitive to Azithromycin (93.75%), followed by Sparfloxacin (81.25%), Ciprofloxacin (68.75%), Ceftriaxone (68.75%), Gatifloxacin (68.75%), Cotrimoxazole (50%), Cefuroxime (50%), Amikacin (50%), Gentamycin (50%) and Cephalexin (25.00%). The remaining antibiotics were effective in less than 25% strains.

High degree of resistance was seen in *E. coli* with reference to Azithromycin (85.25%), Cephalexin (70.5%), Norfloxacin (67.22%), Amoxicillin (65.56%), Cefuroxime (63.96%), Ofloxacin (60.66%), Ceftriaxone (54.10%) and Ciprofloxacin (59%). Ciprofloxacin resistance was more in *E. coli* as compared to resistance seen in the other Gram Negative uropathogens.

High rates of resistance was seen among the Gram Negative uropathogens isolates to cephalosporins like Cephalexin, Cefuroxime and Ceftriaxone. Ceftriaxone resistance was 54.10% in *E. coli*, 42.11% in among *Klebsiella*, 66.66% in *Proteus* and 50% in *Pseudomonas*.

Amongst the Gram Positive isolates the commonest uropathogen was *Staphylococcus aureus*, While most isolates of *Staphylococcus aureus* and CONS were resistant to Norfloxacin (87.50%).

Table 1: Gender-wise distribution among the cases suffering from UTI

Sex	Total number of isolates under study (n=102)	
	Number	Percentage(%)
Male	36	35.29
Female	66	64.71

Table 2: Distribution of different pathogenic organism among cases of UTI

Pathogenic organisms	Total isolates under study	
	Number	Percentage (%)
<i>Escherichia coli</i>	61	59.81
<i>Klebsiella pneumoniae</i>	19	18.62
<i>Proteus mirabilis</i>	3	2.95
<i>Pseudomonas aeruginosa</i>	2	1.96
<i>Staphylococcus aureus</i> and CONS	16	15.68
Mixed (staph/E. coli)	1	0.98

4. Discussion

This study showed that *E. coli* was the commonest pathogen causing complicated and uncomplicated UTI amongst the several organisms known to cause UTI, including *P. aeruginosa*, *S. saprophyticus*, *S. epidermidis*, *Enterococcus spp.*, *P. mirabilis*, *Klebsiella pneumoniae* etc. as reported by earlier workers [10-14]. Among the non-fermenters *Pseudomonas aeruginosa* was isolated as an uropathogen particularly in the intensive care units, although in a very few cases (1.96%). Furthermore, most other workers elsewhere have reported the involvement of *Klebsiella* as the second most important pathogen in UTI cases [15-26, 27-

30], in the same way present study shows a greater incidence of *Klebsiella* (18.62%).

While ciprofloxacin and ofloxacin are the most extensively used fluoroquinolones for the treatment of UTIs, the emergence of resistance for fluoroquinolones is based on several factors [14, 18, 28, 29]. Resistance to ciprofloxacin has emerged in a variety of genera belonging to the family Enterobacteriaceae. Our findings concur with such findings reported earlier [31, 32]. Apart from the notable resistance of *E. coli* to ciprofloxacin, other organisms were also found to be resistant to ciprofloxacin especially *K. pneumoniae*, *Pseudomonas* spp., *Proteus* spp. etc. Also, fluoroquinolone resistance in *E. coli* has emerged particularly in patients with urinary tract infections who have received fluoroquinolone prophylaxis [16-23]. An association between the increase in quinolone prescriptions and an increase in bacterial resistance has been reported from several countries [16-19, 23]. Usually, the prevalence of fluoroquinolone resistance is related to the intensity of antibiotic use [16].

5. Conclusion

The present study shows an increasing antibiotic resistance trends amongst UTI patients. Due to excessive use of antimicrobials for all sorts of infections, uropathogens responsible for UTI are increasingly showing resistance to antibiotics. So it is imperative to rationalize the use of antimicrobials and to use these conservatively. The present study also provides the knowledge of uropathogens and their antimicrobial susceptibility pattern which will help in appropriate and judicious antibiotic usage in our health care setup.

References

- [1] McNulty, C.A., M.R.J. Livermore and D.M. Clinical, 2008. relevance of laboratory-reported antibiotic resistance in acute uncomplicated urinary tract infection in primary care. *Journal of Antimicrobial Chemotherapy*, 58: 1000-8.
- [2] Carl, J., 2006. Urinary tract infections in women: diagnosis and management in primary care. *BMJ*, 332: 94-7.
- [3] A. Masinde, B. Gumodoka, A. Kilonzo and S.E. Mshana. Prevalence of urinary tract infection among pregnant women at Bugando Medical Centre, Mwanza, Tanzania *Journal of Health Research*, Vol. 11, No. 11, July, 2009 pp 154-161.
- [4] Stammand WE and Hooton. T.M. *Dis. Clin. North Am.* 1999, 11(3): 551-559.
- [5] Anderson RU. *Urol. Clin. North Amer.* 1999, 26: 729-735.
- [6] Momoh ARM, Odike MAC, Samuel SO, Momoh AA, Okolo PO. *Berlin Journal of Post Graduate Medicine*. 2007, 9(1): 22-27.
- [7] Jawetz E. Enterobacteriaceae In: Brooks GF, Butel JS, Morse SA eds. *Medical Microbiology* 23rd ed. Stanford-Connecticut. Appleton and Lange. 2004, Pp 248-258.
- [8] Davidson S. Disease due to Infection, In: Nicholas B. Nicki, P.C, Brain, R.W eds. *Principles and Practice of*

- Medicine*. 20th ed. New York: Churchill Livingstone. 2006, Pp 467-470.
- [9] Atae RA, Tavana AM, Hosseini SMJ, Moridi K, Zadegan MG. A Method for Antibiotic Susceptibility Testing: Applicable and Accurate. *Jundishapur J Microbiol.* 2011;5(1):341-5.
- [10] Gatermann SG. Bacterial infections of the urinary tract. In: Borriello P, Murray PR, Funke G. editors. *Topley & Wilson's microbiology & microbial infections*, 10th ed. vol. III. London: Hodder Arnold Publishers; 2007. p. 671-83.
- [11] Karlowsky JA, Jones ME, Thornsberry C, Critchley I, Kelly LJ, Sahm DF. Prevalence of anti microbial resistance among urinary tract pathogens isolated from female outpatients across the US in 1999. *Int J Antimicrob Agents* 2001; 18: 121-7.
- [12] Gorbach SL, Bartlett JG, Balcklow NR. Urinary tract. In: Gorbach SL, Bartlett JG, Balcklow NR, editors. *Infectious diseases*. Philadelphia: Lippincott Williams & Wilkins Publishers; 2004. p. 861-81.
- [13] Hooper DC. Emerging mechanisms of fluoroquinolone resistance. *Emerg Infect Dis* 2001; 2: 338-41.
- [14] Canbaz S, Peksen Y, Tefvik SA, Leblebicioglu H, Sunbul M. Antibiotic prescribing and urinary tract infection. *Int J Antimicrob Agents* 2002; 20: 407-11.
- [15] Zervos MJ, Hershberger E, Nicolau DP, Ritchie DJ, Blackner LK, Coyle EA, et al. Relationship between fluoroquinolone use and changes in susceptibility to fluoroquinolones of selected pathogens in 10 United States teaching hospitals, 1991-2000. *Clin Infect Dis* 2003; 37: 1643-8.
- [16] Arslan H, Azap OK, Ergönül O, Timurkaynak F. Risk factors for ciprofloxacin resistance among *Escherichia coli* strains isolated from community-acquired urinary tract infections in Turkey. *J Antimicrob Chemother* 2005; 56: 914-8.
- [17] Karlowsky JA, Kelly LJ, Thornsberry C, Jones ME, Sahm DF. Trends in antimicrobial resistance among urinary tract in the United States. *Antimicrob Agents Chemother* 2002; 46: 2540-5.
- [18] Kahlmeter G. An international survey of the antimicrobial susceptibility of pathogens from uncomplicated urinary tract infections: the ECO-SENS Project. *J Antimicrob Chemother* 2003; 51: 69-76.
- [19] Anjum F, Kadri SM, Ahmad I, Ahmad S. A study of recurrent urinary tract infection in women attending the outpatient department of SMHS hospital, Srinagar, Kashmir, India. *JK - Practitioner* 2004; 11: 272-3.
- [20] Kauser Y, Chunchanur SK, Nadagir SD, Halesh LH, Chandrashekhar MR. Virulence factors, serotypes and antimicrobial susceptibility patterns of *Escherichia coli* in urinary tract infections. *AJMS* 2009; 2: 47-51.
- [21] Pais P, Khurana R, George J. Urinary tract infections: A retrospective survey of causative organisms and antibiotics prescribed in a tertiary setting. *Indian J Pharmacol* 2002; 34: 278-80.
- [22] Ena J, Amador C, Martinez C, Ortiz de la Tabla V. Risk factors for acquisition of urinary tract infections caused by ciprofloxacin-resistant *Escherichia coli*. *J Urol* 1995; 153: 117-20.
- [23] Tabibian JH, Gornbein J, Heidari A, Dien SL, Lau VH, Chahal P, et al. Uropathogens and host characteristics. *J Clin Microbiol* 2008; 46: 3980-6.

- [24] Collee JG, Duguid JP, Fraser AG, Marmion BP, Simmons A. Laboratory strategy in the diagnosis of infective syndromes. In: Collee JG, Fraser AG, Marmion BP, Simmons A, editors. Mackie & McCartney practical medical microbiology, 14th ed. New York: Churchill Livingstone; 1999. p. 84-90.
- [25] James HJ, John DT. Susceptibility Test Methods: Dilution and Disk Diffusion methods. In: Murray PR, Baron EJ, Jorensen JH, Landry ML, Michael AP, editors. Manual of clinical microbiology, 10th ed. Washington, D.C.: American Society for Microbiology Press; 2007. p. 1152-72.
- [26] Clinical Laboratories Standards Institute (CLSI). Performance of standards for antimicrobial disk susceptibility tests; approved standards. 10th ed. M02-A10. vol. 29. Wayne, PA: CLSI; 2009.
- [27] Boyd LB, Atmar RL, Randall GL, Hamill RJ, Steffen D, Zechiedrich L. Increased fluoroquinolone resistance with time in *Escherichia coli* from >17,000 patients at a large county hospital as a function of culture site, age, sex, and location. *BMC Infect Dis* 2008; 8: 4-10.
- [28] Hooton TM. Fluoroquinolones and resistance in the treatment of uncomplicated urinary tract infection. *Int J Antimicrob Agents* 2003; 22: S65-S72.
- [29] Arjunan M, Al-Salamah AA, Amuthan M. Prevalence and antibiotics susceptibility of uropathogens in patients from a rural environment, Tamil Nadu. *Am J Infect Dis* 2010; 6: 29-33.
- [30] Bhargavi PS, Gopala Rao TV, Mukkanti K, Dinesh Kumar B, Krishna TP. Increasing emergence of antibacterial resistance mainly in uropathogens: southeast part of India. *Intl J Microbiol Res* 2010; 2: 1-6.
- [31] Naber CK, Steghafner M, Kinzig-Schippers M, Sauber C, 27. Sorgel F, Stahlberg HJ, et al. Concentrations of gatifloxacin in plasma and urine and penetration into prostatic and seminal fluid, ejaculate, and sperm cells after single oral administrations of 400 milligrams to volunteers. *Antimicrob Agents Chemother* 2001; 45: 293-7.
- [32] Wagenlehner FM, Naber KG. Fluoroquinolone antimicrobial agents in the treatment of prostatitis and recurrent urinary tract infections in men. *Curr Infect Dis Rep* 2005; 7: 9-16

Author Profile

R R Prasad is Resident, Department of Microbiology, Patna Medical College and Hospital, Patna (Bihar) India

Vijay Shree is Senior Resident, Department of Community Medicine, Indira Gandhi Institute of Medical Sciences, Patna (Bihar) India