

Hahn-street, D-34123 Kassel). The cut-off value was used for demarcating sera into seronegative or seropositive following the manufacturer's instruction. The ELISA OD means titres were used for quantitative comparison.

3. Statistical Analysis

Data generated from this study was analysed using SPSS version 20.0. For qualitative categorized data (seroprevalence), the significance of association was measured by applying Chi square test. ELISA OD mean titres (quantitative categorized) were presented as mean \pm 2 standard error. Only seropositive titres (i.e ELISA OD mean) were used for antibody level comparison in every parameter. Student's T-test was used to compare the only seropositive IgG and IgA mean titres obtained from HNC patients and control. Other parameters assessed were cancer type, disease clinical stage, cancer histology and grade were compared using One-way analysis of variance (ANOVA). Duncan test was further used to indicate the significance of mean differences between a particular variable compared.

4. Results

(Table 1) list the levels of different classes of EBV antibody in patients and controls. All seroprevalence and antibody

levels except for VCA (IgA) were significantly higher in HNC patients in comparison with healthy controls.

(Table 2) compare the antibody levels among various EBV seropositive head and neck cancer types. The seroprevalence of IgG-VCA was 100% for all cancer type. Only IgA-VCA was statistically significant with the highest seroprevalence observed in nasopharyngeal and the lowest in salivary gland cancers. Nasopharyngeal cancer (NPC) had the highest antibody levels for all seropositive EBV antibodies compared to other cancers. Although it was only significant for IgG-EA and IgA-EA. The oral cavity (for IgG-EA) and salivary gland (for IgA-EA) cancers also had antibody level that were in the same range as NPC. Laryngeal cancer had the lowest titres.

(Table 3,4) compares the antibody levels among various seropositive HNC histologies and grades. The seroprevalence of IgG- EA, IgA-VCA and IgA-EA in HNC patients were not statistically significant with the histology and grade of cancer. Only IgA-EA antibody level was significantly associated with cancer histology and grade. lymphoepithelioma histologic type of cancer had the highest mean titres whereas undifferentiated grade type of cancer had the highest mean titre compared to other grades.

Table 1: Comparison of EBV seroprevalence and antibody titres between HNC patients and controls

Antibody	Seroprevalence n(%)		p-value	ELISA OD(means \pm s.e)		p-value
	Patient n=100	Control n=82		Patient	Control	
IgG-VCA	100(100.0)	54 (65.9)	0.000	1.989 \pm 0.05	1.482 \pm 0.07	0.000
IgG-EA	93 (93.0)	47 (57.3)	0.000	1.637 \pm 0.07	1.056 \pm 0.02	0.040
IgA-VCA	30 (30.0)	2 (2.4)	0.000	0.878 \pm 0.11	0.764 \pm 0.07	0.380
IgA-EA	75 (75%)	47 (57.0)	0.009	1.471 \pm 0.09	0.945 \pm 0.04	0.003

Seroprevalence; Chi square df=1; χ^2 value; IgG VCA=40.355, IgG EA=32.317, IgA VCA=23.617, IgA EA=6.375. ELISA OD; Student T-test value; IgG VCA=6.191, IgG EA=0.886, IgA VCA=2.150, IgA EA=5.530

Table 2: Comparison of EBV seroprevalence and antibody titres in various HNC types

Cancer type	Sample number	IgG-VCA	IgG-EA	IgA-VCA	IgA-EA
Seroprevalence n(%)					
Laryngeal	14	14 (100.0)	13(92.9)	2 (14.3)	9(64.3)
Oral cavity	19	19 (100.0)	17(89.5)	5 (26.3)	17(89.5)
Oropharyngeal	11	11 (100.0)	11(100.0)	4 (36.4)	8(72.7)
Hypopharyngeal	9	9 (100.0)	8(88.9)	4 (44.4)	5(55.6)
Nasopharyngeal	24	24 (100.0)	24(100.0)	13 (54.2)	19(79.2)
Salivary gland	11	11 (100.0)	11(100.0)	1(9.0)	8(72.7)
Sinonasal	12	12 (100.0)	9(75.0)	1(8.3)	9(75.0)
p-value*		-	0.123	0.024	0.534
ELISA-OD (means \pm s.e)					
Laryngeal		1.811 \pm 0.09	0.342 \pm 0.05 ^b	1.466 \pm 0.14	0.860 \pm 0.17 ^b
Oral cavity		2.118 \pm 0.08	1.214 \pm 0.25 ^a	1.600 \pm 0.11	1.219 \pm 0.15 ^b
Oropharyngeal		2.065 \pm 0.11	0.730 \pm 0.23 ^{ab}	1.346 \pm 0.11	0.973 \pm 0.21 ^b
Hypopharyngeal		1.959 \pm 0.14	1.011 \pm 0.51 ^{ab}	1.451 \pm 0.06	0.875 \pm 0.17 ^b
Nasopharyngeal		2.041 \pm 0.14	1.249 \pm 0.28 ^a	1.823 \pm 0.13	1.734 \pm 0.20 ^a
Salivary gland		1.865 \pm 0.10	0.790 \pm 0.22 ^{ab}	1.799 \pm 0.00	1.494 \pm 0.30 ^a
Sinonasal		1.938 \pm 0.12	0.202 \pm 0.03 ^b	1.509 \pm 0.00	0.966 \pm 0.19 ^b
p-value**		0.520	0.043	0.302	0.000

Seroprevalence; Chi square df=6; χ^2 IgG VCA=(no statistical value), χ^2 IgG EA=10.032, χ^2 IgA VCA=14.523, χ^2 IgA EA=5.078. Duncan test designated letters= (a,ab,b)

Table 3: Comparison of EBV seroprevalence and antibody titres in various HNC histology

Histology	Sample number	IgG-VCA	IgG-EA	IgA-VCA	IgA-EA
Seroprevalence n(%)					

SCC	62	62(100.0)	56(90.3)	14(22.6)	45 (72.6)
RS	3	3 (100.0)	3(100.0)	2(66.7)	3(100.0)
LEC	19	19(100.0)	19(100.0)	9(47.4)	15(78.9)
AC	3	3 (100.0)	3(100.0)	0(0.0)	2(66.7)
ACC	9	9(100.0)	8(88.9)	4(44.4)	8(88.9)
MEC	4	4 (100.0)	4(100.0)	1(25.0)	2(50.0)
p-value		-	0.685	0.131	0.590
ELISA-OD(means±s.e)					
SCC		1.986±0.05	0.751±0.13	1.579±0.08	1.274±0.09
RS		1.811±0.25	1.168±1.05	1.604±0.09	1.015±0.21
LEC		2.002±0.17	1.158±0.30	1.772±0.18	2.085±0.23
AC		2.215±0.07	1.329±0.09	-	1.848±1.21
ACC		2.132±0.08	1.050±0.31	1.576±0.17	1.376±0.19
MEC		1.618±0.19	0.434±0.27	1.560±0.00	1.980±0.76
p-value		0.483	0.558	0.824	0.006

Seroprevalence; Chi square df=5; χ^2 IgG-VCA=(no statistical value), χ^2 IgG-EA=3.099, χ^2 IgA-VCA=8.503, χ^2 IgA-EA=3.722. ELISA-OD; one way ANOVA

KEY: SCC= Squamous cell carcinoma, RS= Rhabdomyosarcoma, LEC=lymphoepithelioma carcinoma, AC=Adenocarcinoma, ACC= Adenocystic carcinoma, MEC=Mucoepidermoidcarcinoma

Table 4: Comparison of EBV seroprevalence and antibody titres in various HNC grade (differentiation)

Grade of cancer	Sample number	IgG-VCA	IgG-EA	IgA-VCA	IgA-EA
Seroprevalence n(%)					
Well differentiated	40	40(100.0)	38(95.0)	13(33.5)	29 (72.5)
Moderately differentiated	22	22 (100.0)	20(90.0)	5(22.7)	17(77.3)
Poorly differentiated	13	13(100.0)	11(84.6)	1(7.7)	8(61.5)
Undifferentiated	19	19(100.0)	19(100.0)	9(47.4)	15(78.9)
Grade x	6	6(100.0)	5(83.3)	2(33.3)	6(100)
p-value			0.372	0.372	0.531
ELISA-OD(means±s.e)					
Well differentiated		2.024±0.05	0.855±0.15	1.473±0.06	1.335±0.10 ^b
Moderately differentiated		2.030±0.89	0.811±0.02	1.756±0.16	1.430±0.21 ^b
Poorly differentiated		1.744±0.11	0.612±0.28	2.044±0.00	1.256±0.26 ^b
Undifferentiated		2.001±0.17	1.158±0.30	1.772±0.18	2.085±0.23 ^a
Grade x		2.098±0.18	0.845±0.60	1.604±0.09	0.999±0.19 ^b
p-value		0.369	0.702	0.263	0.006

Seroprevalence; Chi square; df=4; χ^2 IgG VCA=(no statistical value), χ^2 IgG EA=0.576, χ^2 IgA VCA=3.454, χ^2 IgA EA=0.053. ELISA-OD; one way ANOVA

Grade X= grade cannot be assessed.

clinical stage of disease. Although the result showed an increase in EBV IgG-VCA, IgG-EA, IgA-VCA and IgA-EA mean titres as the disease advance in stage (i.e stage 2 to stage 4).

(Table 5) compare antibody levels among head and neck cancer clinical stages. The seroprevalence and anti EBV antibody levels of were not significantly associated with the

Table 5: Comparison of EBV seroprevalence and antibody titres in various HNC clinical stage

Clinical stage	Sample number	IgG-VCA	IgG-EA	IgA-VCA	IgA-EA
Seroprevalence n(%)					
Stage 2	7	7 (100.0)	7(100.0)	4(57.1)	5(71.4)
Stage 3	28	28(100.0)	26(92.9)	6(21.4)	21(75.0)
Stage 4	65	65(100.0)	60(92.3)	20(30.8)	49(75.4)
p-value		-	0.750	0.178	0.974
ELISA-OD(means±s.e)					
Stage 2		1.968±0.05	0.819±0.18	1.453±0.13	1.263±0.20
Stage 3		1.981±0.10	0.850±0.13	1.579±0.21	1.476±0.15
Stage 4		2.219±0.18	1.339±0.56	1.692±0.08	1.490±0.12
p-value		0.601	0.473	0.483	0.823

Seroprevalence; Chi square; df=2; χ^2 IgG-VCA= (no statistical value), χ^2 IgG-EA= 0.576, χ^2 IgA-VCA=3.454, χ^2 IgA-EA=0.053. ELISA-OD; one way ANOVA

5. Discussion

In this study the seroprevalence and serum antibody levels of IgG and IgA against VCA and EA were significantly elevated in patients with head and neck cancer when compared with controls except for IgA-VCA. Similarly, Majidi *et al.* (2006) reported significantly elevated serum antibody levels against IgG-VCA, IgA-VCA and IgA-EA in HNC patients than controls. The results reflect the possibility that EBV seropositivity and elevated titres are associated with HCA development and the HNC patients harbour higher titres than healthy individuals because of the cancer burden. In contrast to this result, Abdulmir *et al.*, (2008) and Morshed *et al.*, (2002) also reported no significant difference in anti-VCA titres between patients and controls.

The IgG anti VCA was the least specific serologic parameter in this study. It was elevated to high levels in the majority of HNC patients as observed by Majidi *et al.* (2006). The incidence was 100% for all cancer types as reported to range between 62–100% depending on the site of the lesion by Majidi *et al.* (2006). This titre elevation may represent an activation of a latent EBV infection occurring after a state of immunosuppression develops in the host from the effects of the tumor (Callaghan *et al.*, 1983; Kondo *et al.*, 2004). In this study, the seroprevalence of IgA-VCA alone was associated with the NPC.

The IgG and IgA (VCA) antibody titres were not significantly associated with HNC cancer type ($p>0.05$) and this is in agreement with kamel *et al.*, (2003) who reported no association between EBV IgG-VCA level and HNC tumor site. These results disagrees with Abdulmir *et al.* (2008) who reported that the serum level of EBV IgG- and IgA-(VCA) antibodies were significantly higher in NPC group than other HNC. However NPC had the highest titres for IgG and IgA (VCA and EA) compared to other HNC. But it was only significant for IgG- and IgA- (EA). The seroprevalence and antibody titre of EBV supports several studies on EBV strong association with NPC carcinogenesis than non nasopharyngeal HNC cancers.

The oral cavity and salivary gland also had high antibody titres that were in the same range as NPC. The result agrees with Saemundsen *et al.* (1982) reported that the sera from patients with salivary gland carcinoma showed patterns of EBV IgG and IgA (VCA and EA) level within the same range seen in NPC patients. This suggests the EBV could be associated with non nasopharyngeal HNC particularly the oral cavity and salivary gland in this study. Other non nasopharyngeal HNC having lower titres like the laryngeal, hypopharynx, within the same range in this study supports Majidi *et al* 2006 and Callaghan *et al.*, 1983 who reported no difference in the antibody titers between various sites of head and epidermoid carcinoma

The serum level of EBV IgA-EA titres alone proved to be associated with tumor histology and grade. The results showed that EBV is more associated with lymphoepithelioma (LE) than other HNC histology type in the study. This results agrees with Abdulmir *et al.* (2008) who reported significant association of EBV serum IgG and

IgA levels with tumor histology in NPC patients with lymphoepithelioma (LE) having higher antibody levels than NPC patients with squamous cell carcinoma (SCC) although no similar association was found between laryngeal cancer with LE and laryngeal cancer with SCC.

The EBV IgA-EA antibody titre was significantly higher in patients with undifferentiated cancer compared to others grades having antibody levels within the same range. This is supported by kamel *et al.* (2003) who reported that there was no significant difference of EBV antibody titres among well-, moderately- and poorly- differentiated HNC. More so all the HNC patients with LE histology and undifferentiated carcinoma are the 19 out of 24 nasopharyngeal cases enrolled in the study. The evidence supports Adulamir *et al.* (2008) conclusion that EBV is clearly associated with undifferentiated NPC of LE histology than differentiated NPC of SCC histology.

In this study, the seroprevalence and antibody titres of EBV were not associated with clinical stage of the disease and this agrees with Luo *et al.* (2013) who reported that the positive rates of IgA (VCA and EA) were not related with HNC clinical stages. On the other hand, The EBV antibody titres increases as the disease progresses towards late stage as observed by Cai *et al.* (2010) and Sun *et al.* (2014) who concluded that IgA and IgG (VCA and EA) titre increases with advanced clinical stage of diseases which depicts increase burden of the disease that may affect the immune system of the patients.

6. Conclusion and Recommendation

Based on the present study there is an association between high EBV antibody titres and head and neck cancer particularly NPC, oral cavity and salivary gland cancer. The Undifferentiated-lymphoepithelioma of HNC histology was associated with EBV IgA-EA antibody titre. Further experiments on saliva and tissue samples could investigate the role, of EBV in tumorigenesis of HNC in Nigeria. The significance of one antibody and the insignificance of the other suggest the use of multiple anti EBV antibodies against more than one or more EBV antigen for diagnosis.

7. Authors' Contributions

Jibril, F.L.*¹ carried out the sampling, Data sheet survey, obtained blood samples, ELISA assay, the statistical design, statistical analysis. Aminu-Mukhtar, M.² was the major supervisor, whereas Jatau, E.D.³ and Usman, M.A.⁴ were co-supervisors. They carried out the proofreading of the article language, integrity and approved the final manuscript.

8. Acknowledgements

Special appreciation for the staff of the Department of Oncology and the Department of Hematology, Ahmadu Bello university Teaching Hospital, Shika- Zaria, Nigeria where patient's and healthy control individuals interview and sampling was carried out.

References

- [1] Abdulmir, A. S., Hafidh, R. R., Abdulmuhammen, N., Abubakar, F. and Abbas, K. A. (2008): The distinctive profile of risk factors of nasopharyngeal carcinoma in comparison with other head and neck cancer types. *BMC Public Health*, 8(400): 1-16.
- [2] Cai, Y. L., Zheng, Y. M., Cheng, J. R., Wang, W., Zhang, Y. N., Wang, W. H., Wu, Y. S., Zhong, W. M., Li, J. and Mo, Y. K. (2010): Relationship between clinical stages of nasopharyngeal carcinoma and Epstein - Barr virus antibodies Rta/IgG, EBNA1/IgA, VCA/IgA and EA/IgA. *Journal of Southern Medical University*, 30(3): 509-511.
- [3] Callaghan. D. S., Conner, B. R. and Strauss, M. (1983): Epstein – Barr virus antibody titres in cancer of the head and neck. *Archieve of Otolaryngology*, 109(72): 781-784.
- [4] Chaturvedi, A. K., Anderson, W. F., Lortet-Tieulent, J., Curado, M. P., Ferlay, J., Franchi, S., Rosenberg, P. S., Bray, F. and Gilson M.C. (2013): Worldwide trends in incidence rates of oral cavity and oropharyngeal cancers. *Journal of Clinical Oncology*, 31(36): 4550-559.
- [5] Eduardo, B., Raquel, Jorge F. C., Isabel, A., Marisa, L., Eurico, M. and Rui, M. (2010): Epstein-Barr Virus detection in nasopharyngeal carcinoma-implications in a low-risk area. *Brazilian Journal of Otorhinolaryngology*, 76(3): 310-315.
- [6] Ferlay, J., Shin, H-R., Bray, F., Forman, D., Mathers, C. and Parkin, D. M. (2010): Estimates of worldwide burden of cancer in 2008. *International Journal of Cancer*, 127(2): 2893-2917
- [7] Greespan, J. S., Greenspan, D. and Lennette, E. T. (1985): Replication of Epstein Barr virus within the epithelial cells of oral hairy leukoplakia, an AIDS-associated lesion. *New England Journal of Medicine*, 313: 1564-1571.
- [8] Gupta, K., and Metgud, R. (2013): Evidences suggesting involvement of viruses in oral squamous cell carcinoma. *Pathology Research international*. Article ID 642496, pp 1-18.
- [9] Kamel, A. H., El-Barawy, M. A., Hashish, M. H. and El-Sheikh, S. M. (2003): Epstein–Barr virus in head and neck squamous cell carcinoma. *Eastern Mediterranean Health Journal*, 9(3): 62-71.
- [10] Kondo, S., Horikawa, T., Takeshita, H. and Kanegane, C. (2004): Diagnostic value of serum EBV-DNA quantification and antibody to viral capsid antigen in nasopharyngeal carcinoma patients. *Cancer Science*, 95(6): 508-513
- [11] Lilly-Tariah, O. B., Somefume, A. O. and Adeyemo, W. I. (1999): Current evidence of head and neck cancers in Nigeria. *Head and Neck Oncology*, 1:14
- [12] Luo, Y. L., Chen, H., Peng, S. G., Lin, J. H. and Huang, P. Y. (2013): Assessment of detection assays of Epstein-Barr viral Rta-IgG, VCA-IgA, EA-IgA and Epstein-Barr viral DNA at different clinical stages in the diagnosis of nasopharyngeal carcinoma. *Zhonghua Yi Xue Za Zhi*, 26; 93(44): 3516-3519
- [13] Majidi, M., Rafat panah, H., Zamanian, A. and Hassanzadeh, N. (2006): Epstein – Barr virus antibody titres in non nasopharyngeal head and neck squamous cell carcinoma. *The Iranian Journal of Otorhinolaryngology*, 18(45): 61-65
- [14] Morshed, K., Dacewicz, P. S., Zymanski, M. and Ziaja, S. M. (2002): Epstein– Barr virus antibodies in blood serum of patients with laryngeal cancer. *Otolaryngologia Polska*, 56(1): 45-48.
- [15] Ocheni, S., Olusina D. B., Oyekunle, A. A., Ibegbulam, O. G., Kröger, N., Bacher, U. and Zander, A. R. (2010). EBV-associated malignancies. *The Open Infectious Diseases Journal*, 4: 101-112.
- [16] Onotai, L.O. and Nwagbo, A.C. (2012). Primary head and neck malignant tumor in Port Harcourt, Nigeria: A revisit. *Journal of Medicine and Medical Science*, 3(2): 122-125.
- [17] Rickinson, A.B., and Kieff, E. (2007): Epstein-Barr Virus, In: Knipe, D.M., Howley, P.M., Griffin, D.E., Lamb, R.A., Martin, M.M., Roizman, B. and Straus, S.E. (ed.), *Fields virology*, (5th ed., vol. II). Lippincott Williams & Wilkins, Philadelphia, PA. Pp 2655-2700
- [18] Saemundsen, A., Albeck, H., Hansen, J., Nielsen, N., Anvert, M., Henle, W., Henle, G., Thompson, K., Kristensen, H. and Klein, G. (1982): Epstein-Barr virus in nasopharyngeal and salivary gland carcinomas of Greenland Eskimos. *Brazilian Journal of Cancer*, 46: 721-728.
- [19] Siddiqui, S., Chandra, R., Aziz, A. And Suman, S. (2012): Epidemiology and histopathological spectrum of head and neck cancers in Bihar, Eastern India. *Asian Pacific Journal of Cancer Prevention*, 13: 3949-3953.
- [20] Sun, P. Chen, C., Chen, Y. K., Zeng, Z. J., Chen, X. L., Liu, L. Z. and Gu, M. F. (2014): Serological marker of Epstein-Barr virus Correlate with TNM classification according to the seventh edition of the UICC/AJCC staging system for nasopharyngeal carcinoma. *European Archive of Otorhinolaryngology*, 271(9): 2545-2554
- [21] Thompson, M. P. and Kurzrock, R. (2004): Epstein-Barr virus and cancer. *Clinical Cancer Research*, 10(3): 803-821.
- [22] Wong, M. M., Lye, M. S., Cheng, H. M. and Sam, C. K. (2005): Epstein-Barr virus serology in the diagnosis of nasopharyngeal carcinoma. *Asian Pacific Journal of Allergy Immunology*, 23(1): 65-67.
- [23] Young, L. S. and Rickinson, A. B. (2004): Epstein-Barr virus: 40 years on. *National Review of Cancer*, 4: 757–768.
- [24] Zheng, Y., Xia, P., Zheng, H. C., Takashi, H., Masuda, S. and Takano, Y. (2010): The Screening of viral risk factors in tongue and pharyngolaryngeal squamous carcinoma. *Anticancer Research*, 30:1233-1238.