

$$t = r_1 T, y_i = X_i / K, y_3 = X_3 / K, w_1 = A_1 K / r_1, w_2 = B_1 K, w_3 = B_2 K$$

$$w_4 = r_2 / r_1, w_5 = A_2 K / r_1, w_6 = r_3 K / r_1, w_7 = 1 / S_3, w_8 = S_1 K / S_3, w_9 = S_2 K / S_3,$$

$$w_{10} = h_1 / K$$

In the above, for simplicity, we have further assumed that B_i and $S_i, i=1,2$ are in same proportion i.e. $w_8 = \alpha_1 w_2, w_9 = \alpha_2 w_2$.

3. The existence of Positive Equilibrium Point:

The existence of positive equilibrium point is established in the following theorem:

$$y_2 = \frac{w_5}{w_1 w_4} \left(y_1 - \left(1 - \frac{w_1 w_4}{w_5}\right) \right) > 0 \Rightarrow y_1 - \left(1 - \frac{w_1 w_4}{w_5}\right) > 0 \Rightarrow w_5 > w_1 w_4 \quad (5)$$

$$\hat{y}_3 = \frac{w_{10}}{w_6} \left(\frac{1 + \alpha_1 w_2 \hat{y}_1 + \alpha_2 w_3 \hat{y}_2}{1 + \alpha_1 w_2 \hat{y}_1 + \alpha_2 w_3 \hat{y}_2 - w_7} \right) > 0 \text{ when } 1 + \alpha_1 w_2 \hat{y}_1 + \alpha_2 w_3 \hat{y}_2 > w_7$$

Since $0 < \hat{y}_1 < 1, 0 < \hat{y}_2 < 1$, therefore, the system will have a positive equilibrium point. Thus the system will have a positive nonzero solution under condition (3) in this case. This proves the theorem.

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} -\hat{y}_1 + \frac{w_1 w_2 \hat{y}_1 \hat{y}_3}{(1 + w_2 \hat{y}_1 + w_3 \hat{y}_2)^2} & \frac{w_1 w_3 \hat{y}_1 \hat{y}_3}{(1 + w_2 \hat{y}_1 + w_3 \hat{y}_2)^2} & -\frac{w_1 \hat{y}_1}{(1 + w_2 \hat{y}_1 + w_3 \hat{y}_2)} \\ \frac{w_2 w_5 \hat{y}_2 \hat{y}_3}{(1 + w_2 \hat{y}_1 + w_3 \hat{y}_2)^2} & \hat{y}_2 \left(-w_4 + \frac{w_5 w_3 \hat{y}_3}{(1 + w_2 \hat{y}_1 + w_3 \hat{y}_2)^2} \right) & -\frac{w_5 \hat{y}_2}{(1 + w_2 \hat{y}_1 + w_3 \hat{y}_2)} \\ \frac{\alpha_1 w_7 w_2 w_6 \hat{y}_3 \hat{y}_3}{(1 + \alpha_1 w_2 \hat{y}_1 + \alpha_2 w_3 \hat{y}_2)^2} & \frac{\alpha_2 w_7 w_3 w_6 \hat{y}_3 \hat{y}_3}{(1 + \alpha_1 w_2 \hat{y}_1 + \alpha_2 w_3 \hat{y}_2)^2} & w_6 \hat{y}_3 \left(1 - \frac{w_7}{(1 + \alpha_1 w_2 \hat{y}_1 + \alpha_2 w_3 \hat{y}_2)} \right) \end{bmatrix}$$

The characteristic equation of variational matrix is

$$\lambda^3 + a_0 \lambda^2 + a_1 \lambda + a_2 = 0 \quad (6)$$

where

$$a_0 = -(a_{11} + a_{22} + a_{33}); a_1 = (a_{11} a_{33} + a_{22} a_{33} + a_{11} a_{22} - a_{12} a_{21} - a_{23} a_{32} - a_{13} a_{31});$$

$$a_2 = (a_{13} a_{31} a_{22} + a_{23} a_{32} a_{11} + a_{21} a_{33} a_{12} - a_{12} a_{23} a_{31} - a_{13} a_{21} a_{32} - a_{11} a_{22} a_{33}).$$

$$\text{Let } a_{11} = -m_1; a_{22} = -m_2; a_{33} = -m_3; a_{13} = -m_4; a_{23} = -m_5. \text{ Since}$$

$$a_0 = m_1 + m_2 + m_3 > 0; a_1 = (m_1 m_3 + m_2 m_3 + m_1 m_2 - a_{12} a_{21} + m_5 a_{32} + m_4 a_{31}) > 0;$$

$$a_2 = (m_4 a_{31} m_2 + m_5 a_{32} m_1 - a_{21} m_3 a_{12} + a_{12} m_5 a_{31} + m_4 a_{21} a_{32} + m_1 m_2 m_3) > 0.$$

Then applying Routh's criteria $a_0 > 0$ provided $(a_{11} + a_{22} + a_{33}) < 0$, that is, $a_{11} < 0, a_{22} < 0, a_{33} < 0$. Also $a_2 > 0, a_1 > 0$ and $a_1 a_0 - a_2 > 0$. Therefore positive nonzero equilibrium point is locally asymptotically stable. None of the roots of equation (3) is zero as $a_0 \neq 0$.

Theorem 3.1 The system (2) has positive equilibrium point $(\hat{y}_1, \hat{y}_2, \hat{y}_3)$ provided the following conditions are satisfied:

$$w_5 > w_1 w_4 \text{ and } 1 + \alpha_1 w_2 \hat{y}_1 + \alpha_2 w_3 \hat{y}_2 > w_7 \quad (3)$$

Proof. For nonzero equilibrium point, equating the three equations to zero and solving them we get,

$$f_1(y_1, y_2, y_3) = 0; f_2(y_1, y_2, y_3) = 0; f_3(y_1, y_2, y_3) = 0 \quad (4)$$

by solving first two isoclines of (4); we get

4. HOPF'S Analysis of the Harvested Food Web

Assume $y_1 = \hat{y}_1 + u, y_2 = \hat{y}_2 + v, y_3 = \hat{y}_3 + w$, where u, v and w small perturbations are. The variational matrix about $(\hat{y}_1, \hat{y}_2, \hat{y}_3)$ is given by

Substituting $\lambda = \pm i \omega$ into (3), the real and imaginary partitions of the results lead to the following conditions: (i) $\omega = \pm \sqrt{a_1}$ (ii) $a_0 \omega^2 = a_2$
 (i) and (ii) and (3) results that a pair of purely imaginary roots $\pm i \sqrt{a_1}$ and a real root " $-a_0$ ".

Transversality condition: - Let the characteristic equation be such that it contains a real root, say c_1 , and a pair of purely imaginary roots $\lambda_l = \lambda'_l \pm i \lambda'_2$:

$$(\lambda - \lambda_1)(\lambda - \bar{\lambda}_1)(\lambda - c_1) = 0.$$

Or


$$\lambda^3 - (2\lambda'_1 + c_1)\lambda^2 + (|\lambda'_1|^2 + 2\lambda'_1 c_1) - |\lambda'_1|^2 c_1 = 0 \quad (7)$$

Comparing the coefficients of (3) and (4) gives $a_1(-a_0 - 2\lambda'_1) = -a_2 + 2\lambda'_1(2\lambda'_1 + a_0)^2$ (8)


Differentiating (5) with respect to bifurcation parameter w_7 , and substituting $w_7 = w_7^*$ and $\lambda'_1(w_7^*) = 0$ yields:

$$\left. \frac{\partial \lambda'_1}{\partial w_7} \right|_{w_7 = w_7^*} = - \frac{(a_0 \frac{\partial a_1}{\partial w_7} + a_1 \frac{\partial a_0}{\partial w_7} - \frac{\partial a_2}{\partial w_7})}{2(a_0^2 + a_1)} \Rightarrow (9)$$

$$w_1 = 3.3, w_2 = 1.2, w_3 = 1.3, w_4 = 1.1, w_5 = 2.5, w_6 = 1.0, \alpha_1 = 1.5, \alpha_2 = 0.5, w_{10} = 0.08.$$



$w_7 = 1.7$, Asymptotically behavior




$w_7 = 1.5$, Quasiperiodic behavior

$$\left. \frac{\partial \lambda'_1}{\partial w_7} \right|_{w_7 = w_7^*} \neq 0.$$

Thus the transversality condition is satisfied. So there exists a family of periodic solutions bifurcating from non zero equilibrium in the neighborhood of w_7^* , that is, the Hopf bifurcation will occur when $w_7 \in (w_7^* - \delta, w_7^* + \delta)$.

5. Numerical Simulation for HOPF bifurcation

In this section, the numerical analysis of underlying harvested food web is carried out under the biological feasible conditions. It is observed that numerically the other parameters are fixed at biologically feasible values. Only key parameter w_7 is varied.



6. Conclusion

It is concluded that for a set of parameter values with varying key parameter food web harvesting model shows hopf bifurcation. In this paper hopf bifurcation analysis of food web harvesting model is shown analytically as well as numerically.

References

- [1] S. Gakkhar, R. K. Naji, Existence of chaos in two-prey, one-predator system. *Chaos, Solitons & fractals* 17 (2003) 639-649.
- [2] S. Gakkhar, R. K. Naji, Order and chaos in a food web consisting of a predator and two independent preys. *Communications Nonlinear Science And Numerical Simulations* 10 (2005) 105-120.
- [3] S. Gakkhar, B.Singh, Complex dynamic behavior in a food web consisting of two preys and a predator *Chaos, Solitons & fractals.* 24 (2005) 789-801

- [4] H.I Freedman,. and J.W.H So, Global stability and Persistence of simple food chain, Mathematical Bioscience, 76 (1985) 69-86.
- [5] Brahampal Singh and S.Gakkhar,"Analysis of a Food Web Consisting of Two Logistic Prey and a Predator with Modified Leslie- Gower scheme" accepted for publication in "journal of Natural and physical sciences" Gurukula kangari vishwavidyalaya Haridwar, Uttarakhand. ISSN 097 3799.

