(*k*,*d*)–Mean Labeling of Some Family of Trees

B. Gayathri¹, V. Sulochana²

¹PG and Research Department of Mathematics, Periyar E.V.R. College, Tiruchirappalli, India

²Department of Mathematics, Seethalakshmi Ramaswami College, Tiruchirappalli, India

Abstract: Mean labeling of graphs was discussed in [24-25] and the concept of odd mean labeling was introduced in [22]. k-odd mean labeling and (k,d)-odd mean labeling are introduced and discussed in [1,6-8]. k-mean, k-even mean and (k,d)-even mean labeling are introduced (k,d)-mean labeling and we have obtained results for some family of trees.

Keywords: (k,d)-mean labeling, (k,d)-mean graph

1. Introduction

All graphs in this paper are finite, simple and undirected. Terms not defined here are used in the sense of Harary [20]. The symbols V(G) and E(G) will denote the vertex set and edge set of a graph G.

A graph labeling is an assignment of integers to the vertices or edges or both subject to certain conditions. If the domain of the mapping is the set of vertices (or edges) then the labeling is called a vertex labeling (or an edge labeling).

Graph labeling was first introduced in the late 1960's. Many studies in graph labeling refer to Rosa's research in 1967 [23]. Labeled graphs serve as useful models for a broad range of applications such as X-ray, crystallography, radar, coding theory, astronomy, circuit design and communication network addressing. Particularly interesting applications of graph labeling can be found in [2-5].

Mean labeling of graphs was discussed in [24,25]. Vaidya and et al. [28-31] have investigated several new families of mean graphs. Nagarajan and et al. [27] have found some new results on mean graphs.

Ponraj, Jayanthi and Ramya extended the notion of mean labeling to super mean labeling in [21]. Gayathri and Tamilselvi [18-19,26] extended super mean labeling to k-super mean, (k,d)-super mean, k-super edge mean and (k,d)-super edge mean labeling. Manickam and Marudai [22] introduced the concept of odd mean graphs. Gayathri and Amuthavalli [1,6-8] extended this concept to k-odd mean and (k,d)-odd mean graphs. Gayathri and Gopi [9-17] extended this concept to k-mean, k-even mean and (k,d)-even mean graphs.

In this paper, we extend *k*-mean graphs to (k,d)-mean graphs since there are graphs which are (k,d)-mean for all $k \ge 2$ and $d \ge 2$ but not (k,1)-mean for any $k \ge 1$. Here, we have found (k,d)-mean labeling of some family of trees.

Throughout this paper, k and d denote any positive integer greater than or equal to 1.

For brevity, we use (k,d)-ML for (k,d)-mean labeling and (k,d)-MG for (k,d)-mean graph.

2. Main Results

Definition 2.1

A (p,q) graph G is said to have a **mean labeling** if there is an injective function f from the vertices of G to $\{0,1,2,...,q\}$ such that the induced map f^* defined on E by $f^*(uv) = \left| \frac{f(u) + f(v)}{2} \right|$ is a bijection from E to $\{1,2,...,q\}$.

A graph that admits a mean labeling is called a **mean graph**.

Definition 2.2

A (p,q) graph G is said to have a *k***-mean labeling** if there is an injective function f from the vertices of G to $\{0, 1, 2, ..., k+q-1\}$ such that the induced map f^* defined on E by $f^*(uv) = \left| \frac{f(u) + f(v)}{2} \right|$ is a bijection from E to $\{k, k+1, k+2, ..., k+q-1\}$.

A graph that admits a k-mean labeling is called a k-mean graph.

Observation 2.3

Every 1-mean labeling is a mean labeling.

Definition 2.4

A (p,q) graph G is said to have a (k,d)-mean labeling if there exists an injective function f from the vertices of G to $\{0,1,2,...,k+(q-1)d\}$ such that the induced map f^*

defined on *E* by
$$f^*(uv) = \left| \frac{f(u) + f(v)}{2} \right|$$
 is a bijection from
E to $\{k, k+d, k+2d, \dots, k+(q-1)d\}$.

A graph that admits a (k,d)-mean labeling is called a (k,d)-mean graph.

Observation 2.5

- 1) Every (k,1)-mean labeling is a *k*-mean labeling
- 2) Every (1,1)-mean labeling is a mean labeling.

Theorem 2.6

The path graph P_n is a (k,d)-mean graph for all k and d, when n is even.

Proof

Let $V(P_n) = \{v_1, v_2, ..., v_n\}$ and

$$E(P_n) = \{v_i v_{i+1}, 1 \le i \le n-1\}$$
 be denoted as in the Figure 2.1

First we label the vertices as follows:

Define $f: V \to \{0, 1, 2, ..., k + (q-1)d\}$ by

$$f(v_i) = k + d(i-1) - 1$$
, if *i* is odd

 $f(v_i) = k + d(i-2),$ if *i* is even

Then the induced edge labels are

 $f^*(v_i v_{i+1}) = k + d(i-1),$ for $1 \le i \le n-1$.

The above defined function f provides (k,d)-mean labeling of the graph.

So, the path graph P_n is a (k,d)-mean graph for all k and d, when n is even.

(k,d) mean labeling of P_n for different cases of k and d when n is even are shown in illustration 2.7.

Illustration 2.7

(100, 20)-mean labeling of the graph P_{10} is shown in Figure 2.2 100 120 140 160 180 200 220 240 260

(55, 44)-mean labeling of the graph
$$P_6$$
 is shown in Figure 2.3
 $55 99 143 187 231 54 55 142 143 230 231$
Figure 2.3: (55, 44)-*ML* of P_6

(33, 11)-mean labeling of the graph P_8 is shown in Figure 2.4 $\begin{array}{r} 33 \quad 44 \quad 55 \quad 66 \quad 77 \quad 88 \quad 99 \\
\hline
32 \quad 33 \quad 54 \quad 55 \quad 76 \quad 77 \quad 98 \quad 99 \\
\hline
Figure 2.4: (33, 11)-ML \text{ of } P_8
\end{array}$

(60, 13)-mean labeling of the graph P_{10} is shown in Figure 2.5 <u>60</u> 73 86 99 112 125 138 151 164 59 60 85 86 111 112 137 138 163 164 **Figure 2.5:** (60, 13)-ML of P_{10}

Definition 2.8

A **comb graph** P_n^+ is a tree obtained from a path by attaching exactly one pendant edge to each vertex of the path.

Theorem 2.9

The comb graph P_n^+ is a (k,d)-mean graph for all k and d. **Proof**

Let
$$V(P_n^+) = \{u_1, u_2, ..., u_n, u_1, u_2, ..., u_n\}$$





First we label the vertices as follows:

Define
$$f: V \to \{0, 1, 2, ..., k + (q-1)d\}$$
 by
 $f(u_1) = k$
 $f(u_1) = k - 1$
 $f(u_n) = k + (q-1)d$
 $f(u_n) = k + (q-1)d - 1$
 $f(u_i) = k + 2d(i-1), \text{ for } 2 \le i \le n-1$
 $f(u_i) = k + 2d(i-1) - 1, \text{ for } 2 \le i \le n-1$

Then the induced edge labels are

$$f^*(u_i u_i) = k + 2d(i-1), \quad \text{for } 1 \le i \le n$$
$$f^*(u_i u_{i+1}) = k + d(2i-1), \quad \text{for } 1 \le i \le n-1$$

The above defined function f provides (k,d)-mean labeling of the graph. So, the graph P_n^+ is a (k,d)-mean graph for all k and d. (k,d)-mean labeling of P_n^+ for different cases of k and d are shown in illustration 2.10

Illustration 2.10

(4,6)-mean labeling of the graph P_5^+ is shown in Figure 2.7



(13,3)-mean labeling of the graph P_5^+ is shown in Figure 2.8





(23,8)-mean labeling of the graph
$$P_8^+$$
 is shown in Figure 2.9



Volume 5 Issue 1, January 2016 www.ijsr.net

Licensed Under Creative Commons Attribution CC BY

International Journal of Science and Research (IJSR) ISSN (Online): 2319-7064 Index Copernicus Value (2013): 6.14 | Impact Factor (2014): 5.611



Definition 2.11

A **twig** is a tree obtained from a path by attaching exactly two pendant edges to each internal vertex of the path.

Theorem 2.12

The twig graph T_n is a (k,d)-mean graph for all k and d, when n is even.

Proof

Let $V[T_n] = \{v_i, 0 \le i \le n-1, u_i, w_i, 1 \le i \le n-2\}$

and $E(T_n) = \{v_i u_i, v_i w_i, 1 \le i \le n-2 \text{ and } v_i v_{i+1}, 0 \le i \le n-2\}$ be denoted as in the Figure 2.11.



Figure 2.11: Ordinary labeling of T_n

First we label the vertices as follows:

Define
$$f: V \rightarrow \{0, 1, 2, ..., k + (q-1)d\}$$
 by

$$f(v_i) = \begin{cases} k+3di-1, & \text{if } i \text{ is even} \\ k+3d(i-1), & \text{if } i \text{ is odd} \end{cases}$$

$$f(u_i) = \begin{cases} k+d(3i-4)+1, & \text{if } i \text{ is even} \\ k+d(3i-1), & \text{if } i \text{ is odd} \end{cases}$$

$$f(w_i) = \begin{cases} k+d(3i-2)+1, & \text{if } i \text{ is even} \\ k+d(3i+1), & \text{if } i \text{ is odd} \end{cases}$$
Then the induced edge length are

Then the induced edge labels are

 $f^{*}(v_{i}v_{i+1}) = k + 3di, \quad \text{for } 0 \le i \le n-2$ $f^{*}(v_{i}u_{i}) = k + d(3i-2), \text{ for } 1 \le i \le n-2$ $f^{*}(v_{i}w_{i}) = k + d(3i-1), \text{ for } 1 \le i \le n-2$

The above defined function f provides (k,d)-mean labeling of the graph.

So, the twig graph T_n is a (k,d)-mean graph for all k and d, when n is even.

(k,d)-mean labeling of T_n for different cases of k and d when n is even are shown in illustration 2.13.

Illustration 2.13

(6,6)-mean labeling of the graph T_8 is shown in Figure 2.12



Figure 2.12: (6,6)–ML of T_8 (3,2)–mean labeling of the graph T_4 is shown in Figure 2.13





(5,3)-mean labeling of the graph T_6 is shown in Figure 2.14



Figure 2.14: (5,3)–*ML* of *T*₆

(8,5)-mean labeling of the graph T_{10} is shown in Figure 2.15



Theorem 2.14

The star $K_{1,n}$ $(n \ge 4)$ is a (k,d)-mean graph for all $k \ge n-2$ and for all *d* satisfying $(q-1)d \le k+1$ except when *n* is odd and *d* is even.

Proof

Let $V(K_{1,n}) = \{u, v_1, v_2, ..., v_n\}$ and

$$E(K_{1,n}) = \{uv_i, 1 \le i \le n\}$$
 be denoted as in Figure 2.16.



Figure 2.16: Ordinary labeling of $K_{1,n}$

First we label the vertices as follows:

Define $f: V \to \{0, 1, 2, ..., k + (q-1)d\}$ by Case (1): when *n* is even Subcase (i): d is odd f(u) = k + (q-1)d - 1 $f(v_i) = k - (q-1)d + 2i - 1,$ for $1 \le i \le \frac{n}{2} - 1$ for $i = \frac{n}{2}$ $f(v_i) = k,$ $f(v_i) = k + (q-1)d + (i-n) - 1$, for $\frac{n}{2} + 1 \le i \le n-1$ $f(v_n) = k + (q-1)d$ Subcase (ii): d is even f(u) = k + (q-1)d - 1 $f(v_i) = k - (q-1)d + 2i - 1,$ for $1 \le i \le \frac{n}{2} - 1$ for $i = \frac{n}{2}$ $f(v_i) = k - 1,$ $f(v_i) = k + (q-1)d + (i-n) - 2$, for $\frac{n}{2} + 1 \le i \le n-1$ $f(v_n) = k + (q-1)d$ Then the induced edge labels are $f^*(uv_i) = k + d(i-1),$ for $1 \le i \le n$

Case (2): when *n* is odd subcase (i): *d* is odd f(u) = k + (q-1)d - 1 $f(v_i) = k - (q-1)d + 2i - 1$, for $1 \le i \le \frac{n-1}{2}$ $f(v_i) = k + d$, for $i = \frac{n+1}{2}$ $f(v_i) = k + (q-1)d + (i-n) - 1$, for $\frac{n+3}{2} \le i \le n-1$ $f(v_n) = k + (q-1)d$ Then the induced edge labels are $f^*(uv_i) = k + d(i-1)$, for $1 \le i \le n$

The above defined function f provides (k,d)-mean labeling of the graph.

So, the star $K_{1,n}$ is a (k,d)-mean graph for all $k \ge n-2$ and for all d satisfying $(q-1)d \le k+1$ except when n is odd and d is even.

(k,d)-mean labeling of $K_{1,n}$ for different cases of k and d except when n is odd and d is even are shown in illustration 2.15.

Illustration 2.15

(6,1)-mean labeling of the graph $K_{1,8}$ is shown in Figure 2.17



(3,1)-mean labeling of the graph $K_{1,5}$ is shown in Figure 2.18













Definition 2.16

A **bistar** $B_{m,n}$ is a tree obtained by joining the center vertices of the copies of $K_{1,m}$ and $K_{1,n}$ with an edge.

Theorem 2.17

The Bistar $B_{n,n}$ ($n \ge 2$) is a (k,d)-mean graph for all k and d.

Proof

Let $V(B_{n,n}) = \{u, v, u_1, u_2, ..., u_n, v_1, v_2, ..., v_n\}$

Licensed Under Creative Commons Attribution CC BY



Figure 2.21: Ordinary labeling of B_n

First we label the vertices as follows:

Define $f: V \to \{0, 1, 2, ..., k + (q-1)d\}$ by

$$f(u) = k + (q-1)d$$

$$f(u_i) = k + 2di - 1, \quad \text{for } 1 \le i \le n-1$$

$$f(u_n) = k + (q-1)d - 1$$

$$f(v) = k - 1$$

$$f(v_i) = k + 2d(i-1), \quad \text{for } 1 \le i \le n$$
Then the induced edge labels are
$$f^*(vv_i) = k + d(i-1), \quad \text{for } 1 \le i \le n$$

$$f^*(uv) = k + nd$$

$$f^*(uu_i) = k + (n+i)d, \quad \text{for } 1 \le i \le n$$

The above defined function f provides (k,d)-mean labeling of the graph.

So, the graph $B_{n,n}$ is a (k,d)-mean graph for all k and d. (k,d)-mean labeling of $B_{n,n}$ for different cases of k and d are shown in illustration 2.18.

Illustration 2.18

(6,7)-mean labeling of the graph $B_{7,7}$ is shown in Figure 2.22.





(3,5)-mean labeling of the graph $B_{5,5}$ is shown in Figure 2.23



(5,7)-mean labeling of the graph $B_{6,6}$ is shown in Figure 2.24



Figure 2.24: (5,7)–*ML* of *B*_{6,6}





Figure 2.25: (4,4)–*ML* of *B*_{8.8}

Theorem 2.19

The Bistar $B_{n,n+1}$ $(n \ge 2)$ is a (k,d)-mean graph for all k and for all $d \le k + 1$.

Proof

Let $V(B_{n,n+1}) = \{u, v, u_1, u_2, ..., u_n, v_1, v_2, ..., v_{n+1}\}$ and $E(B_{n,n+1}) = \{uv, uu_i, 1 \le i \le n, vv_i, 1 \le i \le n+1\}$ be denoted as in the Figure 2.26.



Figure 2.26: Ordinary labeling of $B_{n,n+1}$

First we label the vertices as follows: Define $f: V \to \{0, 1, 2, ..., k + (q-1)d\}$ by

$$f(u) = k - d + 1$$

$$f(u_{1}) = k + d - 2$$

$$f(u_{i}) = k + d(2i - 1) - 1, \quad \text{for } 2 \le i \le n$$

$$f(v) = k + (q - 1)d - 1$$

$$f(v_{i}) = k + d(2i - 1), \quad \text{for } 1 \le i \le n + 1$$
Then the induced edge labels are
$$f^{*}(uu_{i}) = k + d(i - 1), \quad \text{for } 1 \le i \le n$$

$$f^{*}(uv) = k + nd$$

$$f^{*}(vv_{i}) = k + (n + i)d, \quad \text{for } 1 \le i \le n + 1$$

Volume 5 Issue 1, January 2016 www.ijsr.net

The above defined function f provides (k,d)-mean labeling of the graph.

So, the graph $B_{n,n+1}$ is a (k,d)-mean graph for all k and for all $d \le k+1$.

(k,d)-mean labeling of $B_{n,n+1}$ for different cases of k and $d \le k + 1$ are shown in illustration 2.20.

Illustration 2.20

(4,5)-mean labeling of the graph $B_{5,6}$ is shown in Figure 2.27





(1,2)-mean labeling of the graph $B_{3,4}$ is shown in Figure 2.28



Figure 2.28: (1,2)–ML of $B_{3,4}$ (3,3)–mean labeling of the graph $B_{4,5}$ is shown in Figure 2.29



Figure 2.29: (3,3)–ML of $B_{4,5}$ (6,6)–mean labeling of the graph $B_{6,7}$ is shown in Figure 2.30



Figure 2.30: (6,6)–ML of *B*_{6,7}

References

- K. Amuthavalli, Graph labeling and its Applications some generalizations of odd mean labeling, Ph.D. Thesis, Mother Theresa Women's University, Kodaikanal, July (2010).
- [2] G.S. Bloom, S.W. Golomb, Applications of numbered undirected graphs, Proc. IEEE, 65 (1977), 562-570.
- [3] G.S. Bloom, S.W. Golomb, Numbered complete graphs unusual rulers and assorted applications, Theory and Applications of Graphs-Lecture notes in Math., Springer Verlag, New York, 642 (1978), 53-65.
- [4] G.S. Bloom, D.F. Hsu, On graceful digraphs and a problem in network addressing, Congressus Numerantium, 35 (1982) 91-103.
- [5] J.A. Gallian, A dynamic survey of graph labeling, Electronic Journal of Combinatorics, 18 (2014) # DS6.
- [6] B. Gayathri and K. Amuthavalli, *k*-odd mean labeling of crown graphs, International Journal of Mathematics and Computer Science, 2(3) (2007) 253-259.
- [7] B. Gayathri and K. Amuthavalli, (k,d)-odd mean labeling of some graphs, Bulletin of Pure and Applied Sciences, 26E(2) (2007) 263-267.
- [8] B. Gayathri and K. Amuthavalli, k-odd mean labeling of $\langle K_{1,n}, K_{1,m} \rangle$, Acta Ciencia Indica, 34(2) (2008) 827-834.
- [9] B. Gayathri and R. Gopi, *k*-even mean labeling of $D_{m,n}@C_n$, International Journal of Engineering Science, Advanced Computing and Bio-Technology, Vol. 1, No. 3, July-September 2010, 137 -145.
- [10] B. Gayathri and R. Gopi, k-even mean labeling of D_{m,n}, Acta Ciencia Indica, Vol. XXXVII, No. 2, 291-300, 2011.
- [11] B. Gayathri and R. Gopi, *k*-even mean labeling of $C_n \cup P_m$, Elixir International Journal of Applied Sciences, No. 36, 2011, 3102-3105.
- [12] B. Gayathri and R. Gopi, k-even mean labeling of $T_{n,m,t}$, International Journal of Engineering Science, Advanced Computing and Bio-Technology, Vol. 2, No. 2, April-June 2011, 77-87.
- [13] B. Gayathri and R. Gopi, International Conference on Mathematics and Computer Science, k-even mean labeling of $C_m \in \overline{K}_n$, Loyola College, Chennai, January 7-8, 2011, Proc. 519-524.

- [14] B. Gayathri and R. Gopi, (k,d)-even mean labeling of P_m e nK_1 , International Journal of Mathematics and soft computing, Vol. 1, No. 1, 17-23, August 2011.
- [15] B. Gayathri and R. Gopi, k-even mean labeling of some graphs, Heber International Conference on Applications of Mathematics and Statistics, Bishop Heber College (Autonomous), January 5-7, 2012 Proc. 420-425.
- [16] B. Gayathri and R. Gopi, International Conference on Mathematics in Engineering and Business Management, *k*-even mean labeling of some trees, Stella Maris College (Autonomous), Chennai, March 9-10, 2012, Proc. 99-102.
- [17] R. Gopi, A Study on Different kinds of Mean Labeling, Ph.D. Thesis, Bharathidasan University, Trichy, February (2013).
- [18] B. Gayathri, M. Tamilselvi and M. Duraisamy, (k,d)-super mean labeling of some graphs, International Journal of Mathematics and Computer Science, 2(3) (2007) 245 252.
- [19] B. Gayathri and M. Tamilselvi, *k*-super mean labeling of some trees and cycle related graphs, Bulletin of Pure and Applied Sciences, Volume 26E(2) (2007) 303-311.
- [20] F. Harary Graph Theory, Addision-Wesley, Reading Masaachusetts, 1972.
- [21] R. Ponraj, Jayanthi and D. Ramya, On super mean graphs of order ≤ 5, Bulletin of Pure and Applied Sciences, 25E (2006) 143-148.
- [22] K. Manickam and M. Marudai, Odd mean labeling of graphs, Bulletin of Pure and Applied Sciences, 25E(1) (2006) 149-153.
- [23] Rosa, On certain valuations of the vertices of a graph Theory of Graphs (Internet Symposium, Rome, July 1966), Gordon and Breach, N.Y. and Duhod, Paris (1967) 349-355.
- [24] S. Somasundaram and R. Ponraj, Mean labeling of graphs, National Academy Science Letter, 26 (7-8) (2003), 10-13.
- [25] S. Somasundaram and R. Ponraj, Non-existence of mean labeling for a wheel, Bull. Pure and Appl. Sciences (Mathematics & Statistics), 22E (2003) 103-111.
- [26] M. Tamilselvi, A study in Graph Theory-Generalization of super mean labeling, Ph.D. Thesis, Vinayaka Mission University, Salem, August (2011).
- [27] R. Vasuki and A. Nagarajan, Further results on mean graphs, Scientia Magna, 6(2) (2010) 26-39.
- [28] S.K. Vaidya and Lekha Bijukumar, Mean labeling for some new families of graphs, Journal of Pure and Applied Sciences, 1(18) (2010) 115-116.
- [29] S.K. Vaidya and Lekha Bijukumar, Some new families of mean graphs, Journal of Mathematics Research, 2(3) (2010) 169-176.
- [30] S.K. Vaidya and Lekha Bijukumar, New mean graphs, International J. Math. Combin., 3 (2011) 107-113.
- [31] S.K. Vaidya and Kanani, Some new mean graphs, International Journal of Information Science and Computer Mathematics, 1(1) (2010) 73-80.