

# Effects of Ingestion of Bonny Light Crude Oil on Sperm Motility and Morphology of Male Wistar Rats

Victor A. Fischer<sup>1</sup>, Christie E. Fischer<sup>2</sup>, Mfon Akpaso<sup>3</sup>, Patrick S. Igbigbi<sup>4</sup>

<sup>1,2,3</sup>Department of Anatomical Sciences, Faculty of Basic Medical Sciences, University of Calabar, Calabar- Nigeria

<sup>4</sup>Department of Anatomy and Cell Biology, Delta State University, Abraka, Nigeria

**Abstract:** This study evaluated the effects of Bonny light crude oil (BLCO) on the sperm morphology and sperm motility of young adult male wistar rats. Eighty (80) adult male wistar rats were used for this study, they were divided into four (4) parallel groups A, B, C, and D consisting each of fifteen (15) experimental animals and five (5) control animals. The experimental animals in groups A, B, C, and D received 0.5, 1.5, 2.5, and 3.5ml/kg B.W of BLCO respectively, while the control animals in corresponding groups received the corresponding doses of normal saline via oral gastric intubation for sixty (60) days alternately. At the end of the sixty days, the animals were sacrificed and the spermatozoa was expressed, a homogenate made and used for semen analysis. The percentage motility for normal motility decreased down the group with increase in administered BLCO (A; 79.30%, B; 44.00%, C; 38.00%, D; 15.80%), while in the control groups, the percentage for both normal motility (A; 90.60%, B; 90.50%, C; 90.10%, D; 89.60%) were fairly stable. The total percentage morphology of the defected sperm increased down the group (A; 23.50%, B; 46.80%, C; 59.90%, D; 70.20%) with increase in BLCO administered, but despite the varying volume of normal saline administered to each control group, the percentage morphology of normal sperm (A; 93.80%, B; 94.60%, C; 94.20%, D; 93.00%) remained fairly the stable. This study shows that administration of BLCO, can cause a reduction in the percentage of motile sperm and also abnormal morphology of sperm cells.

**Keywords:** Crude oil, sperm motility, sperm morphology, wistar rats.

## 1. Introduction

Chemically, crude oil is defined as a complex mixture of thousands of different chemical components, straight and branched chain paraffin, cycloparaffin, aromatic and polynuclear aromatic hydrocarbon, small amounts of sulphur and nitrogen compounds also with trace amounts of metals (Hawley, 1981). Bonny light crude oil (BLCO), is Nigeria's marker crude oil, preferred for its low sulphur content which makes it less corrosive to refinery infrastructures and also for its low specific gravity.

Much dependence on petroleum as a whole has brought with it, cases of environmental pollution, and detrimental effects on human existence and ecosystem (National Research Council, 1985; Ruddel, 1994). Oil spillage during production, exploration and discharge from storage facilities and refineries and bursting of pipelines, is one of the fundamental sources of hazard and pollution in oil producing communities of Nigeria (Dede and Kagbo, 2002; Peters, 1993; Okereke and Ezeanyina, 1987; Lambert akhonbare and Shaw, 1982).

The devastating consequences of spills of this crude oil in Niger delta region, with its eventual hazards on both aerial and terrestrial environs tantamount to an irreversible chain effect on the biodiversity and human safety (Akpofure *et al.*, 2000). Ingestion of crude oil and its products either in the raw or bioaccumulated form in marine life, presents a potential hazard to terrestrial species (Shore and Douben 1994). Studies revealed that there are biochemical and cytotoxic impairment associated with ingestion of marine animals exposed to crude oil polluted water (Eyong, 2000; Eyong *et al.*, 2004). Crude oil is ingested as laxative, anti-

poisoning agent, anti-convulsion agent, used for treatment of arthritis, snake antidote, especially in rural areas, where the conventional antidotes are not available.

The increasing trend of male reproductive impairment observed in some countries has been associated with possible exposure to chemicals that could interfere with endocrine homeostasis (endocrine reproductive chemicals EDC) (Bergstrom *et al.*, 1996; Moller, 2001). Crude oil is an important environmental industrial pollutant that is composed of a widely depending on the location and source (Briggs *et al.*, 1996). In our environment, these chemicals are capable of mimicking the inherent actions of reproductive hormones and thence, have the ability to disrupt the neuroendocrine system or function of gonads directly (Colborn *et al.*, 1993).

The result of this study will thus provide information on the effect of ingestion of BLCO on the motility of sperms and morphology of sperms of male wistar rats.

## 2. Materials and Methods

### 2.1 Bonny Light Crude Oil

The Bonny light crude oil used for this study was obtained from shell petroleum development company, port Harcourt, Rivers state with permission from the Department of petroleum resources, NNPC, Lagos, Nigeria.

### 2.2 Experimental Animals

Eighty (80) male albino rats weighing between 150- 200g and obtained from the Department of pharmacology,

University of Calabar, were used for this study. The rats were given feed and water ad libitum, they were allowed to acclimatize for a period of three (3) weeks in the animal house of the Department of human anatomy prior to experimentation, they were kept in well ventilated cages at a room temperature of about 25±5°C, and 12 hour light and dark cycle.

### 2.3 Experimental Design

The design consisted of eighty (80) rats divided into four parallel groups consisting of Bonny light treated fifteen (15) animals and five (5) control animals in each pair. The dosages of BLCO administered were determined from previous studies (Eyong, 2000; Fischer *et al.*, 2007). The animals in groups A, B, C, and D were given 0.5, 1.5, 2.5 and 3.5ml/kg B.W of BLCO respectively, while the control animals received corresponding doses of normal saline both once on alternate days for sixty (60) days via oral gastric intubation. At the end of the sixty days, the animals were euthanized under chloroform vapour and sacrificed, spermatozoa was expressed out, a homogenate made which was then used for the semen analysis.

### 2.4 Statistical Analysis

Analysis of variance (ANOVA) was used to compare results from the experimental groups and control groups. P values <0.05 were said to be statistical significant.

## 3. Results

### 3.1 Sperm Motility

Table 1 below represents the mean percentage motility of sperm cells that exhibited normal and retarded motility. The percentage of sperm cells showing normal motility decreased down the group (A; 79.30%, B; 44.00%, C; 38.00%, D; 15.80%) with increase in the administered dose of BLCO. The percentage motility of sperm cells showing retarded motility increased down the group (A; 20.70%, B; 56.00%, C; 62.00%, D; 84.20%) with increase in administered dose of BLCO. In the control groups, the percentage motility for both the normal motility (A<sub>1</sub>;

90.60%, B<sub>1</sub>; 90.50%, C<sub>1</sub>; 90.10%, D<sub>1</sub>; 89.60%) and retarded motility (A<sub>1</sub>; 9.40%, B<sub>1</sub>; 9.50%, C<sub>1</sub>; 9.90%, D<sub>1</sub>; 10.40%) fairly stable down the group despite varying doses of normal saline administered.

**Table 1: Mean percentage motility of sperm cells**

Groups	Retarded motility			Normal motility		
	1+ (%)	2+ (%)	Total (%)	3+ (%)	4+ (%)	Total (%)
Control A <sub>1</sub>	1.60	7.80	9.40	30.60	60.00	90.60
Treated A	6.70	14.00	20.70	28.00	51.30	79.30
Control B <sub>1</sub>	1.50	8.00	9.50	31.30	59.20	90.50
Treated B	29.80	26.20	56.00	22.80	21.20	44.00
Control C <sub>1</sub>	1.80	8.10	9.90	30.10	60.00	90.10
Treated C	33.30	28.70	62.00	24.60	13.40	38.00
Control D <sub>1</sub>	1.40	9.00	10.40	32.50	57.10	89.60
Treated D	69.40	14.80	84.20	10.40	5.40	15.80

Values are presented as mean %

1+: Spermatozoa moving but not forward progression

2+: Spermatozoa moving aimlessly with slow forward progression

3+: Spermatozoa moving at moderate speed with forward progression

4+: Spermatozoa moving at high speed with straight line forward progression

### 3.2 Sperm Morphology

Table 2 shows the mean percentage morphology the sperms in each group. The percentage morphology of the normal sperm decreased down the group (A; 76.50%, B; 53.20%, C; 40.10%, D; 29.80%) as the dose of BLCO administered increased, while the total percentage of morphology of the defected sperm increased down the group (A; 23.50%, B; 46.80%, C; 59.90%, D; 70.20%), for each form of defect, a gradual increase in the number of defected sperms down the group with respect to increase in BLCO administered. But despite the varying volume of normal saline administered to each control group, the percentage morphology of normal sperm (A<sub>1</sub>; 93.80%, B<sub>1</sub>; 94.60%, C<sub>1</sub>; 94.20%, D<sub>1</sub>; 93.00%) and defected sperm (A<sub>1</sub>; 6.20%, B<sub>1</sub>; 5.40%, C<sub>1</sub>; 5.80%, D<sub>1</sub>; 7.00%) remained fairly stable.

**Table 2: Mean Percentage Morphology of Sperm Cells**

Group	normal sperm NHAT(%)	defected sperm				
		IH(%)	HOD(%)	TD(%)	FS(%)	TOTAL(%)
Control A <sub>1</sub>	93.80	1.80	2.00	2.20	0.20	6.20
Treated A	76.50	4.80	4.60	4.70	10.40	24.00
Control B <sub>1</sub>	94.60	1.20	2.10	2.10	0.00	5.40
Treated B	53.20	11.60	9.50	10.80	14.90	46.80
Control C <sub>1</sub>	94.20	1.90	1.80	2.00	0.10	5.80
Treated C	40.10	14.50	12.70	16.30	16.40	59.90
Control D <sub>1</sub>	93.00	1.80	2.20	2.80	0.20	7.00
Treated D	29.80	16.80	12.50	20.80	20.10	90.20

Values are presented as percentage, NHAT= normal head and tail, HOD = head only defect, IH = isolated head, TD= tail defect, FS= fused sperm.

## 4. Discussion

Several researches carried out in both laboratory and non-laboratory animals with crude oils of different geologic

origins have pointed out the toxic effects of this important commodity (Ellenton and Hallet, 1981; Mckee *et al.*, 1994; Eyong, 2000; Didia *et al.*, 2003). For normal fertility, a man requires normal spermatogenesis, successful epididymal

storage, and normal sperm transport. The fertility in males depends on normal linear progressive sperm motility and normal morphology. This study showed a significant dose dependent reduction in the percentage of normal motility and normal morphology of sperm cells of rats exposed to BLCO, this is an indication that crude oil interfered with testicular spermatogenesis.

It was proposed that environmental exposure to harmful compounds may cause a variation in sperm quality. Crude oil is a known environmental pollutant and contains polyaromatic hydrocarbons (PAH's) which is a known toxic compound. In this study, the most predominant abnormality was fused sperm 10.40% (least dose) and 20.10% (highest dose) and tail defect 4.70%(least dose), and 20.80 (highest dose), it could be suggested that crude oil decreases the synthesis of sperm membrane coating protein which in turn results in the production of morphologically abnormal sperms (Wang *et al.*, 1997, Dada *et al.*, 2001).

Increase in the number of morphologically abnormal sperms results in impaired motility as normal intact sperm morphology is a prerequisite for high speed with straight line forward progressive motility. It is postulated that sperm function is strictly correlated with sperm morphology and that sperm motility is the least predictor of fertility potential in man (Gandini *et al.*,2000).

## 5. Conclusion

Ingestion of BLCO by male wistar rats showed significant reduction in percentage sperm motility and sperm morphology.

## References

- [1] Agency for Toxic Substances and Disease Registry (ATSDR). Toxicological Profile for Total Petroleum Hydrocarbons (TPH). Atlanta, GA: US Department of Health and Human Services, Public Health Service (2004) pp 1-5
- [2] Akpofure, EA, Efere, ML and Prosper, A. Integrated grass root post-impact assessment of acute damaging effects of continuous oil spill in Niger Delta (January 1998-2000). *Oil spill intelligence report.* (2000) 1-9.
- [3] Ashiru O and Blake E. Stimulation of androgenous FSH release during oestrus by
- [4] administration of exogenous FSH or LH at proestrus in the Phenobarbital-blocked rat.
- [5] *Endocrinology* (1979) 105:1162-1167
- [6] Bergmann M, Belire HM, Niechlag E. Serum FSH and testicular morphology in male infertility. *Clinical endocrinology.* (1999) 40 133-136
- [7] Bergstrom R, Adam H, Mohnner M, Zatonski W, Storm H, Ekbohm A, Tretli S, Teppo L, Akre O, Hakulinen T. Increase in testicular cancer in six European countries. A birth cohort phenomenon. *J Natl Cancer Inst.* (1996) 88 727-733
- [8] Briggs KT, Yoshida SH, Gershwin ME. The influence of petrochemicals and stress on the immune system of seabirds. *Regul Toxicol Pharmacol.* (1996) 23, 145-155.

- [9] Colborn T, Vom Saal FS, Soto AM. Developmental effects of endocrine disrupting chemicals in wildlife and humans. *Environ Health Perspect.* (1993) 101: 378-384
- [10] Dede EB and Kagbo HD. A study on the acute toxicological effects of commercial diesel fuel in Nigeria on rats (*Ratus ratus*) using haematological parameters. *J. Appl. Sc. Environ. Mgt.* (2002) 5-9
- [11] Dede EB, Igboh NM and Ayalogu OA. Chronic Study of the Effect of Crude petroleum (Bonny Light), Kerosine and Gasoline on Rats Using Haematological Parameters. *Journal of applied Sciences and environmental management.* (2002) Vol.6 (1) 60-63.
- [12] Dede, EB and Kagbo, HD. A study on the acute toxicological side effects of commercial diesel fuel in Nigeria on rats (*Rattus rattus*) using haematological parameters. *J. Appl.Sc. Environ. Mgt.*, (2002) 5-9.
- [13] Didia BL, Dede EB and Dapper DV. Effects of crude oil contaminated water on haematocrit and histopathology of Guinea pig: Animal mode for investigating crude oil pollution. *Journal of experimental and clinical anatomy.* (2003) 2 (2): 6-11.
- [14] Environmental Health Criteria (EHC) Selected petroleum products. United Nations Environment programme. *World Health Organisation publication, Geneva.* (1982) 20:
- [15] 5- 118.
- [16] Eyong, E U. Biochemical and toxicological implication of ingestion by rats of shelfish exposed to crude oil polluted. Ph. D. thesis, *University of Calabar, Nigeria.* (2000)
- [17] Eyong, E.U.; Umoh, I.B.; Ebong, P.E.; Eteng, M.U.; Antai, A.B. and Akpa, A.O. Haematotoxic effects following ingestion of crude oil. *Nigeria journal of physiological science* (2004) 1&2: 1-6
- [18] Feuston MH and Hamilton CE. Developmental toxicity study in rats exposed dermally to clarified slurry oil for a limited period of gestation. *J. Toxicol – Environ Health* (1997) 49 (2) 207-220.
- [19] Feuston MH and Mackerer CR. Chronic Dermal Studies of petroleum streams in mice. *Fundamental Application to Toxicology* (1996) 301 (1): 47-54.
- [20] Fischer VA, Akpuaka FC, Ngokere AA. Effects of Bonny Light Crude Oil on the histology of cerebral cortex in fetuses of wistar rats *J.Expt. and Clin. Anat.* (2005) 4(1) 9-11.
- [21] Fischer VA, Anibeze CIP, Igiri AO, Ekanem TB, Mesembe OE, Fischer CE. Effect of maternal administration of Bonny Light Crude Oil on brain dimensions of wistar rat fetuses. *Global Journal of Pure and Applied Sciences.* (2007) 13 (1)95-97
- [22] Fischer VA, Anibeze CIP, Igiri AO, Eyong EO, Mesembe OE, Fischer CE. Effects of Bonny Light Crude Oil on Morphology of litters of wistar rats. *Global Journal of Medical Sciences.* (2006) 5(1)51-53
- [23] Harty WR and Chaman EV. A toxicological assessment of unleaded gasoline contamination of drinking water in petroleum contaminated soils vol. 3. Kosteck PT Calabrese EJ (ed) *Lewis publication inc.* (1990) 324-340
- [24] Khan, S; Codner, M; Payne J.;Rahimtula, A. Effect of prudhoe Bay crude oil on hepatic and renal peroxisomal beta-oxidation and mixed function oxidize

- activities in rats. *Carcinogenesis*. (1999) 10 (2): 269-272.
- [25] Lambert AA. and Shaw H. Environmental effects of crude oil spillage. *Global journal of pure and applied sciences*. (1982) 1:85-90.
- [26] McKee RH, Amoroso MA, Freeman JJ. Evaluation of the genetic toxicity of middle distillate fuels. *Environmental and Molecular Mutagenesis*. (1994) 23:234 – 238
- [27] Moline JM, Golden AL, Bar-chama N, Smith E, Rauch ME, Chapin RE, Perreault SD, Schrader SM, Suk WA, Landrigan PJ. Exposure to hazardous substances and male reproductive health. A research framework. *Environmental health perspectives* (2000) 108:9 803-813
- [28] Moller H. (2001) Trends in incidence of testicular cancer and prostate cancer in Denmark. *Hum Reprod* 16 1007-1011
- [29] Obidike R, Igwebuike UM and Shoyinka SV. Testicular morphology and epididymal sperm reserves of male rats exposed to Nigerian Qua Iboe Brent crude oil. *Journal of pure and applied sciences* (1987) 1: 85-90.
- [30] Okereke, C. and Ezeanyina, I. Environmental side effects of crude oil spillage. *Global journal of pure and applied sciences*. (1987) 1:85-90.
- [31] Orisakwe OF, Afonne OJ, Nwobode E, Asomugha L, Dioka CE. (2004) Testicular toxicity of Nigeria's bonny light crude oil in male albino rats *Reprod toxicol* 18: 439-442
- [32] Orth, J.M.; Gunsalus, G.L.; Lamperti, A.A. Evidence from sertoli cells depleted rats indicate that spermatid number in adults depend on number of sertoli cells produced during prenatal development. *Endocrinology*(1988) 127 pp787 – 794
- [33] Pasquini R, Taningher M, Monarca S, Pala M, Angeli G. Chemical composition and genotoxicity of petroleum derivatives collected in two working environments. *J. Toxical Environ Health*. (1989) 27 (2) 225-238
- [34] Peters, SW. Crises in our environment. Nigeria Environmental Education and Management. *University of Calabar Press*. (1993) 126-148.
- [35] Steger, K; Rey, R; Louis, F; Kleisch, S; Behre, H. M; Nieschlag, E. Reversion of the differentiated phenotype and maturation block in sertoli cells in pathological human testis. *Human Reproduction*(1999) 14 pp 136-143
- [36] Walkinson RH and Holt MS. Biodegradability of produced water effluents; medicinal problems in the off-shore oil industry. Proc. Int. conference of institute of petroleum microbiology committee. Amerdeem. *John Wiley and Sons*. (1987) 165-174.