# A Food Chain Model with Strong Allee Effect in Prey

## Nijamuddin Ali

Department of Mathematics, Katwa College (B.Ed. Section), Dist-Burdwan, West Bengal, India, 713130.E

Abstract: In recent years, there has been an intense focus on the construction of mathematical models involving certain interesting ecological phenomena like Allee effect. Many animal and plant species suffer a decrease of the per capita rate of increase as their populations reach small sizes or low densities. In this article, a three species food chain model is proposed and analysed where prey species is affected by strong Allee effect. I have studied the stability behaviour of the system and bifurcation analysis. The system exhibits saddle-node bifurcation curve. Extensive Numerical simulation results are given to support the validity of the theoretical results. The ecological implications of our analytic and numerical findings are discussed.

Keywords: Food chain model, Allee effect, Stability, Bifurcation, Numerical simulation

## 1. Introduction

Allee effect that refers to a positive relationship between individual fitness and population density provides an important conceptual framework in conservation biology. While declining Allee effect causes reduction in extinction risk in low-density population, it provides a benefit in limiting establishment success or spread of invading species. Population models that incorporated Allee effect confer the fundamental role which plays for shaping the population dynamics. In population dynamics, the Allee effect (named after the seminal works of the zoologist and ecologist Warder Clyde Allee [1]) refers to a process that reduces the growth rate for small population densities [2].

Allee effect can be either strong or weak [3]. Strong Allee effect can induce a critical density below which per capita growth rate is negative and extinction tends to occur, while weak Allee effect may result in reduced, but still positive, growth rate as population size or density decreases [4].

While a variety of species exhibit Allee effect, its underlying mechanisms still remain unclear. Because of this reason, determining which factors regulate or induce Allee effect continues to be an important subject in the field of both theoretical and experimental research [5-8]. Allee effect can arise from many ecological processes under various interactions of functional and aggregative responses. The previous work revealed that increased risk of predation at low population density [9-12], sexual selection [13,14], reduced mating efficiency [15] and reduced foraging efficiency [16] can give rise to Allee effect.

My objective here is not to reveal which species has Allee effect or which factor can cause Allee effect, but to evoke the importance of Allee dynamics and its potential consequences in population dynamical models. I analyze the stability of three-species food chain model with strong Allee effect.

The rest of the article has been organized as follows. In Section 2, I state the formulation of the model under consideration and its assumptions. Section 3 contains some preliminary results. Then in Section 4 the model is analyzed, identifying its equilibrium points, giving conditions for their feasibility, stability and bifurcation. Numerical simulation has been carried out in Section 5. The article concludes with a discussion of the results obtained.

#### 2. The Mathematical Model

The general food chain model system interaction is represented by the system of following differential equations

$$\frac{DX}{DT} = RXg(X) - F_1(X, Y)Y$$
 ...... (2.1*a*)

$$\frac{DY}{DT} = E_1 F_1(X, Y) Y - D_1 Y - F_2(Y, Z) Z \qquad \dots \dots \dots \dots (2.1b)$$

$$\frac{DZ}{DT} = E_2 F_2(Y, Z) Z - D_2 Z \qquad \dots \dots \dots (2.1c)$$

where X, Y, Z are the population densities of prey, predator and top-predator respectively;  $R, E_1, E_2, D_1, D_2, F_1, F_2$  are positive constants that stands for prey intrinsic growth rate, conversion factors, death rates and functional responses for representing predation process of respective predator.

In this paper, an attempt has been made to update the food chain model (2.1) incorporating Allee effect in the prey species. Thus, I obtain the following model

$$\frac{DX}{DT} = RXg(X)(X - L) - F_1(X, Y)Y \quad \dots \quad (2.2a)$$
$$\frac{DY}{DT} = E_1F_1(X, Y)Y - D_1Y - F_2(Y, Z)Z \quad \dots \quad (2.2b)$$
$$\frac{DZ}{DT} = E_2F_2(Y, Z)Z - D_2Z \qquad \dots \quad (2.2c)$$

where *L* is the survival threshold of the prey and other parameters are defined in the previous model (2.1). Here, we consider the predation functions  $F_1(X,Y) = A_1X$ ,  $F_2(Y,Z) = A_2Y$ , that is, the Holling type-I functional responses where the constants  $A_1$ ,  $A_2$  are two parameters characterising the functional response.

Volume 5 Issue 10, October 2016 <u>www.ijsr.net</u> Licensed Under Creative Commons Attribution CC BY Using the transformation  $x = \frac{X}{K}$ ,  $y = \frac{Y}{KE_1}$ ,  $z = \frac{Z}{KE_1E_2}$ , t =*KRT*, the model (2.2) tales the following form  $\frac{dx}{dx} = x(1 - x)(x - m) = z$ 

$$\frac{dx}{dt} = x(1-x)(x-m) - a_1 x y \quad \dots \quad (2.3a)$$

$$\frac{dy}{dt} = a_1 xy - d_1 y - a_2 yz \quad \dots \dots$$
(2.3b)

$$\frac{dz}{dt} = a_2 yz - d_2 z \quad \dots \tag{2.3c}$$

$$x(0) > 0, y(0) > 0, z(0) > 0,$$
 (2.3d)

where  $m = \frac{L}{K}$ ,  $a_1 = \frac{A_1 E_1}{R}$ ,  $a_2 = \frac{A_2 E_1 E_2}{R}$ ,  $d_1 = \frac{D_1}{KR}$ ,  $d_2 = \frac{D_2}{KR}$ .

# 3. Preliminary Results

#### 3.1. Existence and positive invariance

Theorem: Every solution of system (2.3) with initial conditions (2.3d) exists in the interval  $[0, \infty)$  and  $x(t) \ge 1$  $0, y(t) \ge 0, z(t) \ge 0$  for all t > 0.

**Proof.** Since the right hand side of system (2.3) is completely continuous and locally Lipschitzian on C, the solution of (2.3) with initial conditions (2.3d) exists and is unique on  $[0, \eta)$ , where  $0 < \eta \le \infty$  [3]. From system (2.3) with initial conditions (2.3d), we have

$$x(t) = x(0) \exp\left[\frac{1}{2} \int_0^t \{(1 - x(\theta))(x(\theta) - m) - a_1 y(\theta)\} d\theta\right] \ge 0,$$

$$y(t) = y(0) \exp\left[\int_0^t \{a_1 x(\theta) - d_1 - a_2 z(\theta)\} d\theta\right] \ge 0,$$
  
$$z(t) = z(0) \exp\left[\int_0^t \{a_2 y(\theta) - d_2\} d\theta\right] \ge 0,$$
  
which completes the mass for

which completes the proof.

**3.2.** Equilibrium points and their feasibility

The system (2.3) has the following positive equilibrium points:

i) The trivial equilibrium point  $E_0(0,0,0)$ , the origin, exists always;

ii) The predators free equilibrium points  $E_m(m, 0, 0), E_1(1, 0, 0)$  exist provided m > 0;

top-predator free iii) The equilibrium point  $E_2(x_2, y_2, 0)$  provided  $m \le \frac{a_1}{d_1} \le 1$ ; where  $x_2 = \frac{a_1}{d_1}, y_2 =$  $\frac{(d_1-a_1)(a_1-md_1)}{a_1-md_1}$  $a_1 d_1^2$ 

iv) The system has two distinct co-existence equilibrium points  $E_3(x_3, y_3, z_3), E_4(x_4, y_4, z_4)$ , where

$$x_{3} = \frac{1+m+\sqrt{(1-m)^{2}-4a_{1}y_{3}}}{2}, y_{3} = \frac{a_{2}}{d_{2}}, z_{3} = \frac{a_{1}x_{3}-d_{1}}{a_{2}}, x_{4} = \frac{1+m-\sqrt{(1-m)^{2}-4a_{1}y_{4}}}{2}, y_{4} = \frac{a_{2}}{d_{2}}, z_{4} = \frac{a_{1}x_{4}-d_{1}}{a_{2}}.$$

# 4. Statement of the Main Results

The main properties and qualitative behaviour of the system (2.3) are given in this section. Local stability of equilibrium points are determined by the Jacobian matrix:

$$J(x, y, z) = \begin{bmatrix} (1-x)(x-m) + x(1-x) - x(x-m) - a_1y & -a_1x & 0\\ a_1y & a_1x - d_1 - a_2z & -a_2y\\ 0 & a_2z & a_2y - d_2 \end{bmatrix}$$

**4.1.**  $E_0$  is always stable in nature.

**Proof.** The jacobian matrix  $J_0$  of the system (2.3) at  $E_0$  is given by

$$J_0 = \begin{bmatrix} -m & 0 & 0 \\ 0 & -d_1 & 0 \\ 0 & 0 & -d_2 \end{bmatrix}.$$

The eigen values of  $J_0$  are  $-m, -d_1, -d_2$ . Hence, the equilibrium point  $E_0$  is always stable in nature.

# **4.2.** $E_m$ is unstable in nature.

**Proof.** The jacobian matrix  $J_m$  of the system (2.3) at  $E_m$  is given by

$$J_m = \begin{bmatrix} m(1-m) & 0 & 0 \\ 0 & -d_1 & 0 \\ 0 & 0 & -d_2 \end{bmatrix}.$$

The eigen values of  $J_m$  are  $m(1-m), -d_1, -d_2$ . Since,  $0 \le m < 1$ , the first eigenvalue is always positive. Hence, the equilibrium point  $E_m$  is unstable in nature.

#### **4.3.** $E_1$ is unstable in nature.

**Proof.** The jacobian matrix  $J_1$  of the system (2.3) at  $E_1$  is given by

$$J_1 = \begin{bmatrix} 1 - m & 0 & 0 \\ 0 & -d_1 & 0 \\ 0 & 0 & -d_2 \end{bmatrix}.$$

The eigen values of  $J_1$  are  $1 - m, -d_1, -d_2$ . Since,  $0 \le m < m \le m$ 1, the first eigenvalue is always positive. Hence, the equilibrium point  $E_1$  is unstable in nature.

**4.4.**  $E_2$  is stable if  $\theta = x_2(1 - x_2) - x_2(x_2 - m) < 0$  and  $a_2y_2-d_2<0.$ 

**Proof.** The jacobian matrix  $J_2$  of the system (2.3) at  $E_2$  is given by

$$J_2 = \begin{bmatrix} x_2(1-x_2) - x_2(x_2-m) & -a_1x_2 & 0 \\ a_1y_2 & 0 & -a_2y_2 \\ 0 & 0 & a_2y_2 - d_2 \end{bmatrix}.$$

The eigen values of  $J_2$  are  $\lambda_{1,2} = \frac{(\theta) \sqrt{\{(\theta)^2 - 4a_1^2 x_2 y_2\}}}{2}$  $a_2y_2 - d_2$ , where  $\theta = x_2(1 - x_2) - x_2(x_2 - m)$ . If  $\theta < 0, \lambda_3 = a_2 y_2 - d_2 < 0$ , then, the equilibrium point  $E_2$ will be locally asymptotically stable in nature.

**4.5**. The system experiences Hopf-bifurcation around  $E_2$  for  $a_1 = a_1^{[hb]}$  where  $a_1^{[hb]} = \frac{d_1(1+m)}{2}$ .

**Proof.** From sub-section 4.4, we see that  $\lambda_3$  is real and  $\lambda_{1,2}$ will be purely imaginary if and only if there is a  $a_1 = a_1^{\lfloor hb \rfloor}$ 

# Volume 5 Issue 10, October 2016

# www.ijsr.net

# Licensed Under Creative Commons Attribution CC BY

such that  $a_1^{[hb]} = \frac{d_1(1+m)}{2}$ . Now for = 1,2;  $Re\left[\frac{d\lambda_i}{da_1}\right]_{a_1=a_1^{[hb]}} \neq 0$ . Therefore, the system experiences Hopf-bifurcation around  $E_2$  for  $a_1 = a_1^{[hb]}$ .

**4.6.**  $E_3$  is locally asymptotically stable if  $c_{11} = x_3(1 - x^3 - x^3 x^3 - m < 0)$ .

**Proof.** The jacobian matrix  $J_3$  of the system (2.3) at  $E_3$  is given by

$$J_3 = \begin{bmatrix} c_{11} & c_{12} & c_{13} \\ c_{21} & c_{22} & c_{23} \\ c_{31} & c_{32} & c_{33} \end{bmatrix},$$

where  $c_{11} = x_3(1 - x_3) - x_3(x_3 - m), c_{12} = -a_1x_3, c_{13} = 0, c_{21} = a_1y_3, c_{22} = 0, c_{23} = -a_2y_3, c_{31} = 0, c_{32} = a_2z_3, c_{33} = 0.$ 

The characteristic equation of  $J_3$  is  $\lambda^3 + B_1\lambda^2 + B_2\lambda + B_3 = 0$ , where  $B_1 = -c_{11}$ ,  $B_2 = -c_{23}c_{32} - c_{12}c_{21}$ ,  $B_3 = c_{11}c_{23}c_{32}$ .

Therefore,  $B_1B_2 - B_3 = c_{11}c_{12}c_{21}$ . Now, we choose  $c_{11} < 0$ , then  $B_1 > 0$ ,  $B_3 > 0$  and  $B_1B_2 - B_3 > 0$ . Hence, the proof.

**4.7.**  $E_4$  is locally asymptotically stable if  $x_4(1 - x_4) - x_4(x_4 - m) < 0$ .

The proof is similar with the proof of 4.5.

**4.8.** System (2.3) undergoes a saddle-node bifurcation around interior equilibrium point  $E_3$  with respect to bifurcation parameter m with bifurcation threshold  $a_1^{[sn]}$  if **Proof:** One eigenvalue of the Jacobian matrix  $J_3$  will be zero if det $(J_3) = 0$  which gives  $a_1^{[sn]} = \frac{d_2(1-m)^2}{4a_2}$ . The other two eigenvalues of  $J_3$  are evaluated at  $a_1^{[sn]}$  and one of them must be negative in order to get a saddle-node bifurcation. Let u and v are the eigenvectors corresponding to the eigenvalue 0 of the matrix  $J_3$  and its transpose respectively. We obtain that  $u = (u_1, u_2, u_3)^T$  and  $v = (v_1, v_2, v_3)^T$  where  $u_2 = 0, u_3 = -\frac{c_{21}}{c_{23}}u_1, v_2 = 0, v_3 = -\frac{c_{12}}{c_{32}}v_2$  in which  $u_1$  and  $v_1$  are any two real numbers. Since  $v^T [F_{a_1}(E_3, a_1^{[sn]})] = -x_3y_3v_1 \neq 0$ ,  $v^T [D^2 F(E_3, a_1^{[sn]})(u, u)] \neq 0$ , then the system experiences a saddle-node bifurcation around  $E_3 a_1 = a_1^{[sn]}$  [17].

## 5. Numerical Simulation

Analytical results can never be completed without numerical justification of the derived results. In this section, we present computer simulations of some solutions of the system (2.3). Beside justification of our analytical findings, these numerical simulations are very important from practical point of view.



Figure 1: Locally asymptotically stable behaviour of the system (2.3) around boundary equilibrium point  $E_2$  for the set of parameters  $a_1 = 0.6$ ,  $a_2 = 0.5$ ,  $d_1 = 0.38$ ,  $d_2 = 0.8$ , m = 0.2.



Figure 2: Hopf-bifurcation behaviour of the system (2.3) around boundary equilibrium point  $E_2$  for the set of parameters  $a_1 = 0.6$ ,  $a_2 = 0.5$ ,  $d_1 = 0.35$ ,  $d_2 = 0.8$ , m = 0.2.



Figure 3: Locally asymptotically stable behaviour of the system (2.3) around interior equilibrium point for the set of parameters  $a_1 = 0.7$ ,  $a_2 = 1.5$ ,  $d_1 = 0.18$ ,  $d_2 = 0.2$ , m = 0.2.



Figure 4: Periodic behaviour of the system (2.3) around boundary equilibrium point  $E_3$  for the set of parameters  $a_1 = 1.178765$ ,  $a_2 = 2.95$ ,  $d_1 = 0.27$ ,  $d_2 = 0.4$ , m = 0.2.

# 6. Conclusion and Comments

In this paper, three species food chain model where prey species is affected by strong Allee effect is analyzed and possible dynamical behaviour of this system investigated at equilibrium points. It has been shown that, the solutions posses Saddle-node and Hopf-bifurcations. Both analytically and numerical simulation shown that in certain regions of the parameter space, three species food chain model is sensitively depending on the parameter values.

#### 7. Acknowledgment

This research work is supported by Minor Research Project of University Grants Commission, New Delhi, India vide Ref. No. F. No. PSW-021/14-15 (ERO), ID No. WBI-042 dated 03.02.2015.

#### References

- W.C. Allee. Animal aggregations, a study in general sociology. University of Chicago Press, pages 808-817, 1931.
- [2] P. A. Stephens and W. J. Sutherland. Consequences of the allee e\_ect for behaviour, ecology and conservation. Trends in Ecology, 14: 401-405, 1999.
- [3] Deredec, A., Courchmp, F.: Extinction thresholds in hostparasite dynamics. Ann. Zool. Fenn. 40, 115–130 (2003)

- [4] Groom, M.: Allee effects limit population viability of an annual plant. Am. Nat. **151**, 487–496 (1998)
- [5] Berec, L.,Angulo, E.,Courchamp, F.: Multiple allee effects and population management. Trends Ecol. Evol. 22, 185–191 (2007)
- [6] Chen, J.-X., Peng, L., Zheng, Q., Zhao, Y.-H., Ying, H.-P.: Influences of periodic mechanical deformation on pinned spiral waves. Chaos 24, 033103 (2014)
- [7] Chen, J.-X., Peng, L., Ma, J., Ying, H.-P.: Liberation of a pinned spiral wave by a rotating electric pulse. EPL 107, 38001 (2014)
- [8] Lou, Q., Chen, J.-X., Zhao, Y.-H., Shen, F.-R., Fu, Y., Wang, L.-L., Liu, Y.: Control of turbulence in heterogeneous excitable media. Phys. Rev. E 85, 026213 (2012)
- [9] Andrewartha, H.G., Birch, L.C.: The Distribution and Abundance of Animals. University of Chicago Press, Chicago (1954)
- [10]Birkhead, T.R.: The effects of habitat and density on breeding success in the common guillemot (uria aalge).J. Anim. Ecol. 46, 751–764 (1977)
- [11] Clark, C.W.: Possible effects of schooling on the dynamics of exploited fish populations. J. Cons. Int. Explor. Mer 36, 7–14 (1974)
- [12] Gascoigne, J.C., Lipcius, R.N.: Allee effects driven by predation. J. Appl. Ecol. 41, 801–810 (2004)
- [13]Bessa-Gomes, C., Legendre, S., Clobert, J.: Allee effects, mating systems and the extinction risk in

# Volume 5 Issue 10, October 2016

www.ijsr.net

# Licensed Under Creative Commons Attribution CC BY

populations with two sexes. Ecol. Lett. 7, 802-812 (2004)

- [14] Moller, A.P., Legendre, S.: Allee effect, sexual selection and demographic stochasticity. Oikos 92, 27– 34 (2001)
- [15] Dennis, B.: Allee-effects: population growth, critical density, and the chance of extinction. Nat. Res. Model. 3, 481–538 (1989)
- [16] Bertram, B.C.R.: Living in Groups: Predators and Prey. Blackwell Scientific, Oxford (1978)
- [17] Sotomayor, J.: Generic Bifurcations of Dynamical systems. Dynamical Systems, Academic Press, San Diego, 549–560 (1973).
- [18] Hale JK. Theory of functional differential equations. New York: Springer-Verlag; 1977.