

A Comparison between Modified Mallampati Test, Thyromental Distance, Upper Lip Bite Test and Neck Circumference for Predicting Difficult Intubation

Nishtha Sharma¹, V. C. Chandak², Amol Singam³

¹Anaesthesiology/MD Resident/Jawaharlal Nehru Medical College, Sawangi (M), Wardha, Maharashtra, India

²Anaesthesiology/Associate Professor/Jawaharlal Nehru Medical College, Sawangi (M), Wardha, Maharashtra, India

³Anaesthesiology/Associate Professor/Jawaharlal Nehru Medical College, Sawangi (M), Wardha, Maharashtra, India

Abstract: *Background:* Various bedside screening tests are employed for the prediction of difficult intubation. In this study, we compared 4 bedside tests viz. the Modified Mallampati Test (MMT), the Thyromental Distance (TMD) measurement, the Upper Lip Bite Test (ULBT) and the Neck Circumference (NC) measurement in terms of Sensitivity, Specificity, Positive Predictive Value and Negative Predictive Value to determine which is the best individual predictor of difficult intubation, in apparently normal patients. *Methods:* 120 patients of ASA physical status I&II, aged 16-55years, of either gender, undergoing elective surgery under general anaesthesia were enrolled in this prospective study. MMT Classes III&IV, TMD<6.0cm, ULBT Class III and NC≥35cm were considered to be predictive of difficult intubation. A Cormack-Lehane laryngoscopy grade 3 or 4 was considered a difficult intubation. *Results:* The incidence of difficult intubation was 6.67%. The sensitivity, specificity, positive and negative predictive values of the MMT were 62.50%, 86.61%, 25% and 97% respectively. The sensitivity, specificity, positive and negative predictive values of the TMD were 25%, 93.75%, 22.22% and 94.59% respectively. The sensitivity, specificity, positive and negative predictive values of the ULBT were 37.50%, 91.07%, 23.08% and 95.33% respectively. The sensitivity, specificity, positive and negative predictive values of the NC were 87.50%, 43.75%, 10% and 98% respectively. We also found a statistically significant correlation between the MMT and the CL grading and also between the ULBT and the CL grading. *Conclusion:* No single predictor is sufficient for prediction of difficult intubation on its own. All the studied bedside tests are poor to moderate predictors of difficult intubation. All the tests showed poor positive predictive values and high negative predictive values which suggests that they can be more useful predictors of easy intubation than difficult intubation.

Keywords: Modified Mallampati Test, Thyromental Distance, Upper Lip Bite Test, Neck Circumference, Difficult Intubation

1. Introduction

The most important question that baffles us all, the ultimate question, is how do we define difficult intubation? While some authors take into account the number of attempts¹, number of operators or special intubating aids^{2,3,4}, others⁵ argue that actual failed intubation is a very poor criterion for difficult intubation not only because of its very low incidence but also because of its dependence on the skill of the anaesthesiologist. These anaesthesiologists employ the Cormack-Lehane grading of the glottic view on direct laryngoscopy as the criterion for difficult intubation^{6,7}.

The third definition is a modification of the above CL grading system. Wilson et al⁵ used a scale of 1-5 which was further transformed into what we now know as the Modified Cormack Lehane scale introduced by Yentis and Lee⁸ in 1998 whereas the 4th definition that we came across is that of the intubation difficulty scale (IDS) introduced by Adnet et al⁹ in 1997.

Despite all these definitions, the Cormack-Lehane grading still remains the most widely accepted criterion of difficult intubation.

A study pertaining to difficult intubation can be carried out in 2 ways. A prospective study or a retrospective study. The

later seems like a better option since the incidence of difficult intubation is so low, then why not go for a retrospective study and just measure the features of patients with known difficult intubation? It is however, not that simple. The major drawback here would be that there is no control group- nothing to compare these parameters to. There is nothing to tell us about the easy patients. The percentage of false positives cannot be accurately assessed. The better option then, is to study these features in a lot of routine patients and then identify those which are independent predictors of a difficult intubation.

Several authors have combined various predictors and developed different risk scoring systems^{5,10}. Nonetheless, to develop a scoring system we must first identify which predictors are better than the others so that the best indicators can be combined for a more accurate risk-sum scoring system.

The next concern which arises is the diagnostic accuracy of the various predictors of difficult intubation. The sensitivity, specificity, PPV and NPV of these tests varies from trial to trial. This inconsistency can be attributed to many reasons. Firstly, there are gross differences in the incidence of difficult intubation from one study to the next. Where Oates et al¹¹ found an incidence as low as 1.8%, Tse et al⁷ encountered an incidence as high as 13% in their study.

The second reason for the varied performance of different predictors is ofcourse the definition of difficult intubation used. While one study uses the CL grading as the gold standard of difficult intubation⁶, another uses the IDS¹².

The third reason is the different thresholds used for the different tests especially the quantitative tests. While Brodsky et al¹ consider 46cm as the cut-off point of NC measurement, Gonzalez et al¹² consider 43cm for the same whereas Shailaja et al¹³ take it to be 35cm. The same goes for the TMD measurement. While Frerk² considers a cut-off point of <7.0cm, Khan et al¹⁴ considered <6.5cm whereas Ezri et al¹⁵ and Salimi et al¹⁶ considered <6.0cm and <4cm for prediction of difficult intubation respectively. While it is true that decreasing the threshold of a test will decrease the number of false positives, it is also very true that it will lead to a fall in the predicted number of difficult intubations as well.

The last reason to be considered here is the differences in the demographic profiles in different studies. While some studies only enroll apparently normal ASA physical status I & II patients, others consider the obese, the elderly, the obstetric patients as well as those with known anatomic abnormalities. The question here is whether the accuracy of these tests differs in special groups of patients, especially those in whom difficult intubation is a more dreaded possibility like the obese and the obstetric patients.

The important thing to remember, is that no bedside test is perfect. Therefore, becoming disheartened with the results of a particular test in a particular study and abandoning it all together is not the way to go.

We should find and agree upon standardization of not only the cut-off points of various quantitative tests, but also find one universal definition of difficult intubation to minimize the discrepancies in the outcome of different studies, to some extent.

This study was designed to compare the Modified Mallampati Test (MMT), the Thyromental Distance (TMD) measurement, the Upper Lip Bite Test (ULBT) and the Neck Circumference (NC) measurement in terms of Sensitivity, Specificity, Positive Predictive Value and Negative Predictive Value in order to determine which is the best individual predictor of difficult intubation, in apparently normal patients.

2. Material & Methods

This prospective study was conducted in 120 patients aged 16-55years, of ASA physical status I & II, of either gender, after obtaining institutional ethical committee clearance and written informed consent from all the participants.

Patients with cervical spine abnormalities or trauma, edentulous patients, obese patients (BMI > 30kg/m²), obstetric patients, patients with restricted mouth opening, oral pathology or facial injuries and patients with any thyroid pathology or neck mass were excluded from this study.

A thorough pre-anaesthetic evaluation was carried out in all the patients and the procedure was explained to them in detail. Airway assessment was done in all patients, using the Modified Mallampati Test (MMT), the Thyromental Distance (TMD) measurement, the Upper Lip Bite Test (ULBT) and the Neck Circumference (NC) measurement.

In our study, classification of the oropharyngeal view was done according to the Samssoon and Young's modification of the Mallampati classification-the Modified Mallampati Test (MMT), wherein the seated patient was asked to open the mouth fully with the tongue protruded maximally without phonation. The test was performed twice in each patient to minimize the chances for erroneous observation. The patient was observed from eye level and the view was graded as below.

Class I - Soft palate, fauces, uvula, and pillars are seen

Class II - Soft palate, fauces, and uvula are seen

Class III - Soft palate and base of uvula

Class IV - Soft palate not visible

The TMD was measured in centimeters (cm) from the thyroid notch to the bony point of the chin with the head fully extended on the neck with the mouth closed. A ruler was used and the distance was approximated to the nearest 0.5cm. It was classified as below

- Class I: >6.5cm = normal value
- Class II: 6.0-6.5cm = laryngoscopy and intubation are difficult but usually possible
- Class III : <6.0cm = laryngoscopy may be impossible

The ULBT is classified according to the ability of the lower teeth to bite the upper lip. The observer demonstrated the test to the patient before asking them to perform it. It was assessed and classified as under.

- Class I = lower incisors can bite the upper lip above the vermilion line
- Class II = lower incisors can bite the upper lip below the vermilion line
- Class III = lower incisors cannot bite the upper lip

The NC was measured in centimeters (cm) using a measuring tape at the level of the thyroid cartilage.

MMT Classes III&IV, TMD <6.0cm^{10,15}, ULBT Class III and NC ≥ 35cm¹³ were taken as predictors of difficult intubation. Difficult intubation was defined as Cormack-Lehane grades 3&4 of glottic view upon direct laryngoscopy.

The sensitivity, specificity, positive predictive value and negative predictive value of all these tests was calculated and compared. The correlation, if any, between each test and the CL grading was also assessed.

3. Statistical Analysis

Statistical analysis was done by using descriptive and inferential statistics with the help of Chi-square test and Binary classification. Software used in the analysis was

SPSS 17.0 version and Graph Pad prism 6.0 version. The p-value <0.05 was considered to be significant.

4. Results

The incidence of difficult intubation in our study was 6.67%. There was no failed intubation in our study.

Table 1 shows the distribution of patients according to their demographic profile. In our study, the mean Age of the patients was 34.34 ± 9.15 ; the mean Height was 1.65 ± 0.05 ; the mean Weight was 57.88 ± 5.67 and the mean BMI was 21.02 ± 1.49 .

Table 1: Demographic profile

Parameters	Mean \pm SD	Median
Age (yrs)	34.34 ± 9.15	34.50
Height (m)	1.65 ± 0.05	1.65
Weight (kg)	57.88 ± 5.67	57
BMI (kg/m ²)	21.02 ± 1.49	20.68

BMI = Body Mass Index

Table 2 shows the distribution of patients according to the MMT grades. In our study, 68 (56.67%) patients were in Class I, 32 (26.67%) patients in Class II, 20 (16.66%) patients in Class III and 0 patients in Class IV respectively.

Table 2: Distribution of patients according to the MMT

MMT	No of patients	Percentage (%)
Class I	68	56.67
Class II	32	26.67
Class III	20	16.66
Class IV	0	0.00
TOTAL	120	100.00

Table 3 shows the distribution of patients according to the TMD measurement. There were 84 (70%) patients in Class I (>6.5cm); 27 (22.5%) patients in Class II (6.0-6.5cms) and 9 (7.5%) patients in Class III (<6.0cm) respectively.

Table 3: Distribution of patients according to the TMD

TMD (cm)	No of patients	Percentage (%)
Class I >6.5	84	70.00
Class II 6-6.5	27	22.50
Class III <6.0	9	7.50
TOTAL	120	100

Table 4 shows the distribution of patients according to the ULBT. There were 72 (60%) patients in Class I; 35 (29.17%) patients in Class II and 13 (10.83%) patients in Class III respectively.

Table 4: Distribution of patients according to the ULBT

ULBT	No. of patients	Percentage (%)
Class I	72	60.00
Class II	35	29.17
Class III	13	10.83
TOTAL	120	100

Table 5 shows the distribution of patients according to the NC measurement. The <35cm group had 50 (41.67%) patients and the ≥ 35 cm group had 70 (58.33%) patients.

Table 5: Distribution of patients according to the NC

NC (cm)	No of patients	Percentage (%)
<35	50	41.67
≥ 35	70	58.33
TOTAL	120	100

Table 6 shows the distribution of patients according to Cormack-Lehane Classification of the view on direct laryngoscopy. There were 81 (67.5%) patients with Grade 1; 31 (25.83%) patients had Grade 2; 8 (6.67%) patients had Grade 3 and 0 (0%) patients had Grade 4 respectively.

Table 6: Distribution of patients according to Cormack-Lehane Classification

CL Classification	No of patients	Percentage (%)
Grade 1	81	67.50
Grade 2	31	25.83
Grade 3	8	6.67
Grade 4	0	0
TOTAL	120	100

Table 7 shows the comparison between the sensitivity, specificity, PPV, NPV and Accuracy of the MMT, the TMD, the ULBT and the NC.

Table 7: Comparison between the MMT, the TMD, the ULBT and the NC

	MMT	TMD	ULBT	NC
Sensitivity	62.50%	25%	37.50%	87.50%
Specificity	86.61%	93.75%	91.07%	43.75%
PPV	25%	22.22%	23.08%	10%
NPV	97%	94.59%	95.33%	98%
Accuracy	85%	89.16%	87.50%	46.67%
p-value	8.24, p=0.004,S	3.78, p=0.05,NS	6.31, p=0.012,S	3.00, p=0.08, NS

PPV = Positive Predictive Value; NPV = Negative Predictive Value

The sensitivity, specificity, positive and negative predictive values of the MMT were 62.50%, 86.61%, 25% and 97% respectively whereas the sensitivity, specificity, positive and negative predictive values of the TMD were 25%, 93.75%, 22.22% and 94.59% respectively. The sensitivity, specificity, positive and negative predictive values of the ULBT were 37.50%, 91.07%, 23.08% and 95.33% respectively whereas the sensitivity, specificity, positive and negative predictive values of the NC were 87.50%, 43.75%, 10% and 98% respectively.

We also found a statistically significant correlation between the MMT and the CL grading and also between the ULBT and the CL grading.

5. Discussion

In apparently normal patients with no clinically obvious anatomical abnormalities, unanticipated difficult intubation is not only a challenge to the skill of the anaesthesiologist but also carries the risk of significant morbidity and mortality for the patient.

The various bedside screening tests employed for prediction of difficult intubation should not only have the benefit of

ease of demonstration but should also be time saving as well as reliable. Although one wishes for a test that can predict all difficult intubations without any false positives - a test with extremely high sensitivity and specificity, it is however, unattainable.

The Modified Mallampati Test has been criticized over the years for various limitations. It has been mentioned that the demarcation between classes II & III is unclear^{17, 7}. It has also been pointed out that this test has a high inter-observer variability and a large number of false positives¹¹. Another limitation is that many patients involuntarily tend to phonate which affects the classification^{18, 19}. Despite all its shortcomings, the MMT is still the most routinely used bedside test for the prediction of difficult intubation.

In our study we found the sensitivity of the MMT to be 62.5% which is similar to the results found by Isaac and Sharma²⁰ (62.5%). The specificity of the MMT in our study was 86.6% which correlates with the findings of Shiga et al²¹ (86%) and Gonzalez et al¹² (87%). The positive predictive value of the MMT in our study is 25% which is close to the results of Tse et al⁷ (22%). The negative predictive value of the MMT in our study is 97% which is similar to the studies of Khan et al⁶ (98.4%) and Gonzalez et al¹² (96%). The number of false positive cases in our study were 15.

We also found a statistically significant (p-value = 0.002) correlation between the MMT and the CL grading which was similar to the results of Mallampati et al²² and Ezri et al¹⁵.

The measurement of the TMD originated as a quantitative assessment of the receding jaw. Even though some authors²³ have questioned its relevance as a predictor of difficult intubation, it is a popular bedside test because of its ability to assess not only a receding mandible but also the range of head extension. While the MMT predicts difficult intubation by assessing the amount of soft tissue that would be deformed during laryngoscopy, the TMD measurement does so by assessing the size of the region to which the soft tissue has to be deformed²⁴.

In our study we found the sensitivity of the TMD to be 25% which is slightly more than that found by Shiga et al²¹ (20%). The specificity of the TMD in our study is 93.75% which correlates with the results of Shiga et al²¹ (94%). The positive predictive value of the TMD in our study is 22.22% which correlates with the findings of Salimiet al¹⁶ (22%) and Tse et al⁷ (20%). The negative predictive value of the TMD in our study is 94.59% is similar to that found by Mehta et al²⁵ (93.79%).

There was no correlation between the TMD and the CL grading similar to the results found by Brodsky et al¹.

The ULBT is classified according to the ability of the lower teeth to bite the upper lip. The ULBT not only takes into account the presence of buck teeth but also the range and freedom of movements of the mandible. Moreover, the 3 classes are clearly demarcated, making inter-observer variability highly unlikely.

he ULBT has an important limitation that it cannot be used in edentulous patients and also some patients have difficulty in understanding how to perform the test. Since edentulous patients are excluded from our study, we did not encounter this problem and also the observer in our study demonstrated the test to the patient making it easier to perform.

In our study we found the sensitivity of the ULBT to be 37.50% which was close to the findings of Mohan and Mohana²⁶ (40%). The specificity of the ULBT in our study is 91.07% which correlates with the findings of Khan et al¹⁴ (91.9%), Shah et al²⁷ (91.53%), Eberhart et al²⁸ (92.5%) and Ali et al²⁹ (92.9%). The positive predictive value of the ULBT in our study is 23.08% which is slightly less than the original findings of Khan et al⁶ (28.9%). The negative predictive value of the ULBT in our study is 95.33% which correlates with the results of Shah et al²⁷ (95.70%) and Ali et al²⁹ (97.3%).

There was a statistically significant (p-value = 0.009) correlation between the ULBT and the CL grading similar to the results reported by Hester et al³⁰.

The neck circumference has been regarded a good predictor of difficult intubation in terms of sensitivity¹² and inter-observer reliability³¹.

In our study we found the sensitivity of the NC to be 87.50% which is lower than that reported by Gonzalez et al¹² (92%). The specificity of the NC in our study is 43.75% which correlates with that found by Charaet al³² (44.6%). The positive predictive value of the NC in our study is 10% which is lower than that reported by Charaet al³² (15.4%). The negative predictive value of the NC in our study is 98% which is similar to that of Gonzalez et al¹² (99%).

There was no correlation between the NC and the CL grading similar to the results reported by Karciet al³³.

Combining all the results, the TMD shows the highest accuracy but it also has the poorest sensitivity. The NC has the highest sensitivity but a very poor specificity and accuracy.

All the tests showed poor positive predictive values and high negative predictive values which suggests that they can be more useful predictors of easy intubation than difficult intubation.

The limitations of our study were a strictly limited demographic profile. This study can be conducted in a much wider demographic scenario. We did not assess these tests in combination, which was another limitation of our study. Various combinations of these tests should be assessed and applied for better prediction of difficult intubation.

In conclusion, no single predictor is sufficient for prediction of difficult intubation on its own. All the studied bedside tests are poor to moderate predictors of difficult intubation. All the tests showed poor positive predictive values and high negative predictive values which suggests that they can be more useful predictors of easy intubation than difficult intubation.

References

- [1] Brodsky JB, Lemmens HJ, Brock-Utne JG, Vierra M, Saidman LJ. Morbid obesity and tracheal intubation. *AnesthAnalg* 2002; 94: 732–6
- [2] Frerk CM. Predicting difficult intubation. *Anaesthesia* 1991; 46: 1005-8
- [3] Savva D. Prediction of difficult tracheal intubation. *Br J Anaesth* 1994; 73: 149–53
- [4] Salim F, Rehman HU, Khurshid T, Sharif A. Frequency of difficult and failed intubation in apparently normal patients undergoing elective surgery. *Pakistan Armed Forces Medical Journal*. 2015 Aug 31;65(4):494-7.
- [5] Wilson ME, Spiegelhalter D, Robertson JA, Lesser P. Predicting difficult intubation. *British Journal of Anaesthesia* 1988; 61: 211-6.
- [6] Khan ZH, Kashfi A, Ebrahimkhani E. A comparison of the upper lip bite test (a simple new technique) with modified Mallampati classification in predicting difficulty in endotracheal intubation: a prospective blinded study. *AnesthAnalg* 2003; 96: 595–9
- [7] Tse JC, Rimm EB, Hussain A. Predicting difficult endotracheal intubation in surgical patients scheduled for general anesthesia: a prospective blind study. *AnesthAnalg*. 1995; 81: 254-8
- [8] Yentis SM, Lee DJ. Evaluation of an improved scoring system for the grading of direct laryngoscopy. *Anaesthesia* 1998; 53: 1041–4
- [9] Adnet F, Borron SW, Racine SX, Clemessy JL, Fournier JL, Plaisance P, Lapandry C: The Intubation Difficulty Scale (IDS): Proposal and evaluation of a new score characterizing the complexity of endotracheal intubation. *Anesthesiology* 1997; 87:1290–7
- [10] El-Ganzouri AR, McCarthy RJ, Tuman KJ et al. Preoperative airway assessment: predictive value of a multivariate risk index. *AnesthAnalg*. 1996; 82: 1197-204
- [11] Oates JD, Macleod AD, Oates PD, Pearsall FJ, Howie JC, Murray GD. Comparison of two methods for predicting difficult intubation. *British Journal of Anaesthesia*. 1991 Mar 1;66(3):305-9.
- [12] Gonzalez H, Minville V, Delanoue K, Mazerolles M, Concina D, Fourcade O. The importance of increased neck circumference to intubation difficulties in obese patients. *Anesthesia & Analgesia*. 2008 Apr 1;106(4):1132-6.
- [13] Shailaja S, Nichelle SM, Shetty AK, Hegde BR. Comparing ease of intubation in obese and lean patients using intubation difficulty scale. *Anesthesia, Essays and Researches*. 2014 May;8(2):168.
- [14] Khan ZH, Mohammadi M, Rasouli MR, Farrokhnia F, Khan RH. The diagnostic value of the upper lip bite test combined with sternal distance, thyromental distance, and interincisor distance for prediction of easy laryngoscopy and intubation: a prospective study. *Anesthesia & Analgesia*. 2009 Sep 1;109(3):822-4.
- [15] Ezri T, Wartens RD, Szmuk P, et al. The incidence of class 'zero' airway and the impact of Mallampati score, age, sex, and body mass index on prediction of laryngoscopy grade. *AnesthAnalg* 2001; 93: 1073–5
- [16] Salimi A, Farzanegan B, Rastegarpour A, Kolahi AA. Comparison of the upper lip bite test with measurement of thyromental distance for prediction of difficult intubations. *Acta Anaesthesiologica Taiwanica*. 2008 Jun 30;46(2):61-5.
- [17] Wilson ME, John R. Problems with the Mallampati sign. *Anaesthesia* 1990; 45: 486-487.
- [18] Oates JDL, Oates PD, Pearsall EJ, McLeod AD, Howie JC. Phonation affects Mallampati class. *Anaesthesia* 1990; 45:984.
- [19] Tham EJ, Gildersleve CD, Sanders LD, Mapleson WW, Vaughan RS. Effects of posture, phonation and observer on Mallampati classification. *British journal of anaesthesia*. 1992 Jan 1;68(1):32-8.
- [20] Babu Isaac, Sharma M. Comparison of upper lip bite test and modified mallampati test in prediction of difficult endotracheal intubation. *International Journal of Recent Trends in Science and Technology* May 2015; 15(1):118-121
- [21] Shiga T, Wajima Z, Inoue T, Sakamoto A. Predicting difficult intubation in apparently normal patients: a meta-analysis of bedside screening test performance. *Anesthesiology* 2005; 103: 429–37
- [22] Mallampati SR, Gatt SP, Gugino LD, Desai SP, Waraksa B, Freiberger D, Liu PL. A clinical sign to predict difficult tracheal intubation; a prospective study. *Canadian Anaesthetists' Society Journal*. 1985 Jul 1;32(4):429-34.
- [23] Chou HC, Wu TL. Thyromental distance and anterior larynx: misconception and misnomer. *Anesthesia & Analgesia*. 2003 May 1;96(5):1526-7.
- [24] Baysinger CL, Evans W, Middaugh R, Culclasure J, Culling D. Clinical signs of difficult direct laryngoscopy. *Canadian Journal of Anesthesia/Journal canadien d'anesthésie*. 1987 Jan 1;34(1):103-4.
- [25] Mehta T, Jayaprakash J, Shah V. Diagnostic value of different screening tests in isolation or combination for predicting difficult intubation: A prospective study. *Indian journal of anaesthesia*. 2014 Nov;58(6):754.
- [26] Mohan K, Mohana RL. Comparison of upper lip bite test with thyromental distance for predicting difficulty in endotracheal intubation: A prospective study. *Asian Journal of Biomedical and Pharmaceutical Sciences*. 2013 Jul 1;3(22):62.
- [27] Shah PJ, Dubey KP, Yadav JP. Predictive value of upper lip bite test and ratio of height to thyromental distance compared to other multivariate airway assessment tests for difficult laryngoscopy in apparently normal patients. *Journal of Anaesthesiology Clinical Pharmacology*. 2013 Apr 1;29(2):191.
- [28] Eberhart LH, Arndt C, Cierpka T, Schwaneckamp J, Wulf H, Putzke C. The reliability and validity of the upper lip bite test compared with the Mallampati classification to predict difficult laryngoscopy: an external prospective evaluation. *AnesthAnalg* 2005; 101: 284–9
- [29] Ali MA, Qamar-ul-Hoda M, Samad K. Comparison of upper lip bite test with Mallampati test in the prediction of difficult intubation at a tertiary care hospital of Pakistan. *J Pak Med Assoc*. 2012 Oct 1;62(10):1012-5.
- [30] Hester CE, Dietrich SA, White SW, Secrest JA, Lindgren KR, Smith T. A comparison of preoperative airway assessment techniques: the modified Mallampati and the upper lip bite test. *AANA J* 2007; 75: 177–82
- [31] Adamus M, Jor O, Vavreckova T, Hrabalek L, Zapletalova J, Gabrhelik T, Tomaskova H, Janout V.

Inter-observer reproducibility of 15 tests used for predicting difficult intubation. Biomed Pap Med FacUnivPalacky Olomouc Czech Repub. 2011 Sep 1;155(3):275-82.

- [32] Chara L, Eleftherios V, Maria M, Anastasia T, Chryssoula S. Anatomic features of the neck as predictive markers of difficult direct laryngoscopy in men and women: A prospective study. Indian journal of anaesthesia. 2014 Mar 1;58(2):176.
- [33] Karci A, Karagöz S, Girgin P, Bozdogan DG. Comparison of modified Mallampati classification, upper lip bite test and neck circumference in prediction of difficult intubation: 19AP5-5. European Journal of Anaesthesiology (EJA). 2011 Jun 1;28:236.

