

Sensitivity of Carbendazim against *Curvularia Pallescens* Causing Root Rot Disease to Coriander

A. V. Bhilar¹, N. K. Khandare²

¹Department of Botany, Kasturba Walchand College, Sangli, District- Sangli, 416416 (M.S), India
²Department of Botany, Krantisingh Nana Patil College, Walwa, District- Sangli, 416 313 (M.S), India

Abstract: Coriander (*Coriandrum sativum*) belongs to family Apiaceae, is a green herb with strong smell which gives aroma so used in cooking. Coriander is mainly used as a condiment for its medicinal properties as well as for culinary purpose as spice. Such economically and medicinally important plant is attacked by many fungal pathogens and deteriorates quality and yield. The diseased coriander infected by pathogens was selected for evaluation against fungicide, carbendazim. The MIC of carbendazim among 15 isolate of *Curvularia pallescens* on agar plate (in vitro) and inoculated on plants (in vivo) of coriander showed variation in resistance in both and among the samples.

Keywords: *Coriandrum sativum*, *Curvularia pallescens*, MIC, Carbendazim

1. Introduction

Coriander (*Coriandrum sativum*) belongs to family Apiaceae, is cultivated in Indian states like Maharashtra, Rajasthan, Gujarat, Madhya Pradesh, Tamil Nadu, U.P., etc. It is a green herb with strong smell which gives aroma so used in cooking. It is mainly used as a condiment for its medicinal properties as well as for culinary purposes as spice. It provides high values of dietary fibers and is a rich source of magnesium, iron, non-organic phosphorus, essential oils, oleic acid, palmitic acid, stearic acid, and ascorbic acid etc.

Coriander is infected by pathogens like *Helminthosporium* spp., *Fusarium* spp., *Curvularia pallescens*, *Alternaria* spp., and *Erysiphe polygoni* which cause disease like grain mould disease, wilt, root rot, leaf spot and powdery mildew respectively. They affect quality and quantity of host plant. Root rot of coriander caused by *Curvularia pallescens* is serious among these diseases so there is urgent need to investigate management of these diseases. The aim of present investigation was to access the sensitive and resistant isolates among the select pathogens against carbendazim.

2. Material and Method

2.1. Materials

Infected material of coriander (*Coriandrum sativum*) causing root rot disease infected by *Curvularia pallescens*, collected from districts of Maharashtra and carbendazim (trade name Bavistin) a systemic fungicide.

2.2. Methods

Samples of coriander showing symptoms of root rot disease due to *Curvularia pallescens* were collected from Maharashtra state, during a survey. The samples collected from field in clean bags and pathogens were isolated by direct plating analysis (Pitt and Hocking, 1997). The isolated pathogens were identified (Subramanian 1971 and Barnett and Hunter, 1972), and the pathogenicity test was determined by following Koch's postulates (Koch, 1893). Minimum

inhibitory concentration (MIC) was carried by food poisoning test (Dekker and Gielink, 1979).

In vitro Czapek dox agar plates were prepared containing different concentration (300-3000 µg/ml) of fungicide bavistin (source of carbendazim). Mycelial mat (8mm disc) of the isolates was inoculated at the center of the plate in triplicate. The plates were then incubated at 28±20°C in dark or BOD incubator and radial growth was measured at different intervals. Plates without fungicide were served as control. Calculation was done as percentage method by Vincent.

In vivo (on root rot disease) sensitivity tested on healthy plant of host. For this purpose, healthy plant (replicates of three) was treated with concentration of carbendazim solution. After 24 hours, these carbendazim treated plant roots were inoculated with 50ml, spore suspension (10⁶ spores/ml) of *Curvularia pallescens*, prepared from 8 days old culture in sterile deionized water. After inoculating 25 to 35°C for 10 days and thereafter percentage of infection was recorded by 0-4 scale (Kareem, 2007).

3. Results and Discussion

The root rot of coriander was firstly reported by Dwivedi *et al*, (1982). The pathogen was responsible to causes severe infection to host and loss in quality and quantity. Carbendazim is systematic fungicide used to manage fungal disease. There was variation in MIC of carbendazim among 15 isolate, under *in vitro* and *in vivo* conditions. The MIC of carbendazim among 15 isolates of *Curvularia pallescens* on Agar plate (*in vitro*) and on incubated on plant (*in vivo*) of *Coriandrum sativum* ranged from 300 to 3000 µg/ml *in vitro* and *in vivo* respectively (table.1). The resistance was observed more in the isolate Cp- 8 i.e 3000 µg/ml (*in vitro*) while there was 200 µg/ml (0.2%) MIC was observed in Cp-9 (*in vivo*). There are many workers who determined sensitivity of a pathogen against fungicides. Arora *et al* (1992), noted resistance to metalaxyl in *Phytophthora infestans* in Nilgiri hills of South India. Chander and Thind (1995), observed development of carbendazim resistance in *Gloeosporium ampelophagum*. Latha *et al* (2000), reported root rot disease management

through antagonist and chemicals. Apte and Kamble (2008), studied efficacy of carbendazim in combating castor blight in western Maharashtra. Khandare and Kamble(2013), observed carbendazim sensitivity in *Trigonella foenum graecum*. The above study was very useful for further investigation of resistance. The results are in aggregation with earlier workers like Avenot & Michailides (2007), Blixetal. 2011, Kuang *Jetal.* (2011), Miller & Gubler

(2004), Reis *etal.*, (2005), Stehmann & De Waard (1996), Tremblay *et al* (2003). Viljanen-Rollinsonetal(2007).

4. Conclusion

Carbendazim is very effective in management of *Curvularia pallescens*. Under laboratory (*in vitro*) conditions, resistance of pathogen is more than field (*in vivo*) condition.

Table 1: MIC of carbendazim against *Curvularia pallescens* isolates of Coriander

Sr. no.	Locality	Isolate of root rot of <i>Curvularia pallescens</i>	MIC of <i>Curvularia pallescens</i> (<i>in vitro</i>) µg/ml	MIC of <i>Curvularia pallescens</i> (<i>in vivo</i>) %
1	Shirol (Kolhapur)	Cp-1	2500	2.1
2	Kagal(Kolhapur)	Cp-2	1000	1.1
3	Kodoli (Kolhapur)	Cp-3	900	0.7
4	Junnar (Pune)	Cp-4	800	1.7
5	Indapur(Pune)	Cp-5	2000	1.8
6	Tasgaon (Sangli)	Cp- 6	400	0.3
7	Shirala (Sangli)	Cp-7	2700	2.4
8	Walwa (Sangli)	Cp-8#	3000	2.7
9	Jat (Sangli)	Cp-9*	300	0.2
10	Akluj (Solapur)	Cp-10	700	0.6
11	Madha (Solapur)	Cp-11	2100	1.9
12	Sangola(Solapur)	Cp-12	1100	1.0
13	Karad (Satara)	Cp-13	2400	2.1
14	Faltan(Satara)	Cp-14	1300	1.2
15	Koregaon(Satara)	Cp-15	1800	1.8

*Cp-8- *Curvularia pallescens* isolate is more resistant than others.

#Cp-9- *Curvularia pallescens* isolate is more sensitive than others.

5. Abbreviations

MIC –Minimum inhibitory concentration

Cp- *Curvularia pallescens*

* - Sensitive

- Resistant

References

- Apte, A. R & Kamble S. S, "Efficacy of carbendazim in combating castor blight in western Maharashtra," Bioinfolet, 5(1); 73-74.2008.
- Avenot, H.F & Michailides T.J, "Resistance to boscalid fungicide in *Alternaria alternata* isolates from pistachio in California," Plant Dis, 91:1345–1350.2007
- Barnett, H.L & Hunter, B.B, "Illustrated Genera of Imperfect Fungi". Burgess Publishing Co. Minneapolis, Mn, USA, pp. 241.1972
- Blix, t E., Djurle, A., Yuen J & Olson Å, " Fungicide sensitivity in Swedish isolates of *Phaeosphaeria nodorum*," PlantPathology, 58: 655-664.2009
- Chander, M & Thind, T.S, " Development of carbendazim resistance in *Gloeosporium ampelophagum* and strategies for its management," Indian J. Mycol. Pl. Pathol, 25: 25-33.1995
- Chapman, K. S., Sundin G. W & Beckerman J. L, "Identification of resistance to multiple fungicides in field populations of *Venturiainaequalis*," Plant Dis, 95:921-926.2011
- Dekker J & Gielink A. J, "Acquired resistance to pimaricin in cladosporium and *Fusarium oxysporum* f.sp. narcissi associated with decreased virulence". Neth J. Plant Pathol, 85 : 67.. (1979a and b).
- Dwivedi D. K; Shukla D. N & Bhargava, S. N, "Two new root rot disease of spices," Curr. Sci, 51: 243-244.1982
- Kareem, "Research journal of Agriculture and Biological Science," 3(6);767.2007
- Khandare N. K & Kamble S. S. " The diseases of fenugreek in Maharashtra- A survey", Bioinfolet 10 (1 B) : 335 – 336.2013
- Koch, R. "Über den augenblicklichen stand der bakteriologischen cholera diagnose: zeitschrift hygiene und infections krankheiten," (in german). J. Hyg. Inf, 14: 319 – 333.1893
- Kuang J; Hou Y.P; Wang J.X & Zhou M.G. "Sensitivity of *Sclerotinia sclerotiorum* to fludioxonil: in vitro determination of baseline sensitivity and resistance risk," CropProt, 30:876–882. 2011
- Latha, T.K.S., E. Rajeswari & V. Narashiman. "Management of root rot disease complex through antagonists and chemicals" Indian Phytopath, 53(2): 216-218.2000
- Miller, T.C & Gubler W.D, "Sensitivity of Californian isolates of *Uncinula necator* to Trifloxystrobin and Spiroxamine, and update on Triadimefon sensitivity". Am. Phytopathol. Soc, 88: 1205-1212. 2004
- Pitt, J. I & Hocking A. D. 'Fungi and food spoilage', 2nd Edition. Blackie Academic and Professional: London. pp. 137-142.1997
- Reis, A., Ribeiro F.H.S., Maffia L.A & Mizubuti E.S.G, "Sensitivity of Brazilian isolates of *Phytophthora infestans* to commonly used fungicides in

tomato and potato crops,” *PlantDisease*, 89: 1279–1284. 2005.

- [17] Stehmann ,C& De Waard, M. A. “Sensitivity of populations of *Botrytis cinereato* triazoles, benomyl and vinclozolin”. *Eur. J. Plant Pathol*, 102:171-180.1996
- [18] Subramanian, C. V. *Hyphomycetes*, Indian Council of Agricultural Research: New Delhi. pp .930.1971.
- [19] Tremblay, D.M., Talbot B.G &Carisse, O.“Sensitivity of *Botrytis squamosato* different classes of fungicides”. *Plant Disease*, 87: 573-578.2003.
- [20] Viljanen-Rollinson, S. L. H., Marroni M. V & Butler R. C. Reduced sensitivity to carbendazim in isolates of *Botrytis allii*. *New ZealandPlant Protection*, 60:108-113.2007

Author Profile



Dr. Nilkanth K. Khandare is a Ph.D in Botany (specialization in mycology and plant pathology) from Shivaji University Kolhapur. He is assistant Professor of Botany and has 20 years teaching experience at Krantisinh Nana Patil College, Walwa, District-Sangli, (Maharashtra), India.



Aparna Vijay Bhilar is a post graduated in Botany and Ph.D resisted on Shivaji University Kolhapur, (Maharashtra)

