

Isolation and Antimicrobial Potential of *Streptomycessp.* LCC-06 from the Chennai Coast of Bay of Bengal, India

G. Laxshmi¹, Albin T. Fleming²

^{1,2}PG & Research Department of Advanced Zoology and Biotechnology, Loyola College, Chennai-34, India

Abstract: *Streptomycessp.* is the largest genus of Actinobacteria known for producing antibacterial, antifungal, antiparasitic and immunosuppressant drugs. In the present study, *Streptomycessp.* was isolated from the marine soil of Chennai, along the coasts of Bay of Bengal, India. After processing the soil sediment with calcium carbonate and heat, the suspension was inoculated in Starch casein agar medium by serial dilution method. The culture inoculate isolated from colony was identified as *Streptomycessp.* and named as *Streptomycessp.* LCC-06, where LCC represents "Loyola College Chennai". Production medium-I was found to be the best media for antibiotic production. Secondary metabolite was extracted by mixing the culture broth free of biomass with equal volume (1:1, v/v) of ethyl acetate and methanol. The extract was dried in air and scrutinized for their antimicrobial activity. Gram positive and gram negative bacteria were cultured in Mueller-Hinton agar media and fungal pathogen was grown in potato-dextrose agar medium. They showed remarkable antagonistic potential for *Staphylococcus aureus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Klebsiella pneumonia*, *Escherichia coli*, and *Candida albicans*. Marine soil in Chennai, along the coasts of Bay of Bengal hosts numerous potential antibiotic producing microorganisms like *Streptomycessp.* LCC-06.

Keywords: *Streptomyces* sp., isolation, antimicrobial activity, drug discovery and marine soil

1. Introduction

Actinobacteria are gram positive and filamentous bacteria known for turning out extracellular enzymes and bioactive secondary metabolite with broad spectrum of activities^{1,2}. Of the 23,000 bioactive secondary metabolite reported, 42% of the compounds are from Actinobacteria³. Non-ribosomal polyketide synthase (NRPS) and polyketide synthase (PKS) pathways enable several secondary metabolite productions in Actinobacteria⁴. More than 80% of the known antibiotics isolated from them are industrially decisive⁵. Two of four new antibiotics belong to Actinobacterial strains⁶. Antibiotics isolated also depend on the culture media in which they are grown⁷.

Streptomycessp. is the largest genus of Actinobacteria⁸. *Streptomycessp.* unveils antagonistic effects against human pathogens⁹. They produce two-third of the known-antibiotics and demonstrate discrete metabolic diversity¹⁰. But only about 100 antibiotics have been used commercially to treat human, animal, and plant diseases¹¹. The current study purposes at isolating *Streptomyces* sp. along the Chennai coast of Bay of Bengal, India as they are established throughout the ocean, intertidal zones, ocean sediments and seawater¹². In the present study, they were assessed against gram positive, gram negative and fungal pathogens.

2. Materials and Methods

Isolation of *Streptomycessp.* LCC-06

Marine soil samples were collected along the Chennai coast of Bay of Bengal, India. Pre-treatment of soil was carried out by enrichment techniques to stimulate the growth of slow-growing bacteria. Dried soil was suspended in sterilized sea water and diluted up to 10^{-4} . The suspension was inoculated in starch-casein agar (SCA) medium for 14 days. Pure culture from colony was isolated and inoculated

in SCA for 14 days. Based on culture morphology and microscopic examinations, the culture was found to be *Streptomycessp.* and named as LCC-06, where LCC represents "Loyola College Chennai".

Extraction of Secondary metabolite

LCC-06 was inoculated in production medium-I as this media composition intensify the production of antibiotics. After 14 days, broth media free of biomass was mixed with equal concentration of ethyl acetate and methanol (1:1, v/v) for secondary metabolite extraction. The extract so obtained was air-dried and explored for their antimicrobial activity.

Antimicrobial activity

Antimicrobial activity was executed by well diffusion method. Organisms used for study, namely *Staphylococcus aureus*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Klebsiella pneumonia* and *Escherichia coli* were grown in sterilized Mueller-Hinton agar medium and *Candida albicans* was grown in potato-dextrose agar medium. The Secondary metabolite extract was dissolved in DMSO and evaluated for their aggressive activity against test pathogens with concentration extending from 25 uL to 100uL. Zone of Inhibition was measured by antibiotic zone scale. All the pathogenic strains were acquired from Microbial type cell culture (MTCC), Chandigarh, Punjab, India.

3. Result

The following results were attained by weighing the antagonistic activity of secondary metabolite extracted from *Streptomycessp.* LCC-06.

Table 1: Antimicrobial activity of LCC -06

S. No.	Test organism	Zone of inhibition per concentration (mm)			
	Gram Positive	25uL	50uL	75uL	100uL
1.	<i>Staphylococcus aureus</i>	10	11	14	17
2.	<i>Bacillus subtilis</i>	10	12	13	15
	Gram Negative				
3.	<i>Pseudomonas aeruginosa</i>	-	11	12	13
4.	<i>Klebsiella pneumonia</i>	-	11	12	13
5.	<i>Escherichia coli</i>	10	11	13	15
	Fungi				
6.	<i>Candida albicans</i>	-	10	11	13

4. Discussion

Recent years have beheld the evolution of untold multi-drug resistant bacteria and the necessity to combat has surged relatively. In the present study, *Streptomyces* sp. LCC-06 isolated from the marine environment has revealed noteworthy results against gram positive, gram negative and fungal pathogens. The least concentration being 25uL/well controlled gram positive and gram negative bacteria. The highest concentration of 100uL/well, offered antagonistic activity against all the observed test pathogens. The MRSA, multidrug resistant bacteria of *Staphylococcus aureus* is one of the most confounding pathogen in healthcare management. The above studied secondary metabolite could be the possible solution to these microbes and also to numerous others.

5. Conclusion

Secondary metabolite extracted from *Streptomyces* sp. LCC-06 was isolated from the Chennai coast of Bay of Bengal and revealed an outstanding antagonistic activity against human disease causing pathogenic strains.

References

- [1] Balachandran, C., Duraipandian, V., Arun, Y., Sangeetha, B., Emi, N., Al-Dhabi, N. A., Perumal, P. T. (2016). Isolation and characterization of 2-hydroxy-9,10-antraquinone from *Streptomyces olivochromogenes* (ERINLG-261) with antimicrobial and antiproliferative properties. *Revista Brasileira de Farmacognosia*, 26(3), 285–295.
- [2] UshaNandhini, K. L. (2015). Gas Chromatography-Mass Spectrometry Analysis of Bioactive Constituents from the Marine *Streptomyces*. *Asian Journal of Pharmaceutical and Clinical Research*, 8(2), 244–246.
- [3] Karthik, L., Kumar, G., & Rao, K. V. B. (2013). Antioxidant activity of newly discovered lineage of marine actinobacteria. *Asian Pacific Journal of Tropical Medicine*, 6(4), 325–332.
- [4] Manivasagan, P., Venkatesan, J., Sivakumar, K., & Kim, S. K. (2013). Marine actinobacterial metabolites: Current status and future perspectives. *Microbiological Research*, 168(6), 311–332.
- [5] Zarina, A., & Nanda, A. (2014). Antimicrobial, Antioxidant and Cytotoxic Activity of Marine *Streptomyces* MS-60 Isolated from Bay of Bengal. *International Journal of Science and Research*, 3(12), 1634–1640.
- [6] Claveras, F. P., Undabarrena, A., Gonzlez, M., Seeger, M., & Cmara, B. (2015). Culturable diversity and antimicrobial activity of Actinobacteria from marine sediments in Valparaso bay, Chile. *Frontiers in Microbiology*, 6(7), 1–11.
- [7] Duraipandian, V., Al-Dhabi, N. A., & Ignacimuthu, S. (2016). New antimicrobial anthraquinone 6,61-bis (1,5,7-trihydroxy-3-hydroxymethylanthraquinone) isolated from *Streptomyces* sp. isolate ERI-26. *Saudi Journal of Biological Sciences*, 23(6), 731–735.
- [8] Rashad, F. M., Fathy, H. M., El-Zayat, A. S., & Elghonaimy, A. M. (2015). Isolation and characterization of multifunctional i species with antimicrobial, nematicidal and phytohormone activities from marine environments in Egypt. *Microbiological Research*, 175(1), 34–47.
- [9] Balachandran, C., Duraipandian, V., Emi, N., & Ignacimuthu, S. (2015). Antimicrobial and cytotoxic properties of *Streptomyces* sp. (ERINLG-51) isolated from Southern Western Ghats. *South Indian J. Biol.*, 1(1), 7–14.
- [10] Hou, Y. H., Li, F. C., Wang, S. J., Qin, S., & Wang, Q. F. (2008). Intergeneric conjugation in holomycin-producing marine *Streptomyces* sp. strain M095. *Microbiological Research*, 163(1), 96–104.
- [11] Kumar, P. S., Duraipandian, V., & Ignacimuthu, S. (2014). Isolation, screening and partial purification of antimicrobial antibiotics from soil *Streptomyces* sp. SCA 7. *Kaohsiung Journal of Medical Sciences*, 30(9), 435–446.
- [12] Subramani, R., & Aalbersberg, W. (2012). Marine actinomycetes: An ongoing source of novel bioactive metabolites. *Microbiological Research*, 167(10), 571–580.