

Plankton Diversity and Density in Pariyat River, Rural Area of Jabalpur (M.P.)

Neelam Tiwari

Research Scholar-Govt. Model Science College Jabalpur 482001 (M.P.), India

Abstract: The present study was carried out on Pariyat River of Pariyat Village. Pariyat village is situated in Kundam Tehsil of Jabalpur District, M.P. state. Pariyat village is 20 km. far from its district place Jabalpur. Current population of Pariyat village is about 15000. Pariyat lake is situated on western side of the village. The plankton were collected, counted and were identified by using the method suggested by APHA (1995) Present (1970) and Fresh Water Biology (W.T. Edmondson-1959). The plankton were counted by using Sedgwick Rafter counting cell. Different class such as cyanophyceae, chlorophyceae, bacillariophyceae, euglenophyceae, ciliata, cladocera, copepod and specimens from phylum rotifera were identified during the study. Among all these classes the listed phytoplankton such as *Oscillatoria* Sp., *Tetraedron* Sp., *Navicula* Sp., *Nitzschia* Sp., *Euglena* Sp., and zooplankton such as *Ceriodaphnia* Sp., *Cyclops* Sp., *Tropocyclops* Sp., *Brachionus* Sp. and *Platias* Sp. were recorded as a dominant genera in Pariyat Lake. The study was carried out monthly but was tabulated seasonally by using statistical method.

Keywords: Pariyat River, Phytoplankton and Zooplankton

1. Introduction

Jabalpur is the largest city of the Madhya Pradesh state. Today the city becomes the Smart City of the M.P. It has its own indefinable charm, combining many traditional elements and the latest international blend. The Pariyat river is the main river of Jabalpur. The Pariyat Dam is the Second important dam of Jabalpur. It is located on the western part of Pariyat village. The sewage waste of Pariyat village is directly discharge into this River. The people of village also use this river to wash their cloths, take bath, sanitation, discharge of cow dung by dairies etc. The cattle of the villagers also take bath in this River. The length of the river is above 55 Km. long, depth is 21 feet. Pariyat river is geographically located at latitude 23 ° 15'0" and longitude 23 ° 15'0".

2. Materials and Methods

The standard method suggested in APHA used for assessing water quality includes collection, counting and identification of phytoplankton and zooplankton. Plankton net number 25 of mesh size 20µm was used for collecting samples. 50 liters of water was measured in a graduated bucket and filtered through the net and concentrated in a 100 ml. bottle. Samples were collected as close to the water surface as possible in the morning hours. The samples were labeled with the date, time, and study area i.e. name of lake and the volume measured and pasted on the containers. Plankton is preserved by using 4% formalin. The sample was allowed to settle for 24-48 hours and was further concentrated to approximately 30 ml. by decanting. Sedgwick Rafter counting cell is used to count the plankton. Sedgwick Rafter cell is approximately 50 mm. long, 20 mm. wide and 1 mm. deep. The total volume of the cell is 1 ml. A binocular compound microscope is used to count the plankton with different eyepieces such as 10X and 40X. The microscope is calibrated using an ocular micrometer. Formula to convert unit/ml of plankton into unit/liter is

$$n = (a \times 1000) c / l$$

Where,

n = Number of plankton / liter of water.

a = Average no. of plankton in one small counting chamber of S-R cell.

l = Volume of original water filtered in liter.

3. Results and Discussion

Plankton has long been used as indicator of water quality. Because of their short spans, plankton responds quickly to environmental changes. They flourish both in highly eutrophic waters while a few others very sensitive to organic and/or chemical wastes. Some species have also been associated with noxious blooms sometimes creating of offensive tastes and odours or toxic conditions. Because of their short life cycles plankton respond quickly to environmental changes, and hence the standing crop and species composition indicate the quality of the water mass in which they are found.

In the present study 4 different genera of *cyanophyceae* class were recorded from the River. The blue green algae recorded in Pariyat River are *Merismopedia* sp., *Nostoc* sp., *Oscillatoria* sp., and *Spirulina* sp. the minimum algal units were recorded during summer season whereas maximum was recorded during winter season.

In Pariyat River 7 different genera of *chlorophyceae* class were recorded. In Pariyat River the algae recorded are *Closterium* sp., *Closteriopsis* sp., *Coelastrum* sp., *Mugeotia* sp., *Spirogyra* sp., *Tetraedron* sp., *Scenedesmus* sp., the minimum value was recorded during summer season whereas maximum was recorded during monsoon season. From the Pariyat River 7 different genera of *bacillariophyceae* were recorded. In Pariyat River the diatom for *bacillariophyceae* class recorded are *Cyclotella* sp., *Cymbella* sp., *Gomphonema* sp., *Gyrosigma* sp., *Navicula* sp., and *Nitzschia* sp., The minimum unit of diatom were recorded during winter season whereas maximum unit of diatoms were recorded during summer season.

In the Pariyat 2 genera of euglenoids were recorded. The euglenoid recorded in the River are *Euglena sp.* and *Phacus sp.* In Pariyat River the minimum units of euglenoids were recorded during summer season and maximum unit of euglenoids were recorded during summer season.

In Pariyat River 2 genera of class ciliata were recorded. The ciliate recorded in the River are *Pleuronema sp.* and *Strombidium sp.* and 2 genera of cladocera are *Alona sp.* and *Ceriodaphnia sp.* In Pariyat River 5 genera of copepod, *Cyclops sp.*, *Dicacyclops sp.*, *Diaptomus sp.*, *Eucyclops sp.*, and *Tropocyclops sp.*, were recorded. The minimum number of copepods were recorded during summer season whereas maximum unit of copepods were recorded during winter season.

In the Pariyat River 2 genera of phylum rotifer *Brachionus sp.* and *Platyas sp.* were recorded. The minimum number of rotifera were recorded during summer season whereas maximum unit of rotifer were recorded during monsoon season. Density of Phytoplankton and Zooplankton were mention in fig. 1 and 2.

4. Conclusion

Therefore from the above study it is concluded that the total phytoplankton count/mi. is more in summer season and *bacillariophyceae* is dominant while total zooplankton count/ml. is more in winter season. Rotifer is dominant in Pariyat River. The total plankton count/ml. is minimum in monsoon season 256/ml. and maximum in summer season 360ml/l (Table-1). This was proved by the above result as the amounts of *cyanophyceae* algae and rotifer were found in Pariyat River, which are indicator of pollution.

Table 1: Seasonal Variation of Phytoplankton in Priyat River

Phytoplankton		Sampling season		
Class	Genera	Monsoon Mean	Winter Mean	Summer Mean
Chlorophyceae				
	Closteriopsis Sp.	10	0	0
	Coelastrum sp.	8	0	0
	Mugotia sp.	0	14	8
	Pondorina sp.	4	0	0
	Scenedesumus sp.	0	0	2
	Spirogyra sp.	0	12	0
	Tetraedron sp.	6	0	8
Total		28	26	18
Cyanophyceae				
	Merismopedia sp.	4	0	0
	Nostoc sp.	0	4	0

PHYTOPLANKTON COUNT/SEASON




Fig. No.1 Season wise phytoplankton count

	Oscillatoria sp.	14	24	0
	Spirulina sp.	6	0	0
Total		24	28	0
Bacillariophyceae				
	Amphiplura sp.	2	0	0
	Cyclotella sp.	2	0	2
	Cymbella sp.	4	0	4
	Gomphonema sp.	0	8	4
	Gyrosigma sp.	0	0	24
	Navicula sp.	58	16	96
	Nitzschia sp.	22	20	78
Total		88	44	208
Euglenophyceae				
	Euglena acus	4	2	2
	Euglena gracillis	0	10	2
	Euglena sp.	12	42	66
	Phacus sp.	2	2	0
Total		18	56	70
Total phytoplankton count/ml		158	154	296
Total phytoplankton count/l		94800	92400	177600

Table 2: Seasonal Variation of Zooplankton in Priyat River

Zooplankton		Sampling season		
Group	Genera	Monsoon Mean	Winter Mean	Summer Mean
Ciliata				
	Pleuronema sp.	2	6	0
	Strombidium sp.	2	0	0
Total		4	6	0
Clasocera				
	Alona sp.	0	0	6
	Ceriodaphnia sp.	16	28	32
Total		16	28	38
Copepoda				
	Cyclops sp.	10	10	2
	Dicacyclops sp.	0	10	0
	Diaptomus sp.	2	12	0
	Eucyclops sp.	4	8	0
	Tropocyclops sp.	4	12	6
Total		20	52	8
Rotifera				
	Branchionus sp.	56	4	14
	Platyas sp.	2	18	4
Total		58	22	18
Total Zooplankton count/ML		98	108	64
Total Zooplankton count/L		58800	648000	38400
Total Plankton Count/ML		256	262	360
Total Plankton count/L		153600	157200	216000

ZOOPLANKTON COUNT/SEASON




Fig. No.2 Season wise zooplankton count

5. Acknowledgment

We are thankful to the Head Department of Zoology Govt. Model Science College, Jabalpur for providing the necessary laboratory facilities and Dr. K.K. Dube, (Retd.) Dr. Rita Bhandari Pro.OFK College Khamaria, Jabalpur for their help in work of laboratory.

References

- [1] APHA 1995. Standard methods for the examination of water and waste water. 19th Edition, American Public Health Association, American Water Work Association and Water Pollu. Contl. Federation, Washington, D.C., New York.
- [2] A.K. Koorosh, Sadanand and Altaff 2008. Abundance of Zooplankton in Three Contrasting Lakes of Mysore City Karnataka State.
- [3] Battish S.K. Fresh water zooplankton of India. Oxford and IBH Pub. Co. New Delhi 1992.
- [4] Chakrabarty R.D., Roy P. and Singh S.B. 1959. A quantitative study of Plankton and physicochemical condition of river Jamuna at Allahabad in 1945 to 1959. Indian Journal of Fisheries. 6(10): 186-203. De Lange E. 1994, Manual for Simple Water Quality Analysis International Water Tribunal (IWT) Foundation: Amsterdam.
- [5] Descy J.P., Servais P., Smits J.S., Billen G. and Everbecq E. 1987. Phytoplankton Biomass and production in the river Meuse (Belgium) Wat. Res. 21, 12pp: 1557-1566.
- [6] Dhakar M.L. 1979. Studies in some aspects of the hydrobiology of Indrasagar tank (South Rajasthan) Ph.D. thesis, Uni. Of Udaipur.
- [7] Edmondson, W.T. 1959. Fresh Water Biology. 2nd ed John Wiley and sons. New York 1248pp.
- [8] Gawas A.D., Lokhande P.B. and Meijawas H.A. (2006): Study of Physico-Chemical Parameters of surface water in the Mahad Industrial Area. Poll Res. 25 (1): 109-114.
- [9] Henry R., Tundisi J.G. and Curi P.R. (1984). Effect of phosphorus and nitrogen enrichment on the phytoplankton in tropical reservoir, Hydrobiologia, 118(2): 177-186.
- [10] Jones F.H. and Barrington R.I. 1985. A study of the suspended algae in the river Derwent. Derbyshire. U.K. Hydrobiologia, 128: 255-264.
- [11] Kumar A. (1996). Comparative study on diel variation of abiotic factor in lentic and lotic fresh water ecosystem of Santal Paragana (Bihar). J. Environ. Pollut. 3: 83-89.
- [12] Lannineer J., Lea K. and Yrgana E.R. 1982. The role of nitrogen as a growth limiting factor in eutrophic Lake Vesijarvi. Southern Finland, Hydrobiology, 87: 81-85.
- [13] Munawar M. 1972. Ecological studies of euglenae in certain polluted and unpolluted environments. Hydrobiologia 39: 307-320.
- [14] Munawar M. 1970. A Limnological studies of fresh water ponds of Hyderabad, India -1. Journal of the Biotype Hydrobiologia, 35, pp. 127-162.
- [15] Oemke D.V. and Burtan T.M. 1986. Diatom colonization dynamics in a lotic system. Hydrobiologia. 139: 153-166.
- [16] Orihideja Tasevaka, Dafina Guseska 2010 Rotifer based Assessment Of The Lake Dojarn Water Quality. Hydrobiological Institute Ohrid, Macedonia.
- [17] Paransivam M. and Sreenivasan A. 1981. Changes in algal flora due to pollution in Cauvery river. Indian J. Environ. Health. 23(3): 222-238.
- [18] Plankin J. L., M.T. Barbour, K.D. Porter, S.K. Gross, R.M. Hughes. 1989. Rapid Assessment Protocols for.
- [19] Use in Streams and Rivers: Benthic Macroinvertebrates and Fish. EPA:
- [20] Prescott G.W. 1970. Algae of the western great Lakes areas. Pub Cranbrook Institute of science Bulletin. 33: 1-496.
- [21] Rafter, G.W. 1900. The microscopical examination of potable waters. Van. Norstrand Co.
- [22] Saad M.A. and Abbas M.H. 1985. Limnological investigations on the rosetta branch of the Nile II Phytoplankton, Fresh water Biol. 15: 661-669.
- [23] Sarles W.B. 1961. Madisons Lake must Urbanisation destroy their beauty and productivity. Algae and Metropolitan wastes- Tr. Of the 1960 Seminar U.S. Dept. of Health Edu. And Welf: pp: 10-18.
- [24] Sengar R.M.S. and Sharma K.D. 1982. Algal flora of Yamuna river at Agra: Chlorococcales Phycos 21: 164-165.
- [25] Sharma N. 1983. Investigations on limnology of tank ecosystem near Allahabad with particular reference to the abundance and seasonal distribution of plankton and benthos, Ph.D thesis, Sagar University, Sagar.
- [26] Singh, S.R. and Swarup, K. 1979. Limnological studies of Saraha Lake (Billia). II. The periodicity of phytoplankton. J. Ind. Bot. Soc. 58: 319-329.
- [27] Subrahmanyam N.S. and A.V.S. Sambamurty. (2004): "Ecology" Narosa Publication.
- [28] Trivedi, R.K. 1993. Biomonitoring of water pollution in R.K. Trivedi edited, Encyclopedia of Environmental pollution and Control Vol 1 and 2. Environ Media, Karad, India.
- [29] Vyas L.N. 1968. Studies on phytoplankton ecology of Pichola Lake. Udaipur proc. Symp. Recent Adv. Trop. Ecol: 334-347.