

Stopping Power Calculation of Protons using Bloch Correction and Semi Empirical Relation

Dr. Rashid Owaid Kadhim¹, Ahlam Habeeb Hussien²

University of Kufa /College of Education for Girls

Abstract: One of the needs of various accelerators to stimulate recent developments in calculating the energy loss of charged particles in matter is heavy ion ionization energy loss. Bloch incorporates Bethe and Bohr theory in a very transparent way. Thus, Bloch found the bridging formulation between the classical Bohr impact –parameter approach and the quantized Bethe momentum transfer approach to energy loss. In this paper, the behavior of Bloch correction of protons in (C₃H₆, C₂₂H₁₀N₂O₅) and for the energy range (0.1- 1000)MeV has been studied. We have described an implementation of Bloch correction to stopping power (dE / dx) in order to take into account the deviations from the Bethe theory at non – relativistic and relativistic velocities as well as the effects at ultra – relativistic velocities. Bloch correction is expressed in terms of the transport cross section for electron – ion scattering and dependence on the scattering phase shifts. Were calculated stopping power equation by quasi-experimental results showed using both methods showed good agreement with the results SRIM 2012.

Keywords: effective of Bloch, stopping power

1. Introduction

The electronic energy loss has been studied for many years because of its direct application in problems concerning material damage, ion beam analysis and plasma physics. The theoretical treatment of the energy loss in atomic collisions has been greatly improved over the last decades[1]. Stopping power, i.e. energy loss of energetic particles per unit length in matter, has been studied experimentally and theoretically since the beginning of the 20th century because of its wide application area, such as ion implantation, fundamental particle physics, nuclear physics, radiation damage, radiology, and structure analysis of solid target by Rutherford backscattering spectroscopy [2]. The slowing down of energetic ions in matter is dominated by momentum exchanging collision with electrons. The theory of this venerable subject was formulated early on by Bohr [3]and Bethe[4]. The first classical calculation of the energy loss of energetic particles was made by Bohr[3], while the first quantum mechanical treatment was done by Bethe[4]. This latter theory of stopping power is particularly accurate when the projectile's velocity is sufficiently high. It was modified by Bloch [5], and it was shown that at relativistic velocities a Mott correction for spin changing collisions was required[6]. Lindhard and Sorensen performed exact quantum mechanical calculations on the basis of the Dirac equation to produce values for the average energy loss and straggling which are stated to be accurate for any value of projectile charge[7].

2. Stopping Power

The mean ionization energy loss of charged particles heavier than electrons is given by the Bethe expression[8]

$$-\frac{1}{\rho} \frac{dE}{dx} = 4\pi N_a r_e^2 mc^2 \frac{z_1^2 z_2}{A_1 A_2} \frac{1}{\beta^2} L(\beta) \quad (1)$$

Where

$L(\beta)$ is the ionization logarithm and presented in the following form

$$L(\beta) = L_0(\beta) + \sum_i \Delta L_i \quad (2)$$

$$L_0 = \ln \left(\frac{2m_e c^2 \beta^2 \gamma^2}{I} \right) - \beta^2 - \frac{\delta}{2} \quad (3)$$

$\gamma = (1 - \beta^2)^{-1/2}$ is the Lorentz factor

$r_e = \frac{e^2}{4\pi\epsilon_0 mc^2}$ is the classical electron radius, $\beta = \frac{v}{c}$ is the velocity of the particle relative to light velocity.

Z_1 and A_1 are the atomic and mass of the projectile, Z_2 and A_2 are the atomic and mass of the target material, I and δ are the mean excitation energy and density correction, respectively, N_a is Avogadro's number.

When neglecting all the corrections ΔL and dealing only with the $L_0(\beta)$, eq.(1) is referred to as the Bethe equation. The corrections ΔL are to take into account the deviations from the Bethe theory for ions at both low and high energies[8]. This is the form derived originally from the quantum perturbation theory, and the first two terms are typically called the Bethe result. The third term is the density effect. We will refer to as the Bethe result inclusive of the density effect[9]. At high energies, the electronic stopping is determined by Bethe's or Bohr's treatment [10]. The decisive point is whether or not the problem can be treated by classical theory. This is possible for large values of Bohr's parameter [11].

$$K = \frac{2z_1 v_0}{v} = 2\eta$$

In the quantum mechanical limit $K \ll 1$ Bethe's formula based on a first order perturbation treatment gives the high – energy expression in the non relativistic case valid for light ions[10]

$$L_{Bethe} = \ln \frac{2mv^2}{I} \quad (4)$$

Once notes that the condition for quantum can not be expected to hold for slow projectiles or projectiles of high atomic number. In this case the logarithm in eq.(4) is replaced by Bohr's expression[10]

$$L_{Bohr} = \ln \frac{cmv^3}{z_1 e^2} \quad (5)$$

$C = 2e^{-\gamma^-} = 1.1229$; $\gamma^- = 0.5772$ is (Euler's constant).

The major difference is that enters into the argument of the logarithm, and therefore, influences the position of the stopping maximum. is a characteristic frequency of each electron level of the target atom, since the derivation is performed for a harmonic oscillator[10].

3. Bloch Correction (ΔL)

Bloch indicated the exact scattering amplitude (but non – relativistic) should be adopted for collisions with small impact parameter, and derived a correction which is called (Z_1^4) correction term[12]

$$\Delta L_{Bloch} = Z_1 L_2 = \Psi(1) - Re\Psi(1 + i \frac{z_1 v_0}{v}) \quad (6)$$

Here $\Psi(x) = d \ln \Gamma(x) / dx$ is the digamma function, v_0 is the Bohr velocity and with $y = \frac{z_1 v_0}{v}$, eq.(6) becomes

$$\Delta L_{Bloch} = Z_1 L_2 = \Psi(1) - Re\Psi(1 + iy) \quad (7)$$

To bridge the gap between the Bethe's quantum mechanical and Bohr's classical formula. This term is mainly originated in the close collision with small scattering angle which is important for low velocity ions[12]. Eq.(6) may be as a chain

$$\Delta L_{Bloch} = Z_1 L_2 = -y^2 \sum_{n=1}^{\infty} [n(n^2 + y^2)]^{-1} \quad \text{for } y > 1 \quad (8)$$

$$\Delta L_{Bloch} = Z_1 L_2 = \sum_{n=1}^{\infty} (-1)^n \xi(2n + 1) n y^2 \quad \text{for } y < 1 \quad (9)$$

Where $n=1, 2, 3, \dots$ and ξ is the Riemann function. From a practical viewpoint of calculating accurate stopping powers, Bichsel has proposed simple parameterization of the Bloch correction which accurately fits a wide range of high velocity stopping data[13]

$$\Delta L_{Bloch} = Z_1 L_2 = -y^2 [1.202 - y^2 (1.042 - 0.855y^2 + 0.343y^4)] \quad (10)$$

For low velocity, the value of (ΔL_{Bloch}) becomes

$$\Delta L_{Bloch} \rightarrow 0.58 - \ln(y) \quad (11)$$

And thus the Bloch correction provides the transition to the classical stopping formula of Bohr.

For high velocities, i. e. $y \rightarrow 0$, the value of (ΔL_{Bloch}) becomes

$$\Delta L_{Bloch} = -1.202y^2 \quad (12)$$

This term is usually quite small. According to Bloch, the stopping cross section of an atom is determined by the following expression for the stopping number[5]

$$L_{Bloch} = L_{Bethe} + \Delta L_{Bloch}$$

$$L_{Bloch} = \ln \frac{2mv^2}{I} + \Psi(1) - Re\Psi(1 + i \frac{z_1 v_0}{v})$$

therefore, the stopping power that includes the Bloch correction is written as:

$$-\frac{1}{\rho} \frac{dE}{dx} = 4\pi N_a r_e^2 m c^2 \frac{z_1^2 z_2}{A_1 A_2} \frac{1}{\beta^2} L_{Bloch} \quad (13)$$

When Bloch correction (ΔL_{Bloch}) is added to (L_{Bethe}), one arrives at the Bohr stopping formula at the low – velocity end. ΔL_{Bloch} reduces to the Bohr logarithm eq.(5) at low projectile velocity and at high projectile velocity, ΔL_{Bloch} goes to zero[3].

4. SRIM 2012 Program

It is a number of experiments that are related to the measurement of the stopping power and ionic ranges in the elements. It was designed by Ziegler and his group[14]. And because of the high range of power usage which is used in the low, medium, and high power areas, it has spread very fast. Lots of improvements have been added since its first presentation in 1985, and the vehicles stopping power have been improved as well. 2800 new stopping power have been added to the data base of the SRIM program which made it more than 28000 stopping power in the SRIM data base. The new stopping power of the heavy ions emphasized the important enhancements for the precision of the stopping power in the SRIM 2012 program.

5. Semi Empirical Relation for Stopping Power Calculation

Many of the empirical and theoretical studies were performed in order to form standard energy and range relation to account for stopping power. The subject had been reviewed by some researchers like Taylor, Bethe, Askin, Allison, Warsaw, Uehling, Barkas and Berger in the last decades (50-60). And most of the empirical data had been collected by Whaling Bichsel in the form of tables

A. K. Chaubey and H. V. Gupta arrived at the following empirical relation for the stopping power of protons[15]:

$$-\frac{dE}{\rho dx} = \frac{a}{A_2} E^{-b} Z_2^c \log E + d \quad (14)$$

The appropriate values of the constants a, b, c, d are $a=915$, $b=0.85$, $c=0.145$, $d=0.635$, while ρ is the density, A_2 is the atomic weight and Z_2 is the atomic number for the target and E is the kinetic energy of the particle by the units MeV/amu.

The eq(14) is found to be valid in the energy range(0.7-12)MeV/amu, the stopping power is in MeV cm²/gm. The constants c, d are found to be independent of particle type and were obtained by fitting Northcliffe and Schilling[16] (which was called NS later), it accounted the stopping power by the least squares method while a, b were extracted from the empirical data by Whaling[17] and Anderson et al.[18] as well as NS data in the low energies.

The stopping power for the ions heavier than protons can be found by the expressions given by Blann and pierce[19]:

$$\left(-\frac{dE}{\rho dx}\right)_H = \frac{Z_{eff}^2}{\delta_p^2} \left(-\frac{dE}{\rho dx}\right)_p \quad (15)$$

Where $Z_{eff}^2 = \delta^2 Z^2$, the energy is above 0.7 MeV here, so for protons we have taken $\delta_p = 1$ by compensating(14) in (15) we get:

$$\left(-\frac{dE}{\rho dx}\right)_H = \gamma^2 Z^2 a E^{-b} Z_2^c \log E + D \quad (16)$$

In eq.(16) the fractional effective charge γ of the ion of energy E(MeV/amu) can be guessed from the empirical formula of Grant and Booth[20]: $\delta^2 = f(E Z^{-4/3})$

Where

$$f(x) = 1 - \exp(-24.73x + 247.6x^2 - 1131x^3)$$

For the calculation of the stopping power of alpha particles from the equation (16), we assume that the energy is bigger than 4MeV and the fractional effective charge of the alpha particles $\delta^2 = 1$. and when the energy is less than 4MeV or 0.7 to 1 MeV/amu, by fitting NS data with alpha particles by the least squares method and keeping c and d as constants we get a= 3574, b=0.84 so the energy between 0.7 to 1 MeV/amu for Alpha particles and the equation (14) should be modified into:

$$\left(-\frac{dE}{\rho dx}\right) = \frac{3574}{A_2} E^{-0.84} Z_2^{0.145} \log E + 0.635 \quad (17)$$

We can get the stopping power in the compound targets by the summation theory:

$$\left(-\frac{dE}{\rho dx}\right)_{\text{compound}} = \frac{1}{M_2} \sum_i N_i A_i \left(-\frac{dE}{\rho dx}\right)_i \quad (18)$$

Where M_2 is the molecular weight of the compound medium containing N_i atoms of atomic weight A_i .

In addition to, Anderson and Ziegler[21] summarized the empirical data of proton stopping power to many elements of wide range energy. And to get the values for all the elements in a continuous range of protons energies, the researcher had fitted the curves with the available empirical data to generate Coefficient e to be used in the semi empirical stopping power as a function to the proton energy E(keV) and the atomic number Z_2 to the target. It was assumed that S_e to be fit with $E^{0.45}$ to the energy $E < 25$ keV, except $Z_2 \leq 6$ which is fit with $E^{0.25}$ Zielger et al. used the following relation for ($25 \text{keV} \leq E \leq 10 \text{MeV}$):

$$\frac{1}{S_e} = \frac{1}{S_{\text{LOW}}} + \frac{1}{S_{\text{HIGH}}} \quad (19)$$

$$S_{\text{LOW}} = A_1 E^{A_2} + A_3 E^{A_4} \quad (20)$$

$$S_{\text{HIGH}} = \frac{A_5 \ln\left(\frac{A_6}{E} + A_7 E\right)}{E^{A_8}} \quad (21)$$

And for the energy rang $10 \text{MeV} \leq E \leq 2 \text{GeV}$ requires to use the following relation:

$$S_e = A_9 + A_{10} \left(\frac{\ln E}{E}\right) + A_{11} \left(\frac{\ln E}{E}\right)^2 + A_{12} \left(\frac{E}{\ln E}\right) \quad (22)$$

And the Coefficiente A_i for every Z_2 is available in the form of TRIM tables[22].

In this research, The following semi experimental relationships were concluded in order to calculate protons stopping power in the compounds C_3H_6 and $C_{22}H_{10}N_2O_5$:-

Table 1

Compound	Semi Experimental Relation	Constants
C_3H_6	$S_e = ae^{bE} + ce^{dE}$	a= 0.7729 b=-3.321 c=0.2381 d=-0.2002
$C_{22}H_{10}N_2O_5$	$S_e = ae^{bE} + ce^{dE}$	a= 0.8226 b=-3.342 c=0.2554 d=-0.2196

6. The Correlation Coefficient

The Program Curve Expert 1.3 was used, the program through which the Correlation Coefficient is calculated and which in concluded to be $(x_i, y_i) i=1, \dots, n$, a sample of the organized pairs that provides the values of two random variables on (n) of the elements. This factor treats the link between the variables x and y and is referred to as $r=r(x, y)$ and which calculated as follows[23]:

- 1) When $r > 0$ then there is a linear link.
- 2) When $r < 0$ then there is a reverse link.
- 3) When $r = 1$ then there is a full extrusive linear link.
- 4) When $r = -1$ then there is a full reverse linear link.
- 5) When $r = 0$ then there is no linear link.
- 6) Whenever the absolute value goes farther than 1, the linear link goes weaker

7. Result and Discussion

The figure(1-a) shows the results of stopping power as a function of energy for protons in C_3H_6 and The figure(1-b) shows the results of stopping power as a function of energy for protons in $C_{22}H_{10}N_2O_5$

Fig (1) shows a great convergence between the variables of Bloch equation of the experimental variables of the SRIM 2012 program except the first variable the kapton compound $C_{22}H_{10}N_2O_5$ of the starting point of the energy range 0.1MeV because of the reduction in the values of the stopping numbers of Bethe according to the equation (13). That affected the negative Bloch correction values which led to the stopping power values of the Bloch equation. And the greatest stopping power at the value 0.18MeV and at the values higher than 0.18MeV, the stopping power are higher than the results of the SRIM 2012 according to Bloch which indicates that irritation and ionization. Then, we notice a very close convergence between the values of the Bloch formula and the experimental values SRIM 2012 until the end of the used range 1000 MeV.

Figure (2) shows the results of stopping power as a function of energy for protons in C_3H_6 and $C_{22}H_{10}N_2O_5$, when performing a comparison between of the tow mediums for the same projectile we would find the stopping power is that of the multe-broblin compound C_3H_6 highest from the Kapton compound $C_{22}H_{10}N_2O_5$, and that it this is because of two factors:

1) Average Ionization Potential (I)

Considering the variety of the average ionization potential gose to the atomic number Z_i for each element of the compound along its percentage n_i in the compound itself. The multi-broblin compound C_3H_6 has a certain ionization potential ($I=56.5429$ eV)and the kapton compound $C_{22}H_{10}N_2O_5$ has a certain ionization potential as well ($I=80.368$ eV), and since it is the denominator, it is inversely proportional with the stopping power.

2) The Fraction $\frac{Z_2}{A_2}$

The fraction includes the atomic and the mass numbers of the element with its percentage in the compound itself (Z_i, A_i ,

n_i) It is for the multi-broblin compound C_3H_6 (0.5714) and the Kapton compound $C_{22}H_{10}N_2O_5$ (0.5131). And, since it is proportional with the stopping power, the stopping power of the multi-broblin C_3H_6 is greater than that of the Kapton compound $C_{22}H_{10}N_2O_5$.

Figure (3) shows the results of Bloch correction as a function of energy for protons in different targets C_3H_6 ($Z_2=24$) and $C_{22}H_{10}N_2O_5$ ($Z_2=196$) which are calculated from eq.(10) at high velocity. In this figure, the Bloch correction of protons is the same in two target and no variation with Z_2 because it is independent on the target and this is mathematically clear as shown in eq.(10). From the figure, the Bloch correction decreases with increasing the energy and becomes quite small and approaches to zero at very high velocity.

Figure (4) shows the percent contribution of Bloch correction to stopping number as a function of energy for protons in different targets C_3H_6 ($Z_2 = 24$) and $C_{10}H_8O_4$ ($Z_2=100$) which are calculated from eq.(10). From the figure, the percent of Bloch correction decreases with increasing the energy and becomes small at high velocity, therefore there is a divergence in values at low velocities and an convergence in values of percent contribution of Bloch correction at high velocities. The percent Bloch correction in C_3H_6 target is larger than that of $C_{10}H_8O_4$ target because the targets are different and for each one a specific atomic number and there are a number of corrections to the stopping number for each target.

Figure (5) the Correlation Coefficient (r) and the error margin(s) were calculated between the semi experimental formula that has been concluded and which is represented in table (1) along with the results of SRIM 2012. From the preceding results, we conclude that there is only one good match according to the Correlation Coefficient.

Figure (6) the Correlation Coefficient (r) and the error margin(s) between the results of Bloch formula were also calculated and are represented by the formula (13) with the results SRIM 2012. From the preceding results, we conclude that there is only one good match according to the Correlation Coefficient.




Figure 1: a. stopping power for protons in (C_3H_6) b. stopping power for protons in ($C_{22}H_{10}N_2O_5$)




Figure 2: stopping power for protons in ($C_{22}H_{10}N_2O_5$) and (C_3H_6)




Figure 3: Bloch correction for protons in $C_{22}H_{10}N_2O_5$ ($Z_2 = 280$) and C_3H_6 ($Z_2 = 24$)





Figure 4: Percent contribution of Bloch correction to stopping number for protons in C_3H_6 ($Z_2 = 24$) and $C_{22}H_{10}N_2O_5$ ($Z_2 = 280$)






Figure 5: a. the Correlation Coefficient and the error margin between the semi experimental formula along with the results of SRIM 2012 in (C_3H_6)
 b. the Correlation Coefficient and the error margin between the semi experimental formula along with the results of SRIM 2012 in ($C_{22}H_{10}N_2O_5$)




Figure 6: a. the Correlation Coefficient and the error margin between the results of Bloch formula with the results SRIM 2012 in (C_3H_6)
 b. the Correlation Coefficient and the error margin between the results of Bloch formula with the results SRIM 2012 in ($C_{22}H_{10}N_2O_5$)

8. Conclusions

The main stopping mechanism is the slowing down due to interactions with target atoms through excitations and ionizations of target electrons via Coulomb scattering. The Bethe theory of calculating stopping power of a point charge penetrating through matter at a non-relativistic velocity can be used over a wide energy interval for fast charged particles based on first Born approximation for atomic collision events which it provides a stopping dependent on the square of projectile charge (Z_1e). Deviations from the (Z_1^2) dependence are especially large at low velocity. For decreasing velocities, high order of (Z_1) terms (Z_1^3 Barkas effect) and (Z_1^4 Bloch correction) dependence in stopping power may become important. Since the accuracy of this scheme deteriorates with increasing projectile charge and decreasing speed, it seemed reasonable to start at the opposite end, i. e. the classical limit (Bohr theory). Bohr pointed that the regime of validity of classical-orbit models and of quantum perturbation theory are roughly complementary. Bloch evaluated the differences between the classical (Bohr) and quantum-mechanical (Bethe) approaches for particles with velocities much larger than the target electrons. He showed that Bohr's harmonic oscillator approach was valid also in the quantum mechanics of a bound electron if the energy transferred was assumed to be the mean energy loss. For light ions (at high velocity and small Bohr's parameter K), the Bloch correction becomes small and usually approaches to zero and the Bloch stopping number (L_{Bloch}) approaches that of Bethe results eq.(4) but at low velocity, the Bethe theory becomes unphysical because of the negative stopping number, therefore the Bloch correction must be added to Bethe stopping number and arrives at the Bohr stopping number eq.(5) and provides the transition to the classical stopping power. Bloch correction depends upon the projectile and its velocity (energy) and are independent of the sign of the projectile charge (Z_1e).

References

- [1] Grande.P. L. and Schiwietz.G, Nucl.Inst.and Meth.In physics Research, B, 195 (2002) 55 .
- [2] Betz.H.D, Nucl.Inst.and Math.132 (1976) .
- [3] Bohr. N, philos. Mag. 25 (1913) 10.
- [4] Bethe .H, Ann. Physic.5 (1930) 325.
- [5] Bloch.F, Ann.phys. 16 (1933) 285.
- [6] Mott. N. F., Proc. R. Soc. London A, 124 (1929) 1856.
- [7] Lindhard .J and Sorensen .A. H, phys. Rev., A.53 (1996) 2443.
- [8] Bichsel.H, Groom. D. E, and Klein. S.R, Phys. Rev., D. 66 (2002) 195.
- [9] Weaver .B. A. and Westphal. A. J, Nucl. Inst. and Meth. In physics Research, B. 187 (2002) 285.
- [10] P. Sigmund, .151 of Springer Series In Solid – State Sciences, .151 (2006).
- [11] Bohr .N, Mat. Fys. Medd. Dan. Vid. Selsk. 18 (1948) 1.
- [12] Sorensen .A. H, Phys. Rev., A . 55 (1997) 2896.
- [13] Hans Bichsel, phys. Rev., A, Vol.41, N.7, pp.3642(1990).
- [14] Ziegler. J.F and Ziegler .M.D, J.P. Biersack, Nucl.Instr.and Meth.in Phys.Res.B.268 (2010) 1818.
- [15] Chaubey.A.K.and Gupta. H.V, Revuede Physique Appliquee To me 12 (1977) 321.

- [16] Northcliffe .L .C and Schilling, R. F, Nuclear Data A
7(1970)233.
- [17] Whaling, W., Hand.Phys.34(1958) 193.
- [18] Anderson, H.H., Hanke, C.C., Sorensen, H. and Vajda,
P., Phys.Rev.153 (1967)338.
- [19] Pierce, T.E.and Blann, M., Phys. Rev.173(1968)390.
- [20] Booth, W.and Grant, I.S., Nucl. phys. 63(1965)481.
- [21] H.H.Andersen and J.F.Ziegler, pergamon press, New
York .2 (1977) .
- [22] M.S.Abosaleh and A.M.Awed, Introduction to Statistics
(2008).

