

Larvicidal and Antagonistic Activities of Crude Leaf Extracts of *Pyrethrum* (*Chrysanthemum*: *Compositae*), *Eucalyptus camaldulensis* Sm. *Myrtaceae*, and *Nicotiana tabacum* (Tobacco L.) (*Solanaceae*) Against Third Instar Larvae of the Malaria Vector, *Anopheles gambiae* s.s. Giles (Diptera: Culicidae).

Glenn O. Araka (Msc, Lead Researcher)¹, John Ochora (PhD)², Johnstone Wakhisi (PhD)³, Dr. Festus Tolo (PhD)⁴, B.O. Khwa-Otsyula⁵

¹Lecturer, Department of Environmental Health, School of Public Health, Moi University, Eldoret, Kenya

²Professor, Department of Botany, School of Biological Sciences, Jomo Kenyatta University of Agriculture and Technology (JKUAT), Nairobi, Kenya;

³Professor, Department of Medical Biochemistry, Moi University, Eldoret, Kenya

⁴Senior Researcher, Kenya Medical Research Institute (KEMRI), Nairobi, Kenya

⁵Professor of Surgery, Department of Surgery and Anesthesiology, School of Medicine, Moi University, Eldoret, Kenya

Abstract: Crude leaf extracts of *Pyrethrum*, *Eucalyptus camaldulensis* and *Nicotiana tabacum* (Tobacco) were individually and in combination tested for their larvicidal and antagonistic activities against third instar larvae of *Anopheles gambiae* s.s. Giles. The combination tests were targeted on antagonistic activities of the crude leaf extracts. Six different solvents were used namely Ethanol, Methanol, Dichloromethane (DCM), Hexane, Ethyl Acetate and Aqueous for the preparation of crude leaf extracts from the plant leaves. The larval mortality of the third instar larvae of *An. gambiae* s.s. Giles was observed after 24 hours of exposure separately in control using 50, 100, 150, 200, 250, 300, 350 and 400 ppm. of crude leaf concentrations. For the individual crude leaf extracts all the six solvent extracts of the plants showed good larvicidal activity. The highest potency was recorded by DCM extract of pyrethrum (LC_{50} 164.68 ppm, LC_{90} 255.17 ppm) achieving 100% mortality of the larvae. Ethanol extract of pyrethrum also exhibited appreciable larvicidal activity at 167.78 ppm depicting 89.70% larval mortality than same extracts of *Nicotiana tabacum* (189.58 ppm) and *E. camaldulensis* (210.15 ppm) causing 75.0% and 78.5% mortality respectively. The rest of the extracts i.e methanol, hexane, ethyl acetate and aqueous exhibited a range of remarkable and varying activities i.e pyrethrum methanol 224.45 ppm, *E. camaldulensis* hexane 198.56 ppm, *N. tabacum* ethyl acetate 201.52 ppm and pyrethrum aqueous 247.84 ppm exhibiting mortality of 79.41%, 73.65%, 85.08% and 77.24% respectively. In combination activities for antagonism on a combination ratio of 1:1 (v.v) for all solvents used in crude leaf extracts (30 treatments) it was observed that 12 combinations exhibited antagonistic activities ($SF < 1$), 15 indicated synergistic activities while 3 combinations were neither antagonistic nor synergistic.

Keywords: Crude extracts, synergistic, antagonistic, solvents, mortality, individual activities, combination activities

1. Introduction

Milugo et al., (2013) describes the antagonistic effect of alkaloids and saponins on bioactivity in quinine tree from tissue samples of quinine tree (*Rauvolfia caffra* Sond) from a remnant forest in Kuria county of Western Kenya (1). Screening for phytochemicals in crude extracts from leaves and stem bark of *R. caffra* revealed the following classes of compounds: alkaloids, terpenoids, saponin, cardiac glycosides and steroids. This confirmed that *R. caffra* contained molecules known to be antioxidant activity. Although crude extracts from bark and leaf samples of *R. caffra* showed antioxidant activity (free radical inhibition) of 79% and 70% respectively analysis of fractions showed activity to vary with phytochemical composition. Fractions

that included saponins, i.e alkaloids, steroids, terpenoids, cardiac glycosides saponins had a lowered activity of 58.99%. Alkaloids only had 63% activity, but fractions containing a combining of alkaloids and saponins exhibited the poorest antioxidant activity of 15%. Alkaloids and saponins appeared to have antagonistic interaction, at least with regards to antioxidant activity. This potentially lowers their activity as antioxidants.

The co-toxicity factor (CTF) calculated as $COF = (O-E/E) \times 100$, where O is observed % mortality and E is expressed as % mortality categorizes result into three: A positive factor of ≥ -20 which indicates potentiation, a negative factor of ≤ -20 indicates antagonism and the intermediate values of > -20 to < 20 indicate an additive effect. By comparing

mortalities obtained with the expected mortality of the mixture (50%) the resultant synergistic/antagonistic factor (SF) could give an indication to the nature of the effect (i.e SF > 1 means synergism, SF < 1 means antagonism; SF=1 means no obvious effect) (2).

Synergy can result when the synergist (i) inhibits the detoxification of a toxin (3, 4, 5, 6), (ii) modifies an inactive compound rendering it toxic (7, 8,9), (iii) enhances the penetration, transport or accessibility of a toxin to its target (10,4,11,12,13,14) or (iv) attacks two independent steps in a process, such as two stages in development or two steps in a biosynthetic pathway (15,16). Antagonism can result when the reverse occurs. For example, an antagonism might modify a toxin rendering it inactive, or decrease the accessibility of a toxin to its target. Diawara et al. (1993) (17) used χ^2 analysis to test for antagonistic effects of chemical combinations on larval mortality. The expected mortality of a chemical combination was determined from the observed mortality of each compound using the formula : $E=Oa + Ob(1-Oa)$. The expected and observed values were analyzed by χ^2 , a significant difference from the expected being interpreted as synergy and other parameters such as larval weight and development time were analysed by ANOVA followed by a second unspecified test to determine which treatments differed and none of the ANOVA results were interpreted in terms of synergy or antagonism.

Greco et al. (1995) (18) have come up with isobolographic analysis method of determining synergistic and antagonistic interactions. In this null model, the combinations of two compounds, A and B, are assumed noninteractive and the compounds A and B are two names for the same compound. Hence, in combination they act in additive manner. Additivity can be further generalized to two compounds that act independently on the same target such that their effects are additive. The combinations of A and B giving the same effect as A or B alone (the isoeffective dose combination) are represented by a straight, dashed line on an isobolograph that connects the effective doses of A and B when alone (Fig. 1). This line is called the zero -interaction isobole (Berenbaum, 1989) (19). The isobolograph (or isobologram) in Figure 1 depicts equivalent activities (GK.iso ζ = same + boli = effect) (Gessner, 1988) (20).




Figure 1: Zero-interaction isobole under the Loewe additivity null model

The dosages for A and B producing the same level of effect are plotted as the intercepts for each axis. The straight,

dashed line represents the isoeffective dose combinations, or the combinations of A and B giving the same effect as either A or B alone.

Further as a hypothetical example, consider two hypothetical dose response curves for compounds A and B (Fig. 2). Use LD₅₀ values (the dose killing 50% of the test organisms) for compounds A and B on the isobolograph, though any constant level of effect (e.g. LD₉₀, LD₂₅) can be used (Greco et al, 1995) (18).




Figure 2: Hypothetical dose response curves for compounds A and B.

On the isobolograph the LD₅₀ values for each compound alone are plotted as the intercepts for each axis (Fig. 3). As in Fig.1 the straight, dashed line between the intercepts in Fig. 3 represents the isobole, or the null model: Loewe additivity with no interaction between the compounds. Thus any combination of A and B having concentrations that fall on this line should cause 50% mortality of the test organisms. If the LD₅₀ value for a particular combination of

A and B lies above the zero- interaction line, A and B at that particular dose combination are said to be antagonistic (Fig.3a). That is they are less potent in combination than one would expect based on their individual effective doses. If the LD₅₀ value lies below the zero-interaction line, A and B are synergistic (Fig.3b). The combination is more effective than A and B are individually. Thus, a dose response curve resulting from various isoeffective combinations of A and B for a synergistic interaction will be concave up and down the zero-interaction line (Fig. 3b). Conversely, an antagonistic interaction will be represented by a concave down curve above the line (Fig. 3a). The null model for the Loewe additivity is also represented by the equation: $d_a/D_a + d_b/D_b = 1$ (Brenhaum, 1989 (19) ;Carter and Genings 1994) (21), where d_a and d_b are the concentrations of A and

B, respectively, used in combination and D_a and D_b are the individual concentrations of A and B producing a specific level of effect (LD₅₀, for example). If the sum of the two ratios equals one, no interaction, or Loewe additivity, exists. If the sum is less than one, synergy is said to occur. If the sum is greater than one, antagonism is said to occur.

As a finality statistics can then be applied in order to determine the intensity of the synergistic or antagonistic interactions. The simplest method for representing the intensity of the interaction , a potency ratio, is illustrated in Figure 4 (Gressner1988) (20).




Figure 3: Isobolograph for compounds A and B




Figure 4: Isobolographic Analysis of a single Experiment

Isobolographic analysis of a single experiment indicating antagonism between indole and caryophylens. The dotted line is the zero interaction isobole constructed from experiments with each compound alone. The solid lines represent 95% confidence intervals. The third point above the zero-interaction isobole is the concentration of the combination that caused 50% mortality along with the 95% confidence intervals. The ratio of the two lines originating from the origin the solid plus the long-dashed line to the solid line is a measure of the intensity of antagonism.

2. Materials and Methods

2.1 Collection of plant materials and extraction of essential oils

A total of three plants were selected for use in these tests and were collected from various sources in Kenya in the month of May 2015. Pyrethrin (1 litre) as crude extract was purchased from Pyrethrum Board of Kenya (PBK), Stanley Mathenge Road, Nakuru, Kenya as a reserve extract. From the same Board 5kg of dried pyrethrum flowers were as well purchased to be able to extract crude oil using similar solvent as those for *Eucalyptus camaldulensis* and *Nicotiana tabaccum*. PBK is a pyrethrum processing and marketing industry located 98 miles (156.8 km) east of Eldoret municipality and similar distance west of Kenya capital city, Nairobi. *Eucalyptus* leaves from mature gum trees (*Eucalyptus camaldulensis*) were collected from Molo sub-county, Kenya a distance of 68 miles (108.8 km) east of Eldoret and 30 miles (48 km) west of Nakuru. Tobacco leaves from the tobacco plant *Nicotiana tabaccum* were purchased from Mr. Meshack Wasike tobacco farm in Malakisi Location, Bungoma county, Kenya, a distance of 88 miles (140.8 km) west of Eldoret municipality and 15 miles (24 km) to the boarder of Kenya and Uganda. Pyrethrum which was purchased from PBK and extracted as crude oil using hexane was stored at 4⁰C in airtight amber or blue bottle until later when required for use. The dried pyrethrum flowers (*Compositae cinerariaefolium* one grown in Kenya) was extracted for pyrethrin mechanically using a commercial stainless steel blender, then 1 kg of powdered leaves was mercerated using six nonpolar to polar solvents:

dichloromethane (DCM), ethyl acetate, ethanol, methanol, hexane and aqueous and similarly were stored under 4°C until required for use. The leaves of *Eucalyptus camaldulensis*, and *Nicotiana tabaccum* (2kg each) were dried in shed for 20-30 days. The dried leaves were then separately powdered mechanically by the same commercial electrical stainless steel blender. One kg of each powdered leaves was extracted successfully by maceration using six nonpolar to polar solvents namely ethanol, methanol, dichloromethane (DCM), ethyl acetate, hexane and water (aqueous). In each solvent the plant material was soaked for 48 hours at 35°C and filtered twice first using a fine cloth and then using Whatman number 1 filter paper (12x15cm) to obtain the extract and to the residue the same solvent was added again. The procedure was repeated twice to obtain maximum extract. The extracts were concentrated at reduced temperature using a rotary vacuum evaporator and stored in air tight amber or blue bottles at 4°C until when required for use. From the stock solutions of the extracts, varying concentrations of each extract were prepared and these concentrations were used for larvicidal bioassays. All chemicals used in this study were of extreme pure grade obtained from Kenya Medical Research Institute (KEMRI), Kisumu, Kenya.

2.2 Mosquito Collection

Larvae of *Anopheles gambiae s.s.* Giles mosquito were grown in a laboratory (insectary) at the Human Anatomy Department, School of Medicine, Moi University, Eldoret. Using a mouth aspirator, male and female adult *Anopheles gambiae s.s.* Giles mosquitoes were collected into test tubes from Langas sub-urban area and taken for rearing in the laboratory. The mosquitoes were placed in cages (30 x 30 x 30cm) in the ratio 3:1 male:female and were fed on 10% sucrose solution soaked in cotton wool. The rearing of larvae adult mosquitoes and larvae were maintained under favourable conditions (temperature 27±2°C, RH 70-80%). Larvae were fed in the laboratory with brewers yeast, dog biscuits and algae (3:1:1) on water surface.

2.3 Larvicidal bioassays

Larvicidal activity of each extract derived from the leaves of Pyrethrum, *Eucalyptus camaldulensis* and *Nicotiana tabaccum* (Tobacco) were tested. The test medium (250ml glass beakers) was prepared by adding 1ml of appropriate dilution of essential oil in ethanol and mixed with 249 ml of distilled water to make up 250ml of test solution (Dhamagadda et al, 2005) (39). Note that as the essential oil does not dissolve in water, it was first dissolved in ethanol (99.0%). From the standard solution varying concentrations of each extract by dilution with distilled water, was prepared in various concentrations of 50, 100, 150, 200, 250, 300, 350 and 400 ppm and these concentrations were used for larvicidal bioassays (An alternative to this method is to take 1gm of the concentrated plant extract and dissolve in 100ml of 1:1 acetate: diethyl sulphoxide (DMSO) and consider as 1% stock solution. From this stock solution varying concentrations as indicated above but expressed as percent could be prepared for use in larvicidal bioassays). Third instar of *Anopheles gambiae s.s.* Giles were exposed to these broad range of test concentrations of each leaf extract to

determine the activity range of each extract. Susceptibility tests were carried out using WHO insecticide susceptibility test-kits (however, slightly modified) and standard procedures (1981) (40). The laboratory reared (27±200C and 75±5% RH) late third instar larvae of *Anopheles gambiae s.s.* Giles were used for experiments. By use of a mouth aspirator batches of 25 late third instar larvae were transferred to 300 ml wide mouth disposable bowls containing serial concentrations of each plant extract. Four replicates were performed for each concentration. Larvae were confirmed dead when they failed to move after probing them with a needle at their cervical region. Moribund larvae were those incapable of rising to the surface when the test solutions were disturbed gently. Moribund larvae were counted after 24 hours of exposure (and added to dead larvae WHO, (2005) (41) and percentage mortality was calculated for each test as follows:-

Number of dead larvae ÷ Number of larvae introduced × 100.

The final percentage was calculated from the average of four replicates. Solutions containing unchlorinated tap water and 1:1 v/v acetone: DMSO but without the plant extract, served as controls. The control mortalities were corrected by using Abbott's formula (1925) (42).

2.4 Statistical Analysis

The average larval mortality data were subjected to Probit analysis for calculating LC₅₀ and LC₉₀ and other statistics at 95% fiducial limits of upper confidence limits (UCL) and lower confidence limit (LCL) and chi-square values were calculated using the SPSS 18.0 (Statistical Package of Social Sciences) software – Finney, (1971) (1).

3. Methodology

3.1 Effects of individual crude leaf extracts.

Three experiments were performed under individual crude leaf extracts activities:

- i) **Experiment 1:** The effect of Pyrethrum crude leaf extract on the third instar larvae of *Anopheles gambiae s.s.* Giles
- ii) **Experiment 2:** The effect of *Eucalyptus camaldulensis* crude leaf extract on the third instar larvae of *Anopheles gambiae s.s.* Giles
- iii) **Experiment 3:** The effect of *Nicotiana tabaccum* crude leaf extract on the third instar of *Anopheles gambiae s.s.* Giles

The effects of Pyrethrum, *E. camaldulensis* and *N. tabaccum* crude leaf extracts on third instar larvae of *An. gambiae s.s.* Giles in the experiments 1, 2 and 3 above were investigated. Six solvents were used for each plant to extract the crude oil resulting to a total of 18 similar treatments for the three plants. Concentrations ranging from 0 ppm (control), 50, 100, 150, 200, 250, 300, 350 and 400 ppm from each plant were made and used. Twenty (20) ml of each concentration for each plant was put into separate 50 ml beakers and then twenty five (25) third instar larvae were dipped into the beakers. Mortality of the larvae was observed after 24 hour exposure by counting the number of dead third instar larvae. Larvae were confirmed dead when

they failed to move after probing them with a needle at their cervical region. Moribund larvae were those incapable of rising to the surface when the test solutions were disturbed gently. Calculations of the percentage of mortality of the larvae from three replications of each treatment determined lethal concentration (LC₅₀).

3.2 The efficacy of various solvent crude leaf extracts combination on the mortality of third instar larvae of *Anopheles gambiae* s.s. Giles

Three experiments were performed under crude leaf extracts combination activities:

3.2.1 Experiment 1: The effect of Pyrethrum crude leaf extract in combination with *Eucalyptus camaldulensis* crude leaf extract on third instar larvae of *Anopheles gambiae* s.s. Giles

3.2.2 Experiment 2: The effect of Pyrethrum crude leaf extract in combination with *N. tabaccum* crude leaf extract on the third instar larvae of *Anopheles gambiae* s.s. Giles

3.2.3 Experiment 3: The effect of *Eucalyptus camaldulensis* crude leaf extract, in combination with *N. tabaccum* crude leaf extract on the third instar larvae of *Anopheles gambiae* s.s. Giles.

The effects of the crude leaf extracts in their combination of Pyrethrum/*E.camaldulensis* (P+E) Pyrethrum/*N. tabaccum* (P+T) and *E. camaldulensis*/*N.tabaccum* (E+T) on third

instar larvae of *An. gambiae* s.s. Giles in the experiments 1, 2, and 3 above were investigated. Six solvents were used for each plant to extract the crude oil. In the crude extract combinations a total of 30 similar treatments for the three plants were carried out. Concentrations ranging from 0 ppm (control), 50,100,150, 200, 250, 300, 350 and 400 ppm from each plant were made and used. Twenty (20) ml of each concentration for each plant was put into separate 50 ml beakers and then twenty five (25) third instar larvae were dipped into the beakers. Mortality of the larvae was observed after 24 hour exposure by counting the number of dead third instar larvae. Larvae were confirmed dead when they failed to move after probing them with a needle at their cervical region. Moribund larvae were those incapable of rising to the surface when the test solutions were disturbed gently. Calculations of the percentage of mortality of the larvae from three replications of each treatment determined lethal concentration (LC₅₀).

4. Results and Discussion

Individual crude leaf extract activities exhibited tremendous results as all solvents achieved far beyond 50% mortality, ranging from pyrethrum DCM crude leaf extract 164.86 ppm (100 % larval mortality), *E. camaldulensis* DCM crude leaf extract 168.65 ppm (100% larval mortality) as highest activities and ethyl acetate 260.56 ppm (65.55% larval mortality). All crude leaf extracts were competitive in their activities.

Table 1: Individual crude leaf extract activities of six various solvents.

Name of plant	Extract solvent	LC ₅₀ (ppm)	Fiducial limits		Regression equation	Chi-square value (x2)
			Upper	Lower		
Pyrethrum	Ethanol	187.78	179.78	196.53	0.7470+1.7758x	4.5217
	Methanol	222.45	209.85	238.71	0.4684+1.9089x	10.6452
	DCM	164.86	161.57	176.28	0.7336+1.7893x	14.2584
	Hexane	230.66	214.79	252.67	0.5566+1.9098x	19.5759
	Ethylacetate	227.56	219.77	269.96	0.4899+1.9078x	19.5759
	Aqueous	247.84	233.37	267.72	0.6758+1.2123x	18.6202
<i>Eucalyptus (Camaldulensis)</i>	Ethanol	210.15	193.88	232.07	0.7686+1.9694x	4.6621
	Methanol	197.46	189.61	208.69	0.4868+1.9227x	13.256
	DCM	168.65	152.44	176.95	0.6975+1.9567x	10.4532
	Hexane	198.56	181.66	220.45	0.7086+1.8685x	9.5033
	Ethylacetate	260.56	240.77	289.96	0.05886+1.7825x	14.0773
	Aqueous	259.58	239.87	288.87	0.05977+1.8365x	6.7556
<i>Nicotiana tabaccum</i>	Ethanol	189.58	181.50	298.42	0.7376+1.8898x	3.5463
	Methanol	224.35	211.73	240.86	0.4988+1.8985x	3.8642
	DCM	229.72	216.80	246.63	0.5878+1.9874x	15.5740
	Hexane	235.85	221.25	240.53	0.5663+1.9096x	4.6542
	Ethylacetate	201.52	191.00	213.84	0.6455+1.8990x	14.0773
	Aqueous	258.42	238.79	287.58	0.0608+2.1015x	6.4444

Antagonistic activities i.e those with SF< 1 are shown in table 2 amongst them synergistic activities and those neither synergistic nor antagonistic. The results indicated antagonistic activities in 12 crude leaf extracts combinations, synergistic activities in 15 crude leaf extract combinations

and 3 combinations were neither synergistic nor antagonistic.

From table 2 antagonistic activities i.e those with SF<1 can be selected and listed as shown in Table 3.

Table 2: Antagonistic activities of the crude leaf extracts combination derived from 30 treatments.

Plant	Ratio	Combination solvent extracts	Individual LC ₅₀ ppm		Combination LC ₅₀ ppm	SF		effect
Pyr. + <i>Eucalyptus</i>	1:1	Ethanol +Aqueous	187.78	210.15	152.85	1.2285	1.3749	S
		Methanol +Ethyl acetate	224.45	260.56	189.54	1.736	1.3747	S
		DCM + Hexane	164.86	168.65	127.85	1.2894	1.3191	S
Pyr + <i>N. tabaccum</i>	1:1	Ethanol +Aqueous	187.78	189.58	146.64	1.2855	1.2928	S
		Methanol +Ethyl acetate	224.45	224.35	232.66	0.9561	0.9643	A
		DCM + Hexane	164.86	229.72	130.78	1.2606	1.7565	S
<i>E. camald.</i> + <i>N. tabaccum</i>	1:1	Ethanol +Aqueous	210.15	258.42	242.80	0.8655	1.0643	SA
		Methanol +Ethyl acetate	197.46	201.52	219.45	0.8998	0.9182	A
		DCM + Hexane	168.65	235.85	135.55	1.2442	1.7399	S
Pyr. + <i>E. camald.</i>	1:1	Methanol + Aqueous	224.45	259.58	201.55	1.1136	1.2879	S
		DCM + Aqueous	164.86	259.58	151.75	1.0864	1.7106	S
		DCM +Methanol	164.86	197.46	118.65	1.3895	1.6642	S
		DCM +Ethyl acetate	164.86	260.56	121.45	1.3574	2.1454	S
		Methanol +Ethyl acetate	224.45	260.56	230.50	0.9651	1.1304	SA
		Methanol + Hexane	197.46	198.56	147.46	1.3391	1.3465	S
		Ethylacetate+ Hexane	227.56	198.56	236.65	0.9616	0.8390	A
Pyr + <i>N. tabaccum</i>	1:1	DCM +Methanol	164.86	224.35	225.80	0.7301	0.9936	A
		DCM +Ethyl acetate	164.86	201.52	160.75	1.0256	1.2536	S
		Methanol + Aqueous	224.45	259.58	263.15	0.8453	0.9864	A
		DCM + Aqueous	164.86	259.58	155.80	1.0582	2.6164	S
		Methanol +Ethyl acetate	224.45	201.52	160.75	1.0256	1.2536	S
		Methanol + Hexane	224.45	235.85	231.65	0.9360	0.9924	A
		Ethylacetate+ Hexane	227.56	235.85	241.45	0.9425	0.9768	A
<i>E. camald.</i> + <i>N. tabaccum</i>	1:1	DCM +Methanol	164.65	224.35	149.35	1.1292	1.5022	S
		DCM +Ethyl acetate	164.65	201.52	211.55	0.7972	0.9526	A
		DCM + Aqueous	164.86	259.58	158.43	1.0406	1.6385	S
		Methanol + Aqueous	222.45	259.58	264.65	0.8483	0.9808	A
		Methanol + Hexane	197.46	235.85	220.75	0.8945	1.0684	SA
		Ethylacetate+ Hexane	260.56	235.85	262.60	0.9922	0.8981	A

Key: S- Synergistic; A – Antagonistic; SA- neither Synergistic nor Antagonistic

Of the five highest synergistic activity to note are the following combinations in their order: pyrethrum DCM extract + *Eucalyptus camaldulensis* methanol extract; pyrethrum DCM extract + *Eucalyptus camaldulensis* ethyl acetate extract; pyrethrum DCM extract + *Eucalyptus camaldulensis* hexane extract; pyrethrum DCM extract + *Nicotiana tabaccum* hexane extract and *Eucalyptus camaldulensis* DCM extract + *Nicotiana tabaccum* hexane extract. All these combinations yielded 100% larval mortality at the concentrations of 118.65, 121.45, 127.85, 130.78 and 135.55 ppm respectively. These were indeed reduced concentrations compared to those used as individual extracts. From table 1 individual extracts activities are low:

pyrethrum DCM (164.86 ppm); *E. camaldulensis* methanol (197.46 ppm); *Eucalyptus camaldulensis* ethyl acetate (260.56 ppm); and *E. camaldulensis* – hexane (198.56 ppm); and *N. tabaccum* hexane (235.85 ppm). This comparison raises the need for combination extracts as they prove to be economical. From the above highest synergistic activities it is observed that DCM extract of pyrethrum is a synergist (as it appears 5 times in the highest 5 activities shown above) and DCM extract of *E. Camaldulensis* is another synergist (as it appears once in the highest 5 activities shown above). Even in the lower activities DCM extract of all the plants is seen to be synergistically useful.

Table 3: Actual antagonistic activities of the crude leaf extracts derived from 30 treatments

Plant	Ratio	Combination solvent extracts	Concentration (LC ₅₀ ppm)		Combination (LC ₅₀ ppm)	SF		Effect
Pyrethrum + <i>N. tabaccum</i>	1:1	Methanol + Ethylacetate	224.45	201.52	232.66	0.9561	0.9645	A
<i>E. camald.</i> + <i>N. tabaccum</i>	1:1	Methanol + Ethylacetate	197.46	201.52	219.45	0.8998	0.9183	A
Pyrethrum + <i>E. camald.</i>	1:1	Ethylacetate +Hexane	227.56	198.56	236.65	0.9616	0.8390	A
Pyrethrum + <i>N. tabaccum</i>	1:1	DCM +Methanol	164.86	224.35	225.80	0.7301	0.996	A
Pyrethrum + <i>N. tabaccum</i>	1:1	Methanol +Ethylacetate	222.45	201.52	243.40	0.9139	0.8279	A
Pyrethrum + <i>N. tabaccum</i>	1:1	Methanol + Hexane	225.45	235.85	237.65	0.9360	0.9924	A
Pyrethrum + <i>N. tabaccum</i>	1:1	Ethylacetate +Hexane	227.56	235.85	241.45	0.9425	0.9768	A
<i>E. camald.</i> + <i>N. tabaccum</i>	1:1	DCM +Ethylacetate	168.65	201.52	211.55	0.7972	0.9526	A
<i>E. camald.</i> + <i>N. tabaccum</i>	1:1	Methanol +Ethylacetate	197.46	201.52	210.30	0.9389	0.9583	A
<i>E. camald.</i> + <i>N. tabaccum</i>	1:1	Ethylacetate +Hexane	260.56	235.85	262.60	0.9922	0.8981	A
Pyrethrum + <i>N. tabaccum</i>	1:1	DCM +Aqueous	222.45	259.58	263.15	0.8453	0.9864	A
<i>E. camald.</i> + <i>N. tabaccum</i>	1:1	Methanol +Aqueous	222.45	258.42	264.65	0.8483	0.9808	A

From these combinations, it was observed that crude leaf extracts with the highest activity (Pyrethrum) are easily

antagonized. From table 3 actual antagonists can be listed as follows:-

Table 4: List of Antagonists

S/NO	Antagonistic crude leaf extract	Lowered concentration (ppm)
1	Methanol of pyrethrum	232.66
2	Ethyl acetate of <i>N. tabaccum</i>	219.45
3	Ethyl acetate of pyrethrum	236.65
4	Methanol of <i>N. tabaccum</i>	225.80
5	Methanol of pyrethrum	243.40
6	Hexane of pyrethrum	237.65
7	Hexane of pyrethrum	241.45
8	Aqueous of <i>N. tabaccum</i>	263.15
9	Ethyl acetate of <i>N. tabaccum</i>	211.55
10	Ethyl acetate of <i>N. tabaccum</i>	210.30
11	Ethylacetate of <i>E. camaldulensis</i>	262.60
12	Aqueous of <i>N. tabaccum</i>	264.65

5. Conclusion

Antagonistic act in opposition of synergistic. Antagonistic exhibit toxification of a toxin (3,4,5,6) demodify an active compound rendering it un toxic (7,8,9), hinders the penetration, transport or accessibility of a toxin to its target (4, 10, 11, 12, 13, 14), and do not attack two independent steps in a process e.g stages in development or biosynthetic pathways (15,16). In this respect antagonists can render mosquito programme control difficult and unsuccessful. It is important one to have prior knowledge on antagonistic activities before any crude leaf extract combinations is attempted to be used in mosquito control strategies. However, there is need for further studies in antagonism to come out clearly which compound(s) in each crude leaf extract is antagonistic to other compound(s) of the combining extract. Antagonistic compounds can render mosquito control programmes expensive since there will be unanticipated purchasing and repeated use of insecticides thus incurring large financial expenditure.

References

- [1] Milugo, T.Z., Omosa, L.K., Ochanda, J.O., Owuor, B.O., Wamunnyokoli, F.A., Oyugi, J.O., and Ochieng, J.W., (2013). Antagonistic in the quinine tree (*Rauvolfia caffra* Sond); further evidence to support biotechnology in traditional medicinal plants. *BMC complimentary and alternative medicine BMC Series open, inclusive and trusted* 2013 13: 285. DOI: 10, 1186/1472-6882-13-285.
- [2] Mansour, S.A., Bakr, R.F.A., Mohamed, R.I., and Hasaneen N.M., (2011). Larvicidal activity of some botanical extracts, commercial insecticides and their binary mixtures against the housefly, *Musca domestica* L. *The Open Toxinology Journal*, 2011, 4, 1-13.
- [3] Wilkinson, C.F., (1976). Insecticide synergism. Pp 195-222 in Metcalf R. McKelvey Jr JJ (eds). *The future for insecticides*. New York: John Wiley and Sons.
- [4] McKey, D., (1979). The distribution of secondary compounds within plants. Pp 55-133 in Rosenthal G.A., and Janzen D.H (eds) *Herbivores: Their Interactions with Secondary Plant Metabolites*. New York: Academic Press.
- [5] Berenbaum, M.R., and Zanger. A.R., (1993). Furanocoumarin metabolism in *Papilio Polyxenes*: biochemistry, genetic variability and ecological significance. *Oecologia* 95:370-375.
- [6] Zangeri A.R. in Ann C. Nelson and Thomas Kursar (1999). Interactions among plant defense compounds: a method for analysis. *Chemoecology* 9:81-92 (1999).
- [7] Spence K.C., (1988) Chemical mediation of coevolution in the *Passiflora-Heliconious* interaction Pp 167-240 in Spencer K.C., (ed) *Chemical Mediation of Coevolution*. San Diego/CA: Academic Press.
- [8] Cipollin, M.L., Stiles, E.W., (1992). Antifungal activity of ripe *Ericaceae* fruits: phenolic – acid interactions and palatability for dispersers. *Biochem Syst Ecol*. 20:501-514.
- [9] Appel, H.M., (1993). Phenolics in ecological interactions: the importance of oxidation. *J Chem Ecol* 19:1521-1552.
- [10] Sun, Y.P., and Johnson, E. R., (1972). Quasi-synergism and penetration of insecticides. *J Econ Entomol* 55:349-353.
- [11] Applebaum, S.W., Birk, Y., (1979). Saponins. Pp. 539-566 in Rosenthal G.A., Janzen, D.H., (eds). *Herbivores: Their Interaction with Secondary Plant Metabolites*, New York: Academic Press.
- [12] Berenbaum, M., (1985). Brentown revisited: interactions among allelochemicals in plants. *Recent Adv Phytochem* 19:139-169.
- [13] Kubo I., and Taniguchi, M., (1988). Polygodial, antifungal potentiator. *J Nat Prod* 51:22-29.
- [14] Vaara, M., (1992). Agents that increase the permeability of the outer membrane. *Microbial Rev* 56:395-411.
- [15] Georghiou, G.P., (1990). Overview of insecticides resistance. Pp. 18-41 in Green M.B., LeBaron, H.M., Moberg W.K., (ed). *Managing Resistance to Agrochemicals*. Washington/DC. American Chemical Society.
- [16] Deminie, C.A, Bechtold, C.M, Stock, D., Alam, M., Djang, F., Balch, A.H., Chou T.C., Prichard, M., Colonna R.J., and Lin P.F., (1996). Evaluation of reverse transcriptase and protease inhibitors in two-drug combinations against human immunodeficiency virus replication. *Antimicrob Agents chemother* 40:1346-1351.
- [17] Diawara, M.M., Trumble J.T., White K.K., Carson, W.G., and Martinez, L.A., (1993). Toxicity of linear furanocoumarins to *Spodoptera exigua*: evidence for antagonistic interactions. *J Chem Ecol* 19: 2473-2484.
- [18] Greco, W.R., Bravo G., and Parsons J.C., (1995). The search for synergy: a critical review from a response surface perspective. *Pharmacol Rev* 47: 331-385.
- [19] Berenbaum, M.C., (1989). What is synergy? *Pharmacol Rev* 41:93-141.
- [20] Gessner, P.K., (1988). A straight forward method for the study of drug interactions: An isobolographic analysis primer. *J An Coll Toxicol* >: 987-1012.
- [21] Carter Jr, W.H., and Genning C., (1994). Analysis of Chemical combinations: relating isobolograms to response surfaces. Pp. 643-653 in Yang R.S.H. (ed) *Toxicology of Chemical Mixtures*. New York: Academic Press.