

Micropropagation through Nodal Culture of an Economically Important Forest Tree Species *Gmelina Arborea* Roxb

M. Rambabu¹, D. Ujjwala², M. Rajinikanth³, N. Rama Swamy⁴

¹Department of Botany, Govt. Degree College, Mahabubabad, Warangal (Dist.) –506101, Telangana, India

²Department of Botany, RMS-Nereda-Kuravi, Warangal (Dist)-506104, Telangana, India

^{3,4}Department of Biotechnology, Kakatiya University, Warangal (Dist.) –506009, Telangana, India

Abstract: *Micropropagation technique plays a vital role in tree improvement program. The species Gmelina arborea (Verbanaceae) is an economically and medicinally important forest tree. In the present investigation, the nodal explants containing single node were cultured on MS medium containing 30 g/L sucrose supplemented with 0.2 - 5.0 mg/L BAP/Kn and also without plant growth regulators. Maximum percentage of response and high frequency number of multiple shoots formation was observed at 0.6mg/L BAP compare with Kn in the specie G.arborea.*

Keywords: micropropagation, *in vitro*, *Gmelina arborea*, auxins, cytokinins, BAP, Kn.

1. Introduction

Micropropagation through proliferation of axillary buds is a common technique in *in vitro* multiplication of forest trees. Axillary bud proliferation approach typically results in many fold increase in shoot number. Each culture passage makes feasible to obtain as many as possible propagules from a single explant and they are true-to-type. The induction of axillary bud proliferation seems to be applicable as a means of micropropagation in many woody forest trees. Proliferation of these axillary buds from tree species may be difficult due to contamination, phenolic exudation and tissue maturity (Bonga, 1982a, b). Rapid clonal propagation has gained considerable importance as a promising tool for multiplication of woody plants. So there are few reports of timber trees of Combretaceae successfully established by tissue culture technology (Joshi *et al.*, 1991). Commercially important forest tree species have been multiplied by using this technique. They are: *Eucalyptus* (Lakshmisita, 1979; Grewal *et al.*, 1980, Gupta *et al.*, 1981); *Melia azaderach* (Raghuraman and Ramanujam, 1998); *Celastrus paniculatus* (Arya *et al.*, 2002); *Maytenus emerginata* (Rathore *et al.*, 1992); *Zizyphus mauritiana* (Sudershan *et al.*, 2000); *Balanites aegyptica*, *Citrus lemon*, *Syzygium cuminii* (Rathore *et al.*, 2004a, b); *Swietenia mahagoni* (Nagarajan *et al.*, 2006).

Gmelina arborea, Roxb. (Vern: Tel. Gummati Teaku) belongs to Verbanaceae is a large to medium sized, fast grown, deciduous tree up to 40m tall and 140cm in diameter. The wood is yellow-grayish or reddish-white with soft and light. It is one of the best timber for making furniture, constructions, plywood, black doors, general carpentry and packages. It is also used in carriages, carvings, musical instruments and ornamental work etc. (Alam *et al.*, 1988). The total plant parts of *G.arborea* were used as a very good medicine. Roots of this species are one of the constituents of “*Dashamula*”, is well known Ayurvedic formulation. Roots and bark are useful in treatment of hallucinations, fever, dyspepsia, stomachalgia and burning

sensation (Tiwari, 1995; Hartwell, 1967-1971; Shirwaiker *et al.*, 2003). Leaf Juice is used for ulcers. Flowers and fruits are used for treating leprosy, anemia, constipation, and also known to inhibit platelet aggregation. The leaves are harvested for fodder for animals and silkworms (Sharma *et al.*, 2001; Duke, 1984; Faiza and Dharakhshanda, 1996, 1998; Little, 1983).

2. Materials Methods

Young and healthy nonwoody branches from two years old plants of *G.arborea* were used as explants. Shoots were excised free of leaves, trimmed into (3cm) segments containing a single node. These explants were washed for 30 minutes under running tap water. Then treated with 5% (w/v) Teepol for 5 minutes, followed by rinsing in distilled water for five times. Surface sterilization was carried out with 0.1% (w/v) HgCl₂ for 5 minutes followed by five rinses in sterile distilled water. These sterilized explants were dried on sterile tissue paper. The nodal segments were trimmed into 1.0-2.0 cm long having one node and were used as explants.

The sterilized nodal segments containing single node were inoculated on MS medium containing 30 g/L sucrose without growth regulators (MSO) and also supplemented with different concentrations (0.2-5.0 mg/L) of BAP/Kn.

After inoculation, all the cultures were incubated under cool white-fluorescent lights at an intensity of 40-60 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for 16 hrs photoperiod at $25 \pm 2^{\circ}\text{C}$.

3. Results

The initial response exhibited by the nodal explants was an enlargement and subsequent breaking of the axillary bud. The time taken for bud breaking was found to be varied depending on the concentrations and combinations of growth regulators used. New shoots were developed after

two weeks of culture and attained height of 2 cm after 3 weeks. Maximum percentage of response (90%) was recorded at 0.6 mg/L BAP followed by 0.8 mg/L (Table-1). Average number of shoots per explant was also recorded at the same (Fig.1). As the concentration of BAP increased, lowest percentage of response was observed. Nodal segments cultured on MS medium fortified with different concentrations of Kn, showed varied results. Maximum percentage (80%) of response was observed at 0.6 mg/L Kn. Lowest percentage of response was recorded at high concentrations of Kn.

4. Discussion

Axillary bud proliferation and elongation was observed in all the concentrations and combinations of growth regulators used except on MS basal medium in *G.arborea*. Early bud breakage and shoot emergence was observed at 0.6 mg/L BAP/Kn. Maximum percentage of response and high frequency number of multiple shoots formation were found at the same concentration of BAP. It was also showed superiority in inducing more number of multiple shoots in comparison to Kn. Similarly Purohit and Kukda (2004) have reported the multiple shoots formation from nodal explants of forest tree *Wrightia tinctoria* on MS medium with 2.0 mg/L BAP.

Anitha and Pullaiah (2002) found that BAP was the best cytokinin in enhancing the shoot buds proliferation in *Sterculia foetida* than to Kn and TDZ. Sunnichan *et al.*, (1998) have recorded the highest frequency number of shoots from axillary bud on MS + 6.62 µm BAP than to Kn in *Sterculia urens*. At higher concentration of BAP and Kn reduced markedly the number of shoots as found in the present investigation. These finding are similar to our present observations that BAP showed superiority over Kn.. Superiority of BAP was reported in many other woody tree species too (Joshi *et al.*, 1991; Gupta *et al.*, 1993; Purohit and Dave, 1996; Sunnichan *et al.*, 1998). Whereas Tiwari *et al.*, (2004) have reported the multiple shoots induction from nodal culture on MS/B₅ medium supplemented with 0.2 mg/L IBA in an endangered forest tree *Pterocarpus marsupium*. Sunnichan *et al.*, (1998) have reported the multiple shoots induction on BAP/Kn in gum-karaya (*Sterculea urens*) and Jagdish Chandra *et al.*, (1999) in *Pisonia alba* as found in *G.arborea*. But Appa Rao (2004) had observed the synergistic effect of both cytokinins BAP and Kn combination by enhancing the shoot buds proliferation in *Sapindus trifoliatus*. Similarly Kathiravan and Ignacimuthu (1999) found the efficiency of both the cytokinins BAP and Kn together in the medium in *Canavalia virosa*.

The size of the nodal explant was also found to play an important role in initiation, proliferation and elongation of

the shoots. Sharon and D'souza (2000) and D'souza and Sharon (2001) and Rama Swamy *et al.*, (2004) have reported that the smaller explants could initiate more multiples than that of longer nodal explants. Similar findings were also noted in *Sapindus trifoliatus* (Appa Rao, 2004). Nodal segment orientation at the time of culture also influences on the proliferation of shoots. Vertical orientation of *Wrightia tinctoria* explants was found better than horizontal orientation in terms of number of proliferated shoots (Purohit *et al.*, 2004) as observed in *G.arborea*.

5. Future Scope

The species *Gmelina arborea* is a very good economic and medicinal important forest tree. This protocol for *in vitro* nodal culture has been established in *G.arborea* which plays a vital role in rapid multiplication of the species

6. Acknowledgement

The corresponding author would like thanks to University Grants Commission for providing financial assistance

Table 1: Effect of BAP/Kn on *in vitro* nodal culture in *G.arborea*

Concentration of PGR (mg/L)	% of response	Average No. of shoots/explant ± (SE) ^a	Mean length of shoot (cm) ± (SE) ^a
M50	b	b	b
BAP			
0.2	75	2.5 ± 0.5	5.0 ± 0.15
0.4	80	3.5 ± 0.10	6.0 ± 0.5
0.6	90	5.0 ± 0.01	6.5 ± 0.5
0.8	84	3.5 ± 0.05	6.0 ± 0.13
1	72	3.0 ± 0.20	5.5 ± 0.10
2	70	2.5 ± 0.02	5.0 ± 0.12
3	65	1.5 ± 0.13*	4.0 ± 0.01
4	60	1.0 ± 0.15*	3.5 ± 0.20
5	52	1.0 ± 0.13*	3.0 ± 0.10
Kn			
0.2	65	2.0 ± 0.10	3.5 ± 0.10
0.4	75	3.0 ± 0.5	4.0 ± 0.12
0.6	80	3.5 ± 0.15	5.0 ± 0.50
0.8	70	2.5 ± 0.10	4.5 ± 0.12
1	65	2.0 ± 0.13	4.0 ± 0.01
2	60	2.0 ± 0.01	3.0 ± 0.05
3	50	1.0 ± 0.15*	2.0 ± 0.20
4	40	1.0 ± 0.10*	1.5 ± 0.15
5	40	1.0 ± 0.02*	1.0 ± 0.13

^aSE = Standard Error; * = With callus induction; b=no response




Figure 2




Figure 2: Nodal explants were cultured on MS medium supplemented with 0.6 mg/L BAP

- single node containing two axillary buds cultured on the medium
- sprouting of axillary buds after two weeks of culture
- multiple shoots formation after four weeks of culture
- elongation of multiple shoots after two months of culture

References

- Alam, M.K., Siddiqui, N.A. and Das, D. (1988). Fodder free of Bangladesh forest research institute, Chittagang, Bangladesh, 167 p.
- Anitha, S. and Pullaih, T. (2002). Shoot regeneration from hypocotyls and shoot tip explants of *Sterculia foetida* L. derived seedlings. *Taiwania*, 47 : 62-69.
- Appa Rao, K. (2004). *In vitro* micropropagation of *Sapindus trifoliatus* L. M.Phil. Thesis, submitted to Kakatiya University, Warangal.
- Arya, V., Singh, R.P. and Shekhawat, N.S. (2002). A micropropagation protocol for mass multiplication and off-site conservation of *Celastrus paniculatus* – A vulnerable medicinal plant of India. *J. Sustainable Forestry*, 14 : 107-120.
- Bonga, J.M. (1982a). Vegetative propagation in relation to juvenility, maturity and rejuvenation. In : *Tissue Culture in Forestry*, (eds.) Bonga, J.M. and Durzan, D.J., Martinus Nijhoff Publications, The Hague, pp.387-412.
- Bonga, J.M. (1982b). *Tissue Culture of economically important plants*, (ed.) Rao, A.N. Costed publication, Singapore, pp.191-196.
- Das, T.D. and Mitra, G.C. (1990). Micropropagation of *Eucalyptus tereticornis* Smith. *Plant Cell Tiss. & Org. Cult.*, 22 : 95-103.
- Duke, J.A. in ed. *An herb a day Ayurvedic Typescript*, (1984).
- D'Souza, M.C. and Sharon, M. (2001). *In vitro* clonal propagation of Annatto (*Bixa orellana* L.). *In vitro Cell Dev. Biol. Plant.*, 37 : 168-172.
- Faiza, H. and Darakhshanda, S. The inhibition of platelet aggregation and related physiological responses with crude drug extract of *Gmelina arborea*, in: Sixth International Symposium on New trends in Natural Products Chemistry, Publication, pp.229-286; (1996, 1998).
- Jagadish Chandra, K.S., Rachappaji, S., Gowda, K.R.D. and Tharasaraswathi, K.J. (1999). *In vitro* propagation of *Pisonia alba* (L.) Spanogar (Lettence tree)-A threatened species. *Phytomorphology*, 49:43-47.

- [12] Joshi, M.S., Shekhawat, N.S. and Rathore, T.S. (1991). Micropropagation of *Anogeissus pendula* Edgew and arid forest tree. *Indian J. Expt. Biol.* 29:615-618.
- [13] Lakshmi Sita, G. and Vaidyanathan, C.S. (1979). Rapid multiplication of *Eucalyptus* by multiple shoot production. *Curr. Sci.*, 48:350-351.
- [14] Little, E.L. Jr. Common fuelwood crops: a handbook for their identification McClain printing Co., Persons. Wv. (1983).
- [15] Grewal, S., Ahuja, A. and Atal, C.K. (1980). *In vitro* proliferation of shoot apices of *Eucalyptus citrioidera* Hook. *Indian J. Exp., Biol.*, 18:775-777.
- [16] Gupta, P.K., Nadgir, A.L., Mascarenhas, A.F. and Jagannathan, V. (1980). Tissue culture of forest trees. Clonal multiplication of *Tectona grandis* (teak) by tissue culture. *Plant Sci. Lett.*, 17:259-268.
- [17] Gupta, P.K., Mascarenhas, A.F. and Hagannathan, V. (1981). Tissue culture of forest trees-clonal propagation of mature trees of *Eucalyptus citrioidera* Hook, by tissue culture. *Plant Sci., Lett.*, 20:195-201.
- [18] Gupta, P.K., Pullman, G., Timmis, R., Kretinger, M., Carison, W.C., Grob, J. and Welty, E. (1993). Forestry in the 21st century – The Biotechnology of somatic embryogenesis. *Biotechnology*, 11:454-459
- [19] Hartwell. J.L. *Plants used against cancer*. A survey Lloydia 30-34;(1967-1971).
- [20] Kathiravan, K. and Ignacimuthu, I. (1999). Micropropagation of *Canavalia virosa* (Roxb.) Wight & Arn. A medicinal plant. *Phytomorphology*, 49:61-66.
- [21] Kukda, G. (1994). *Clonal propagation of toy wood tree Wrightia tinctoria through tissue culture and biotechnology*, Ph.D. Thesis, Mohanlal Sukhadia University, India.
- [22] Nagarajan, S.M., Rajashekhara, S., Kandasamy, S. and Jaychandra, R. (2006). *In vitro* studies on *Swietenia mahagoni* (L.) Jacq. *Asian J. Microbiol. Biotech. Environ. Sci.*, 8: 473-474.
- [23] Purohit, S.D. and Kukda, G. (2004). Micropropagation of an adult tree – *Wrightia tinctoria*. *Indian J. Biotech.*, 3:216-220. Purohit, S.D. and Kukda, G. (2004). Micropropagation of an adult tree – *Wrightia tinctoria*. *Indian J. Biotech.*, 3:216-220.
- [24] Purohit, S.D. and Dave, A. (1996). Micropropagation of *Sterculia urens* Roxb. an endangered tree species. *Plant Cell Rep.*, 15:704-706.
- [25] Raghuraman, G. and Ramanujam, M.P. (1998). Micropropagation of *Melia azadirach* (L.). *J. Swamy Bot. Cl.*, 15:1-5.
- [26] Rama Swamy, N., Ugandhar, T., Praveen, M., Venkataiah, P., Rambabu, M., Upender, M. and Subhash, K. (2005). Somatic embryogenesis and plantlet regeneration from cotyledon and leaf explants of *Solanum surattense*. *Indian J. Biotechnol.* 4: 414-418.
- [27] Rathore, T.S., Singh, R.P. and Shekhawat, N.S. (1991). Clonal propagation of desert teak (*Tecomella undulata*) through tissue culture. *Plant Science*, 79:217-222.
- [28] Rathore, T.S., Deora, N.S. and Shekhawat, N.S. (1992). Cloning of *Maytenus emerginata* (wild) Ding Hou – a tree of the Indian Desert, through tissue culture. *Plant Cell Rep.*, 11:449-451.
- [29] Rathore, J.S., Vinod Rathore, Shekhawat, N.S., Singh, R.P., Liler, G., Mahendra Phulwaria and Dagla, H.R. (2004a). Micropropagation of woody plants. In : *Plant Molecular Biology and Molecular Markers* (eds.) Srivastava, P.S., Alka Narula and Sheela Srivastava, Anamaya Publishers, New Delhi, pp.195-205.
- [30] Rathore, J.S., Shekhawat, N.S., Singh, R.P., Rathore, J.S. and Dagla, H.R. (2004b). Cloning of adult trees of Jamun (*Syzygium cumini*). *Indian J. Biotech.*, 3:241-245.
- [31] Sharon, M. and D'Souza, M. (2000). *In vitro* clonal propagation of annatto (*Bixa orellana* L.). *Curr. Sci.*, 7: 1532-1535.
- [32] Sharma, P.C., Yelne, M.B. and Dennis, T.J. Database on medicinal plants used in Ayurveda, Vol.3, Central Council for Research in Ayurveda and Siddha, Department of ISM and Ministry of health and family welfare. Government of India, pp. 217-228;(2001).
- [33] Shirwaiker., A., Ghosh., S. and Rao, P.G.M. Effect of *Gmelina arborea* Roxb. Leaves on wound healing in rats, *J.Nat. Remedies* 3/1:45-48; (2003).
- [34] Shekhawat, N.S., Rathore, T.S., Singh, R.P., Deora, N. and Rao, S.R. (1993). Factors affecting *in vitro* clonal propagation of *Prosopis cineraria*. *Plant Growth Reg.*, 12:273-280.
- [35] Sudershan, C., Aboel, M.N. and Hussain, J. (2000). *In vitro* propagation of *Zizyphus mauritiana* cultivar umrdn by shoot tip and nodal multiplication. *Curr. Sci.*, 80:290-292.
- [36] Sunnichan, V.G., Shivanna, K.R. and Mohan Ram, H.Y. (1998). Micropropagation of gum karaya (*Sterculia urens*) by adventitious shoot formation and somatic embryogenesis. *Plant Cell Rep.*, 17:951-956.
- [37] Hartwell. J.L. *Plants used against cancer*. A survey Lloydia 30-34;(1967-1971).
- [38] Tiwari, S.K., Kashyap, M.K., Ujjaini, M.M. and Agarwal, A.P. (2002). *In vitro* propagation of *Lagerstromia parviflora* Roxb. from adult tree. *Indian J. Expt. Biol.*, 40: 212-215.
- [39] Tiwari, S., Shah, P. and Singh, K. (2004). *In vitro* propagation of *Pterocarpus marsupium* Roxb. an endangered medicinal tree. *Indian J. Biotech.*, 3: 323-464.